

Omar Hijab

INTRODUCTION TO CALCULUS AND CLASSICAL ANALYSIS



Springer

Undergraduate Texts in Mathematics

Editors

S. Axler

F.W. Gehring

P.R. Halmos

Springer

New York

Berlin

Heidelberg

Barcelona

Budapest

Hong Kong

London

Milan

Paris

Santa Clara

Singapore

Tokyo

Undergraduate Texts in Mathematics

Anglin: Mathematics: A Concise History and Philosophy.

Readings in Mathematics.

Anglin/Lambek: The Heritage of Thales.

Readings in Mathematics.

Apostol: Introduction to Analytic Number Theory. Second edition.

Armstrong: Basic Topology.

Armstrong: Groups and Symmetry.

Axler: Linear Algebra Done Right.

Bak/Newman: Complex Analysis. Second edition.

Banchoff/Wermer: Linear Algebra Through Geometry. Second edition.

Berberian: A First Course in Real Analysis.

Brémaud: An Introduction to Probabilistic Modeling.

Bressoud: Factorization and Primality Testing.

Bressoud: Second Year Calculus. *Readings in Mathematics.*

Brickman: Mathematical Introduction to Linear Programming and Game Theory.

Browder: Mathematical Analysis: An Introduction.

Buskes/van Rooij: Topological Spaces: From Distance to Neighborhood.

Cederberg: A Course in Modern Geometries.

Childs: A Concrete Introduction to Higher Algebra. Second edition.

Chung: Elementary Probability Theory with Stochastic Processes. Third edition.

Cox/Little/O'Shea: Ideals, Varieties, and Algorithms. Second edition.

Croom: Basic Concepts of Algebraic Topology.

Curtis: Linear Algebra: An Introductory Approach. Fourth edition.

Devlin: The Joy of Sets: Fundamentals of Contemporary Set Theory. Second edition.

Dixmier: General Topology.

Driver: Why Math?

Ebbinghaus/Flum/Thomas:

Mathematical Logic. Second edition.

Edgar: Measure, Topology, and Fractal Geometry.

Elaydi: Introduction to Difference Equations.

Exner: An Accompaniment to Higher Mathematics.

Fine/Rosenberger: The Fundamental Theory of Algebra.

Fischer: Intermediate Real Analysis.

Flanigan/Kazdan: Calculus Two: Linear and Nonlinear Functions. Second edition.

Fleming: Functions of Several Variables. Second edition.

Foulds: Combinatorial Optimization for Undergraduates.

Foulds: Optimization Techniques: An Introduction.

Franklin: Methods of Mathematical Economics.

Gordon: Discrete Probability.

Hairer/Wanner: Analysis by Its History. *Readings in Mathematics.*

Halmos: Finite-Dimensional Vector Spaces. Second edition.

Halmos: Naive Set Theory.

Hämmerlin/Hoffmann: Numerical Mathematics.

Readings in Mathematics.

Hijab: Introduction to Calculus and Classical Analysis.

Hilton/Holton/Pedersen: Mathematical Reflections: In a Room with Many Mirrors.

Iooss/Joseph: Elementary Stability and Bifurcation Theory. Second edition.

Isaac: The Pleasures of Probability. *Readings in Mathematics.*

James: Topological and Uniform Spaces.

Jänich: Linear Algebra.

Jänich: Topology.

Kemeny/Snell: Finite Markov Chains.

Kinsey: Topology of Surfaces.

Klambauer: Aspects of Calculus.

Lang: A First Course in Calculus. Fifth edition.

(continued after index)

Omar Hijab

Introduction to Calculus and Classical Analysis

With 68 Illustrations



Springer

Omar Hijab
Department of Mathematics
Temple University
Philadelphia, PA 19122
USA

Editorial Board

S. Axler
Department of
Mathematics
Michigan State University
East Lansing, MI 48824
USA

F.W. Gehring
Department of
Mathematics
University of Michigan
Ann Arbor, MI 48109
USA

P.R. Halmos
Department of
Mathematics
Santa Clara University
Santa Clara, CA 95053
USA

Mathematics Subject Classification (1991): 26-01

Library of Congress Cataloging-in-Publication Data
Hijab, O.

Introduction to calculus and classical analysis / Omar Hijab.

p. cm. — (Undergraduate texts in mathematics)

Includes bibliographical references and index.

ISBN 0-387-94926-7 (hc : alk. paper)

1. Calculus. 2. Mathematical analysis. I. Title. II. Series.

QA303.H54 1997

515'.83—dc21

96-49574

Printed on acid-free paper.

© 1997 Springer-Verlag New York, Inc.

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer-Verlag New York, Inc., 175 Fifth Avenue, New York, NY 10010, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

Production managed by Natalie Johnson; manufacturing supervised by Joe Quatela.

Photocomposed pages prepared from the author's $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$ file.

Printed and bound by R.R. Donnelley and Sons, Harrisonburg, VA.

Printed in the United States of America.

9 8 7 6 5 4 3 2 1

ISBN 0-387-94926-7 Springer-Verlag New York Berlin Heidelberg SPIN 10523597

**This book is dedicated to my daughter
Sonia A. Hijab
who waited patiently for its completion.**

Preface

This is an undergraduate one-variable analysis text. For undergraduate students, the transition from *calculus* to *analysis* is often disorienting and mysterious. What happened to the beautiful calculus formulas? Where did ϵ - δ and open sets come from? It is not until later that one integrates these seemingly diverse points of view. When teaching “advanced calculus”, I always had a difficult time answering these questions.

Now, every mathematician knows that analysis arose naturally in the nineteenth century out of the calculus of the previous two centuries. Believing that it was possible to write a book reflecting, explicitly, this organic growth, I set out to do so.

I chose several of the jewels of classical eighteenth and nineteenth century analysis and inserted them at the end of the book, inserted the axioms for reals at the beginning, and filled in the middle with (and only with) the material necessary for clarity and logical completeness. In the process, every little piece of one-variable calculus assumed its proper place, and theory and application were interwoven throughout.

Let me describe some of the unusual features in this text, as there are other books that adopt the above point of view. First is the systematic avoidance of ϵ - δ arguments. Continuous limits are defined in terms of limits of sequences, limits of sequences are defined in terms of upper and lower limits, and upper and lower limits are defined in terms of \sup and \inf . Everybody thinks in terms of sequences, so why do we teach our undergraduates ϵ 's and δ 's? (In calculus texts, especially, doing this is unconscionable.)

The second feature is the treatment of integration. Since the integral is supposed to be the area under the graph, why not define it that way? What goes wrong? Why don't we define¹ the area of all subsets of \mathbb{R}^2 ? This is the point of view we take in our treatment of integration. As is well known, this approach remains valid, with almost no modifications, in higher dimensions.

The third feature is the treatment of the theorems involving interchange of limits and integrals. Ultimately, all these theorems depend on the *monotone convergence theorem* which, from our point of view, follows from the Greek mathematicians'

¹As in geometric measure theory.

method of exhaustion. Moreover, these limit theorems are stated only after a clear and nontrivial need has been elaborated. For example, *differentiation under the integral sign* is used to compute the area under the graph of the bell-shaped curve.

As a consequence of our treatment of integration, uniform convergence and uniform continuity can be dispensed with. (If the reader has any doubts about this, a glance at the range of applications in Chapter 5 will help.) Nevertheless, we give a careful treatment of uniform continuity, and use it, in the exercises, to discuss an alternative clumsy definition of the integral that was important in the nineteenth century.

The fourth feature is our complete avoidance of complex numbers. We do this to bring out the fundamental simplicity of the material in Chapter 5, which is usually presented in a complex setting.

The fifth feature is our heavy emphasis on computational problems. Computation, here, is often at a deeper level than expected in calculus courses and varies from the high school quadratic formula in §1.4 to $\zeta'(0) = -\log(2\pi)/2$ in §5.8.

Because we take the real numbers as our starting point, basic facts about the natural numbers, trigonometry, or integration are rederived in this context, either in the text or as exercises. Although it is helpful for the reader to have seen calculus prior to reading this text, the development does not presume this. We feel it is important for undergraduates to see, at least once in their four years, a nonpedantic, purely logical development that starts from scratch, is self-contained, and leads to nontrivial and striking results.

We have attempted to present applications from many parts of analysis, many of which do not usually make their way into advanced calculus books. For example we discuss a specific transcendental number, convex conjugates, Machin's formula, the Cantor set, the Bailey–Borwein–Plouffe series, continued fractions, Laplace and Fourier transforms, Bessel functions, Euler's constant, the AGM, the gamma and beta functions, the entropy of the binomial coefficients, infinite products and Bernoulli numbers, theta functions, the zeta function, primes in arithmetic progressions, the Euler–Maclaurin formula, and the Stirling series. Again and again, in discussing these results, we show how the “theory” is indispensable.

As an aid to self-study and assimilation, there are 347 problems with all solutions at the back of the book. If some of the more “theoretical parts” are skipped, this book is suitable for a one-semester course (the extent to which this is possible depends on the students' calculus abilities). Alternatively, covering thoroughly the entire text fills up a year-course, as I have done at Temple teaching our advanced calculus sequence.

Acknowledgements

While writing I benefited from² Courant [5], Lang [9], and Whittaker–Watson [12]. Over and over, I opened Euler’s calculus book (translated into English [7]) to see how *he* would do something, and Birkhoff’s book [2] gave me a feeling for the historical development. The historical remarks at the beginning of Chapter 4 are from *Numbers* [6]. I also looked at Pinsky’s PDE book [10] and the recent advanced calculus book by Strichartz [11].

I learned Euler’s theorem on primes in arithmetic progressions from Knapp’s recent *Notices* article [8], the Euler–Maclaurin formula from a lecture³ of Guillemin at the University of Pennsylvania (Spring 1995), the arithmetic–geometric mean and theta functions from Borwein and Borwein’s book [3], and the Bailey–Borwein–Plouffe series [4] from the worldwide web.

I spent many hours discussing this book, in particular, and calculus issues, in general, with my colleagues. I would like especially to thank Eric Grinberg, Seymour Haber, and Jack Schiller. I am grateful to the students, especially Natalie Boon, in my 1995–96 advanced calculus course, who pointed out errors and made helpful comments. I am also grateful to Temple University and the National Science Foundation (DMS-9121317) for their support.

The text was produced with⁴ the $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$ macro, the figures were drawn with Adobe Illustrator, and the graphs were plotted with Waterloo Software’s Maple.

²References are at the end of the book.

³E. L. Grinberg graciously lent me his notes.

⁴Using Blue Sky Research’s Textures on an Apple Macintosh.

Contents

Preface	vii
Acknowledgements	ix
1 The Set of Real Numbers	1
1.1 Sets and Mappings	1
1.2 The Set \mathbb{R}	3
1.3 The Subset \mathbb{N} and the Principle of Induction	7
1.4 The Completeness Property	12
1.5 Sequences and Limits	16
1.6 Nonnegative Series and Decimal Expansions	25
1.7 Signed Series and Cauchy Sequences	30
2 Continuity	39
2.1 Compactness in \mathbb{R} and \mathbb{R}^2	39
2.2 Continuous Limits	42
2.3 Continuous Functions	45
3 Differentiation	61
3.1 Derivatives	61
3.2 Mapping Properties	68
3.3 Graphing Techniques	74
3.4 Power Series	83
3.5 Trigonometry	94
3.6 Primitives	102
4 Integration	110
4.1 The Cantor Set	110
4.2 Area	114
4.3 The Integral	127
4.4 The Fundamental Theorem of Calculus	143
4.5 The Method of Exhaustion	153
5 Applications	163
5.1 Euler's Gamma Function	163
5.2 The Number π	168
5.3 Gauss' Arithmetic-Geometric Mean (AGM)	182

5.4 The Bell-Shaped Curve	188
5.5 Stirling's Approximation of $n!$	197
5.6 Infinite Products	203
5.7 Jacobi's Theta Functions	211
5.8 Riemann's Zeta Function	217
5.9 The Euler–Maclaurin Formula	226
Appendix Solutions	234
A.1 Chapter 1	234
A.2 Chapter 2	248
A.3 Chapter 3	255
A.4 Chapter 4	269
A.5 Chapter 5	285
References	307
Index	309

1

The Set of Real Numbers

A Note to the Reader

This text consists of many assertions, some big, some small, some almost insignificant. These assertions are obtained from the properties of the real numbers by logical reasoning. Assertions that are especially important are called *theorems*. An assertion's importance is gauged by many factors, including its *depth*, how many other assertions it depends on, its *breadth*, how many other assertions are explained by it, and its level of *symmetry*. The later portions of the text depend on *every single assertion*, no matter how small, made in Chapter 1.

The text is self-contained, and the exercises are arranged linearly: Every exercise can be done using only previous material *from this text*.

Doing the exercises is essential for understanding the material in the text. Some exercises develop additional optional material; these are indicated by a ★.

Sections are numbered linearly within each chapter; for example, §4.3 means the third section in Chapter 4. Equations are numbered linearly within each section; equation numbers are always written within parentheses. Thus, a reference to (3.1) in Chapter 4 is a reference to the first numbered equation in §4.3, whereas a reference to (3.1) in Chapter 2 is a reference to the first numbered equation in §2.3.

Throughout, we use the abbreviation 'iff' to mean 'if and only if' and \square to signal the end of a derivation.

§1.1 Sets and Mappings

We assume the reader is familiar with the usual notions of sets and mappings, but we review them to fix the notation. Strictly speaking, some of the material in this section should logically come after we discuss natural numbers (§1.3). However we include this material here for convenience.

A *set* is a collection A of objects, called *elements*. If x is an element of A we write $x \in A$. If x is not an element of A , we write $x \notin A$. Let A, B be sets. If every element of A is an element of B , we say A is a *subset* of B , and we write

$A \subset B$. Equivalently, we say B is a *superset* of A and we write $B \supset A$. When we write $A \subset B$ or $A \supset B$, we allow for the possibility $A = B$, i.e., $A \subset A$ and $A \supset A$.

The *union* of sets A and B is the set C whose elements lie in A or lie in B ; we write $C = A \cup B$, and we say C equals A union B . The *intersection* of sets A and B is the set C whose elements lie in A and lie in B ; we write $C = A \cap B$ and we say C equals A inter B . Similarly, one defines the union $A_1 \cup \dots \cup A_n$ and the intersection $A_1 \cap \dots \cap A_n$ of finitely many sets A_1, \dots, A_n .

More generally, given any infinite collection of sets A_1, A_2, \dots , their *union* is the set $\bigcup_{n=1}^{\infty} A_n$ whose elements lie in at least one of the given sets. Similarly, their *intersection* $\bigcap_{n=1}^{\infty} A_n$ is the set whose elements lie in all the given sets.

Let A and B be sets. If they have no elements in common, we say they are *disjoint*, $A \cap B$ is *empty*, or $A \cap B = \emptyset$, where \emptyset is the *empty set*, i.e., the set with no elements. By convention, we consider \emptyset a subset of every set.

The set of all elements in A , but not in B , is denoted $A \setminus B = \{x \in A : x \notin B\}$ and is called the *complement of B in A* . For example, when $A \subset B$, the set $A \setminus B$ is empty. Often the set A is understood from the context; in these cases, $A \setminus B$ is denoted B^c and called the *complement of B* .

We will have occasion to use *De Morgan's law*,

$$\begin{aligned} \left(\bigcup_{n=1}^{\infty} A_n \right)^c &= \bigcap_{n=1}^{\infty} A_n^c \\ \left(\bigcap_{n=1}^{\infty} A_n \right)^c &= \bigcup_{n=1}^{\infty} A_n^c. \end{aligned}$$

We leave this as an exercise. Of course these also hold for finitely many sets A_1, \dots, A_n .

If A, B are sets, their *product* is the set $A \times B$ whose elements consist of all ordered pairs (a, b) with $a \in A$ and $b \in B$. A *relation* between two sets A and B is a subset $f \subset A \times B$. A *mapping* is a relation $f \subset A \times B$, such that, for each $a \in A$, there is exactly one $b \in B$ with $(a, b) \in f$. In this case, it is customary to write $b = f(a)$ and $f : A \rightarrow B$.

If $f : A \rightarrow B$ is a mapping, the set A is the *domain*, the set B is the *codomain*, and the set $f(A) = \{f(a) : a \in A\} \subset B$ is the *range*. A *function* is a mapping whose codomain is the set of real numbers \mathbb{R} , i.e., the values of f are real numbers.

A mapping $f : A \rightarrow B$ is *injective* if $f(a) = f(b)$ implies $a = b$, whereas $f : A \rightarrow B$ is *surjective* if every element b of B equals $f(a)$ for some $a \in A$, i.e., if the range equals the codomain. A mapping that is both injective and surjective is *bijective*.

If $f : A \rightarrow B$ and $g : B \rightarrow C$ are mappings, their *composition* is the mapping $g \circ f : A \rightarrow C$ given by $(g \circ f)(a) = g(f(a))$ for all $a \in A$. In general, $g \circ f \neq f \circ g$.

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are mappings, we say they are *inverses* of each other if $g(f(a)) = a$ for all $a \in A$ and $f(g(b)) = b$ for all $b \in B$. A mapping

$f : A \rightarrow B$ is *invertible* if it has an inverse g . It is a fact that a mapping f is invertible iff f is bijective.

Exercises 1.1

1. Give an example where $f \circ g \neq g \circ f$.
2. Verify De Morgan's law.
3. Show that a mapping $f : A \rightarrow B$ is invertible iff it is bijective.
4. Let $f : A \rightarrow B$ be bijective. Show that the inverse $g : B \rightarrow A$ is unique.

§1.2 The Set \mathbf{R}

We are ultimately concerned with one and only one set, the set \mathbf{R} of real numbers. The properties of \mathbf{R} that we use are

- the arithmetic properties,
- the ordering properties, and
- the completeness property.

Throughout, we use 'real' to mean 'real number', i.e., an element of \mathbf{R} .

The *arithmetic properties* start with the fact that reals a, b can be added to produce a real $a + b$, the *sum* of a and b . The rules for addition are $a + b = b + a$ and $a + (b + c) = (a + b) + c$, valid for all reals a, b , and c . There is also a real 0 , called *zero*, satisfying $a + 0 = 0 + a = a$ for all reals a , and each real a has a *negative* $-a$ satisfying $a + (-a) = 0$. As usual, we write subtraction $a + (-b)$ as $a - b$.

Reals a, b can also be multiplied to produce a real $a \cdot b$, the *product* of a and b , also written ab . The rules for multiplication are $ab = ba$, $a(bc) = (ab)c$, valid for all reals a, b , and c . There is also a real 1 , called *one*, satisfying $a1 = 1a = a$ for all reals a , and each real $a \neq 0$ has a *reciprocal* $1/a$ satisfying $a(1/a) = 1$. As usual, we write division $a(1/b)$ as a/b .

Addition and multiplication are related by the property $a(b + c) = ab + ac$ for all reals a, b , and c and the assumption $0 \neq 1$. Let us show how the above properties imply there is a unique real number 0 satisfying $0 + a = a + 0 = a$ for all a . If $0'$ were another real satisfying $0' + a = a + 0' = a$ for all a , then, we would have $0' = 0 + 0' = 0' + 0 = 0$, hence, $0 = 0'$. Also it follows that there is a unique real playing the role of one and $0a = 0$ for all a . These are the arithmetic properties of the reals.

The *ordering properties* start with the fact that there is subset \mathbf{R}^+ of \mathbf{R} , the set of *positive* numbers, that is closed under addition and multiplication, i.e., if $a, b \in \mathbf{R}^+$, then $a + b, ab \in \mathbf{R}^+$. If a is positive, we write $a > 0$ or $0 < a$, and we say a is *greater than* 0 or 0 is *less than* a , respectively. Let \mathbf{R}^- denote the set of *negative* numbers, i.e., $\mathbf{R}^- = -\mathbf{R}^+$ is the set whose elements are the negatives

of the elements of \mathbf{R}^+ . The rules for ordering assume the sets \mathbf{R}^- , $\{0\}$, \mathbf{R}^+ are pairwise disjoint and their union is all of \mathbf{R} . We write $a > b$ and $b < a$ to mean $a - b > 0$. Then, $0 > a$ iff a is negative and $a > b$ implies $a + c > b + c$. These are the ordering properties of the reals.

From the ordering properties, it follows, for example, that $0 < 1$, i.e., one is positive, $a < b$ and $c > 0$ imply $ac < bc$, $0 < a < b$ implies $aa < bb$, and $a < b$, $b < c$ imply $a < c$. As usual, we also write \leq to mean $<$ or $=$, \geq to mean $>$ or $=$, and we say a is *nonnegative* or *nonpositive* if $a \geq 0$ or $a \leq 0$.

If S is a set of reals, a number M is an *upper bound* for S if $x \leq M$ for all $x \in S$. Similarly, m is a *lower bound* for S if $m \leq x$ for all $x \in S$ (Figure 1.1). For example, 1 and $1 + 1$ are upper bounds for the sets $J = \{x : 0 < x < 1\}$ and $I = \{x : 0 \leq x \leq 1\}$ whereas 0 and -1 are lower bounds for these sets. S is *bounded above* (*below*) if it has an upper (lower) bound. S is *bounded* if it is bounded above and bounded below.

Not every set of reals has an upper or a lower bound. Indeed, it is easy to see that \mathbf{R} itself is neither bounded above nor bounded below. A more interesting example is the set \mathbf{N} of natural numbers (next section): \mathbf{N} is not bounded above.




FIGURE 1.1. Upper and lower bounds for A .

A given set S of reals may have several upper bounds. If S has an upper bound M , such that $M \leq b$ for any other upper bound b of S , then, we say M is a *least upper bound* or M is a *supremum* or *sup* for S , and we write $M = \sup S$. Since there cannot be more than one least⁵ upper bound, the sup, whenever it exists, is uniquely determined. For example, in the next section, we see that $\sup I = 1$ and $\sup J = 1$. Similarly, a real m that is a lower bound for S and satisfies $m \geq b$ for all other lower bounds b is called a *greatest lower bound* or an *infimum* or *inf* for S , and we write $m = \inf S$. Again the inf, whenever it exists, is uniquely determined.

The *completeness property* of \mathbf{R} asserts that every nonempty set $S \subset \mathbf{R}$ that is bounded above has a sup, and every nonempty set $S \subset \mathbf{R}$ that is bounded below has an inf.

We introduce a convenient abbreviation, two symbols ∞ , $-\infty$, called *infinity* and *minus infinity*, subject to the ordering rule $-\infty < x < \infty$ for all reals x . If a set S is not bounded above, we write $\sup S = \infty$. If S is not bounded below, we write $\inf S = -\infty$. For example, $\sup \mathbf{R} = \infty$, $\inf \mathbf{R} = -\infty$; in §1.4 we show that $\sup \mathbf{N} = \infty$. Recall that the empty set \emptyset is a subset of \mathbf{R} . Another convenient abbreviation is to write $\sup \emptyset = -\infty$, $\inf \emptyset = \infty$. Clearly, when S is nonempty, $\inf S \leq \sup S$.

⁵If a and b are least upper bounds, then, $a \leq b$ and $a \geq b$.

With this terminology, the completeness property asserts that *every subset of \mathbf{R} , bounded or unbounded, empty or nonempty, has a sup and has an inf; these may be reals or $\pm\infty$.*

We emphasize that ∞ and $-\infty$ are not reals but just convenient abbreviations. As mentioned above, the ordering properties of $\pm\infty$ are $-\infty < x < \infty$ for all real x ; it is convenient to define the following arithmetic properties of $\pm\infty$:

$$\begin{aligned}\infty + \infty &= \infty, \\ -\infty - \infty &= -\infty, \\ \infty - (-\infty) &= \infty, \\ \infty \pm c &= \infty, & c \in \mathbf{R}, \\ -\infty \pm c &= -\infty, & c \in \mathbf{R}, \\ (\pm\infty) \cdot c &= \pm\infty, & c > 0, \\ \infty \cdot \infty &= \infty, \\ \infty \cdot (-\infty) &= -\infty.\end{aligned}$$

Note that we have not defined $\infty - \infty$, $0 \cdot \infty$, ∞/∞ , or $c/0$.

Let a be an upper bound for a set S . If $a \in S$, we say a is a *maximum* of S , and we write $a = \max S$. For example, with I as above, $\max I = 1$. The max of a set S need not exist; for example, in §1.4, we will see that $\max J$ does not.

Similarly, let a be a lower bound for a set S . If $a \in S$, we say a is a *minimum* of S , and we write $a = \min S$. For example, $\min I = 0$ but (§1.4) $\min J$ does not exist.

Theorem. *Let $S \subset \mathbf{R}$ be a set. The max of S and the min of S are uniquely determined whenever they exist. The max of S exists iff the sup of S lies in S , in which case the max equals the sup. The min of S exists iff the inf of S lies in S , in which case the min equals the inf.*

To see this, note that the first statement follows from the second since we already know that the sup and the inf are uniquely determined. To establish the second statement, suppose that $\sup S \in S$. Then, since $\sup S$ is an upper bound for S , $\max S = \sup S$. Conversely, suppose that $\max S$ exists. Then, $\sup S \leq \max S$ since $\max S$ is an upper bound and $\sup S$ is the least such. On the other hand, $\sup S$ is an upper bound for S and $\max S \in S$. Thus, $\max S \leq \sup S$. Combining $\sup S \leq \max S$ and $\sup S \geq \max S$, we obtain $\max S = \sup S$. For the inf, the derivation is completely analogous. \square

Because of this, when $\max S$ exists we say *the sup is attained*. Thus, the sup for I is attained whereas the sup for J is not. Similarly, when $\min S$ exists, we say *the inf is attained*. Thus, the inf for I is attained whereas the inf for J is not.

Let A, B be subsets of \mathbf{R} , let a be real, and let $c > 0$; let $-A = \{-x : x \in A\}$, $A+a = \{x+a : x \in A\}$, $cA = \{cx : x \in A\}$, and $A+B = \{x+y : x \in A, y \in B\}$.

Here are some simple consequences of the definitions that must be checked at this stage:

- $A \subset B$ implies $\sup A \leq \sup B$ and $\inf A \geq \inf B$ (monotonicity property).
- $\sup(-A) = -\inf A$, $\inf(-A) = -\sup A$ (reflection property).
- $\sup(A + a) = \sup A + a$, $\inf(A + a) = \inf A + a$ for $a \in \mathbb{R}$ (translation property).
- $\sup(cA) = c \sup A$, $\inf(cA) = c \inf A$ for $c > 0$ (dilation property).
- $\sup(A + B) = \sup A + \sup B$, $\inf(A + B) = \inf A + \inf B$ (addition property), whenever the sum of the sups and the sum of the infs are defined.

These properties hold whether A and B are bounded or unbounded, empty or nonempty.

We verify the first and the last properties, leaving the others as Exercise 7. For the monotonicity property, if A is empty, the property is immediate since $\sup A = -\infty$ and $\inf A = \infty$. If A is nonempty and $a \in A$, then $a \in B$, hence, $\inf B \leq a \leq \sup B$. Thus, $\sup B$ and $\inf B$ are upper and lower bounds for A , respectively. Since $\sup A$ and $\inf A$ are the least and greatest such, we obtain $\inf B \leq \inf A \leq \sup A \leq \sup B$.

Now, we verify $\sup(A + B) = \sup A + \sup B$. If A is empty, then, so, is $A + B$; in this case, the assertion to be proved reduces to $-\infty + \sup B = -\infty$ which is true (remember we are excluding the case $\infty - \infty$). Similarly, if B is empty.

If A and B are both nonempty, then, $\sup A \geq x$ for all $x \in A$, and $\sup B \geq y$ for all $y \in B$, so, $\sup A + \sup B \geq x + y$ for all $x \in A$ and $y \in B$. Hence, $\sup A + \sup B \geq z$ for all $z \in A + B$, or $\sup A + \sup B$ is an upper bound for $A + B$. Since $\sup(A + B)$ is the least such, we conclude that $\sup A + \sup B \geq \sup(A + B)$. If $\sup(A + B) = \infty$, then, the reverse inequality $\sup A + \sup B \leq \sup(A + B)$ is immediate, yielding the result.

If, however, $\sup(A + B) < \infty$ and $x \in A$, $y \in B$, then, $x + y \in A + B$, hence, $x + y \leq \sup(A + B)$ or, what is the same, $x \leq \sup(A + B) - y$. Thus, $\sup(A + B) - y$ is an upper bound for A ; since $\sup A$ is the least such, we get $\sup A \leq \sup(A + B) - y$. Now, this last inequality implies, first, $\sup A < \infty$ and, second, $y \leq \sup(A + B) - \sup A$ for all $y \in B$. Thus, $\sup(A + B) - \sup A$ is an upper bound for B ; since $\sup B$ is the least such, we conclude that $\sup B \leq \sup(A + B) - \sup A$ or, what is the same, $\sup(A + B) \geq \sup A + \sup B$. Since we already know that $\sup(A + B) \leq \sup A + \sup B$, we obtain $\sup(A + B) = \sup A + \sup B$.

To verify $\inf(A + B) = \inf A + \inf B$, use reflection and what we just finished to write

$$\begin{aligned}\inf(A + B) &= -\sup[-(A + B)] = -\sup[(-A) + (-B)] \\ &= -\sup(-A) - \sup(-B) = \inf A + \inf B.\end{aligned}$$

This completes the derivation of the addition property.

Every assertion that follows in this book depends only on the arithmetic, ordering, and completeness properties of \mathbb{R} , just described.

Exercises 1.2

1. Show that $a0 = 0$ for all real a .
2. Show that there is a unique real playing the role of 1. Also show that each real a has a unique negative $-a$ and each nonzero real a has a unique reciprocal.
3. Show that $-(-a) = a$ and $-a = (-1)a$.
4. Show that negative times positive is negative, negative times negative is positive, and 1 is positive.
5. Show that $a < b$ and $c \in \mathbb{R}$ imply $a + c < b + c$, $a < b$ and $c > 0$ imply $ac < bc$, $a < b$ and $b < c$ imply $a < c$, and $0 < a < b$ implies $aa < bb$.
6. Let $a, b \geq 0$. Show that $a \leq b$ iff $aa \leq bb$.
7. Verify the properties of sup and inf listed above.

§1.3 The Subset N and the Principle of Induction

A subset $S \subset \mathbb{R}$ is *inductive* if

- (1) $1 \in S$ and
- (2) S is closed under addition by 1: $x \in S$ implies $x + 1 \in S$.

For example, \mathbb{R}^+ is inductive. The subset $\mathbb{N} \subset \mathbb{R}$ of *natural numbers* or *naturals* is the intersection of all inductive subsets of \mathbb{R} ,

$$\mathbb{N} = \bigcap \{S : S \subset \mathbb{R} \text{ inductive}\}.$$

Then, \mathbb{N} itself is inductive. Indeed, since $1 \in S$ for every inductive set S , we conclude that $1 \in \bigcap \{S : S \subset \mathbb{R} \text{ inductive}\} = \mathbb{N}$. Similarly, $n \in \mathbb{N}$ implies $n \in S$ for every inductive set S . Hence, $n + 1 \in S$ for every inductive set S . Hence, $n + 1 \in \bigcap \{S : S \subset \mathbb{R} \text{ inductive}\} = \mathbb{N}$. This shows that \mathbb{N} is inductive.

From the definition, we conclude that $\mathbb{N} \subset S$ for any inductive $S \subset \mathbb{R}$. For example, since \mathbb{R}^+ is inductive, we conclude that $\mathbb{N} \subset \mathbb{R}^+$, i.e., every natural is positive.

From the definition, we also conclude that \mathbb{N} is the only inductive subset of \mathbb{N} . For example, $S = \{1\} \cup (\mathbb{N} + 1)$ is a subset of \mathbb{N} , since \mathbb{N} is inductive. Clearly, $1 \in S$. Moreover, $x \in S$ implies $x \in \mathbb{N}$ implies $x + 1 \in \mathbb{N} + 1$ implies $x + 1 \in S$, so, S is inductive. Hence, $S = \mathbb{N}$ or $\{1\} \cup (\mathbb{N} + 1) = \mathbb{N}$, i.e., $n - 1$ is a natural for every natural n other than 1.

The conclusions above are often paraphrased by saying \mathbb{N} is the *smallest inductive subset* of \mathbb{R} , and they are so important they deserve a name.

Principle of Induction. If $S \subset \mathbb{R}$ is inductive, then, $S \supset \mathbb{N}$. If $S \subset \mathbb{N}$ is inductive, then, $S = \mathbb{N}$. \square

Let $2 = 1 + 1 > 1$; we show that there are no naturals between 1 and 2. For this, let $S = \{1\} \cup \{n \in \mathbb{N} : n \geq 2\}$. Then, $1 \in S$. If $n \in S$, there are two possibilities. Either $n = 1$ or $n \neq 1$. If $n = 1$, then, $n + 1 = 2 \in S$. If $n \neq 1$, then, $n \geq 2$, so, $n + 1 > n \geq 2$ and $n + 1 \in \mathbb{N}$, so, $n + 1 \in S$. Hence, S is inductive. Since $S \subset \mathbb{N}$, we conclude that $S = \mathbb{N}$. Thus, $n \geq 1$ for all $n \in \mathbb{N}$, and there are no naturals between 1 and 2. Similarly (Exercise 1), for any $n \in \mathbb{N}$, there are no naturals between n and $n + 1$.

In fact, \mathbb{N} is closed under addition and multiplication by any natural. To see this, fix a natural n , and let $S = \{x : x + n \in \mathbb{N}\}$, so, S is the set of all reals x whose sum with n is natural. Then, $1 \in S$ since $n + 1 \in \mathbb{N}$, and $x \in S$ implies $x + n \in \mathbb{N}$ implies $(x + n) + 1 = (x + 1) + n \in \mathbb{N}$ implies $x + 1 \in S$. Thus, S is inductive. Since \mathbb{N} is the smallest such set, we conclude that $\mathbb{N} \subset S$ or $m + n \in \mathbb{N}$ for all $m \in \mathbb{N}$. Thus, \mathbb{N} is closed under addition. This we write simply as $\mathbb{N} + \mathbb{N} \subset \mathbb{N}$. Closure under multiplication $\mathbb{N} \cdot \mathbb{N} \subset \mathbb{N}$ is similar and left as an exercise.

In the sequel, when we apply the principle of induction, we simply say ‘by induction’.

To show that a given set S is inductive, one needs to verify (1) and (2). Step (2) is often referred to as *the inductive step*, even though, strictly speaking, induction is both (1) and (2), because, usually, most of the work is in establishing (2). Also, the hypothesis in (2), $x \in S$, is often referred to as *the inductive hypothesis*.

Let us give another example of the use of induction. A natural is *even* if it is in $2\mathbb{N} = \{2n : n \in \mathbb{N}\}$. A natural n is *odd* if $n + 1$ is even. We claim that every natural is either even or odd. To see this, let S be the union of the set of even naturals and the set of odd naturals. Then, $2 = 2 \cdot 1$ is even, so, 1 is odd. Hence, $1 \in S$. If $n \in S$ and $n = 2k$ is even, then, $n + 1$ is odd since $(n + 1) + 1 = n + 2 = 2k + 2 = 2(k + 1)$. Hence, $n + 1 \in S$. If $n \in S$ and n is odd, then, $n + 1$ is even, so, $n + 1 \in S$. Hence, in either case, $n \in S$ implies $n + 1 \in S$, i.e., S is closed under addition by 1. Thus, S is inductive. Hence, we conclude that $S = \mathbb{N}$. Thus, every natural is even or odd. Also the usual parity rules hold: even plus even is even, etc.

Let A be a nonempty set. We say A has n elements if there is a bijection between A and the set $\{k \in \mathbb{N} : 1 \leq k \leq n\}$. We often denote this last set by $\{1, 2, \dots, n\}$. If $A = \emptyset$, we say that the number of elements of A is zero. A set A is *finite* if it has n elements for some n . Otherwise, A is *infinite*. Here are some consequences of the definition that are worked out in the exercises. If A and B are disjoint and have n and m elements, respectively, then, $A \cup B$ has $n + m$ elements. If A is a finite subset of \mathbb{R} , then, $\max A$ and $\min A$ exist. In particular, we let $\max(a, b)$, $\min(a, b)$ denote the larger and the smaller of a and b . However, $\max A$ and $\min A$ may exist for an infinite subset of \mathbb{R} .

Theorem. *If $S \subset \mathbb{N}$ is nonempty, then, $\min S$ exists.*

To see this, note that $c = \inf S$ is finite since S is bounded below. Since $c + 1$ is not a lower bound, there is an $n \in S$ with $c \leq n < c + 1$. If $c = n$, then, $c \in S$. Hence, $c = \min S$ and we are done. If $c \neq n$, then, $n - 1 < c < n$, and n is not a

lower bound for S . Hence, $n > 1$, and there is an $m \in S$ lying between $n - 1$ and n . But there are no naturals between $n - 1$ and n . \square

The two other subsets, mentioned frequently, are the *integers* $\mathbf{Z} = \mathbf{N} \cup \{0\} \cup (-\mathbf{N}) = \{0, \pm 1, \pm 2, \dots\}$, and the *rational*s $\mathbf{Q} = \{m/n : m, n \in \mathbf{Z}, n \neq 0\}$. Then, \mathbf{Z} is closed under subtraction (Exercise 3), and \mathbf{Q} is closed under all four arithmetic operations, except under division by zero. As for naturals, we say that the integers in $2\mathbf{Z} = \{2n : n \in \mathbf{Z}\}$ are *even*, and we say that an integer n is *odd* if $n + 1$ is even.

Fix a real a . By (an extension of) induction, one can show (Exercise 9) that there is a function $f : \mathbf{N} \rightarrow \mathbf{R}$ satisfying $f(1) = a$ and $f(n + 1) = af(n)$ for all n . As usual, we write $f(n) = a^n$. Hence, by construction $a^1 = a$ and $a^{n+1} = a^n a$ for all n . Since the set $\{n \in \mathbf{N} : (ab)^n = a^n b^n\}$ is inductive, it follows also that $(ab)^n = a^n b^n$ for $n \in \mathbf{N}$.

Now, $(-1)^n$ is 1 or -1 according to whether $n \in \mathbf{N}$ is even or odd, $a > 0$ implies $a^n > 0$ for $n \in \mathbf{N}$, and $a > 1$ implies $a^n > 1$ for $n \in \mathbf{N}$. These are easily checked by induction.

If $a \neq 0$, we extend the definition of a^n to $n \in \mathbf{Z}$ by setting $a^0 = 1$ and $a^{-n} = 1/a^n$ for $n \in \mathbf{N}$. Then (Exercise 10), $a^{n+m} = a^n a^m$ and $(a^n)^m = a^{nm}$ for all integers n, m .

Let $a > 1$. Then, $a^n = a^m$ with $n, m \in \mathbf{Z}$ only when $n = m$. Indeed, $n - m \in \mathbf{Z}$, and $a^{n-m} = a^n a^{-m} = a^n / a^m = 1$. But $a^k > 1$ for $k \in \mathbf{N}$, and $a^k = 1/a^{-k} < 1$ for $k \in -\mathbf{N}$. Hence, $n - m = 0$ or $n = m$. This shows that powers are unique.

As another application of induction, we establish, simultaneously, the validity of the inequalities $1 < 2^n$ and $n < 2^n$ for all naturals n . This time, we do this without mentioning the set S explicitly, as follows. The inequalities in question are true for $n = 1$ since $1 < 2^1 = 2$. Moreover, if the inequalities $1 < 2^n$ and $n < 2^n$ are true for a particular n (the inductive hypothesis), then, $1 < 2^n < 2^n + 2^n = 2^n 2 = 2^{n+1}$, so, the first inequality is true for $n + 1$. Adding the inequalities valid for n yields $n + 1 < 2^n + 2^n = 2^n 2 = 2^{n+1}$, so, the second inequality is true for $n + 1$. This establishes the inductive step. Hence, by induction, the two inequalities are true for all $n \in \mathbf{N}$. Here, the set S is $S = \{n \in \mathbf{N} : 1 < 2^n, n < 2^n\}$.

Using these inequalities, we show that every nonzero $n \in \mathbf{Z}$ is of the form $2^k p$ for a uniquely determined $k \in \mathbf{N} \cup \{0\}$ and an odd $p \in \mathbf{Z}$. We call k the *number of factors of 2 in n* .

If $2^k p = 2^j q$ with $k > j$ and odd integers p, q , then, $q = 2^{k-j} p = 2 \cdot 2^{k-j-1} p$ is even, a contradiction. On the other hand, if $j > k$, then, p is even. Hence, we must have $k = j$. This establishes the uniqueness of k .

To show the existence of k , by multiplying by a minus, if necessary, we may assume $n \in \mathbf{N}$. If n is odd, we may take $k = 0$ and $p = n$. If n is even, then, $n_1 = n/2$ is a natural $< 2^{n-1}$. If n_1 is odd, we take $k = 1$ and $p = n_1$. If n_1 is even, then, $n_2 = n_1/2$ is a natural $< 2^{n-2}$. If n_2 is odd, we take $k = 2$ and $p = n_2$. If n_2 is even, we continue this procedure by dividing n_2 by 2. Continuing in this manner, we obtain n_1, n_2, \dots naturals with $n_j < 2^{n-j}$. Since this procedure ends in less than n steps, there is some k natural or 0 for which $p = n/2^k$ is odd.

The final issue we take up here concerns square roots. Given a real a , a *square root* of a , denoted \sqrt{a} , is any real x whose square is a , $x^2 = a$. For example 1 has the square roots ± 1 , 0 has the square root 0. On the other hand, not every real has a square root. For example, $\sqrt{-1}$ does not exist within \mathbf{R} , i.e., there is no real x satisfying $x^2 = -1$, since $x^2 + 1 > 0$. In fact, this argument shows that negative numbers never have square roots.

At this point, we do not know whether $\sqrt{2}$ exists within \mathbf{R} . Now, we show that $\sqrt{2}$ does not exist within \mathbf{Q} .

Theorem. *There is no rational a satisfying $a^2 = 2$.*

We argue by contradiction. Suppose that $a = m/n$ is a rational whose square is 2. Then, $(m/n)^2 = 2$ or $m^2 = 2n^2$, i.e., there is a natural N , such that $N = m^2$, $N = 2n^2$. Then, $m = 2^k p$ with odd p and $k \in \mathbf{N} \cup \{0\}$, so, $N = m^2 = 2^{2k} p^2$. Since p^2 is odd, we conclude that $2k$ is the number of factors of 2 in N . Similarly $n = 2^j q$ with odd q and $j \in \mathbf{N} \cup \{0\}$, so, $N = 2n^2 = 2^{2j+1} q^2$. Since q^2 is odd, we conclude that $2j + 1$ is the number of factors of 2 in N . Since $2k \neq 2j + 1$, we arrive at a contradiction. \square

Note that \mathbf{Q} satisfies the arithmetic and ordering properties. The completeness property is all that distinguishes \mathbf{Q} and \mathbf{R} .

As usual, in the following, a *digit* means either 0, 1, 2 or $3 = 2 + 1$, $4 = 3 + 1$, $5 = 4 + 1$, $6 = 5 + 1$, $7 = 6 + 1$, $8 = 7 + 1$, or $9 = 8 + 1$. Also, the letters n , m , i , j will usually denote integers, so, $n \geq 1$ will be used interchangeably with $n \in \mathbf{N}$, with similar remarks for m , i , j .

We say that a nonzero $n \in \mathbf{Z}$ *divides* $m \in \mathbf{Z}$ if $m/n \in \mathbf{Z}$. Alternatively, we say that m is *divisible* by n , and we write $n \mid m$. A natural n is *composite* if $n = jk$ for some $j, k \in \mathbf{N}$ with $j > 1$ and $k > 1$. A natural is *prime* if it is not composite and is not 1. Thus, a natural is prime if it is not divisible by any smaller natural other than 1.

For $a \geq 1$, let $[a] = \max\{n \in \mathbf{N} : n \leq a\}$ denote the *greatest integer* $\leq a$ (Exercises 7 and 8). Then, $[a] \leq a < [a] + 1$, and the *fractional part* of a is $(a) = a - [a]$. Note that the fractional part is a real in $[0, 1)$. More generally, $[a] \in \mathbf{Z}$ and $0 \leq (a) < 1$ are defined⁶ for all $a \in \mathbf{R}$.

Exercises 1.3

1. Let n be a natural. Show that there are no naturals between n and $n + 1$.
2. Show that the product of naturals is natural, $\mathbf{N} \cdot \mathbf{N} \subset \mathbf{N}$.
3. If $m > n$ are naturals, then, $m - n \in \mathbf{N}$. Conclude that \mathbf{Z} is closed under subtraction.
4. Show that no integer is both even and odd. Also, show that even times even is even, even times odd is even, and odd times odd is odd.

⁶ $\{n \in \mathbf{Z} : n \leq a\}$ is nonempty since $\inf \mathbf{Z} = -\infty$ (§1.4).

5. If n, m are naturals and there is a bijection between $\{1, 2, \dots, n\}$ and $\{1, 2, \dots, m\}$, then, $n = m$ (use induction on n). Conclude that the number of elements $\#A$ of a nonempty set A is well defined. Also, show that $\#A = n$, $\#B = m$, and $A \cap B = \emptyset$ imply $\#(A \cup B) = n + m$.
6. If $A \subset \mathbb{R}$ is finite and nonempty, then, show that $\max A$ and $\min A$ exist (use induction).
7. If $S \subset \mathbb{Z}$ is nonempty and bounded above, then, show that S has a max.
8. If $x \geq y > 0$ are reals, then, show that $x = yq + r$ with $q \in \mathbb{N}$, $r \in \mathbb{R}^+ \cup \{0\}$, and $r < y$. (Look at the sup of $\{q \in \mathbb{N} : yq \leq x\}$.)
9. ★ Fix a real a . A set $f \subset \mathbb{R} \times \mathbb{R}$ is *inductive* if $(1, a) \in f$ and $(x, y) \in f$ implies $(x + 1, ay) \in f$. For example, $\mathbb{N} \times \mathbb{R}$ is inductive. Now, let f be the *smallest* inductive set in $\mathbb{R} \times \mathbb{R}$ and let $A = \{x \in \mathbb{R} : (x, y) \in f \text{ for some } y \in \mathbb{R}\}$.

(1) Show that $A = \mathbb{N}$.

(2) Show that f is a mapping with domain \mathbb{N} and codomain \mathbb{R} .

(3) Show that $f(1) = a$ and $f(n + 1) = af(n)$ for all $n \geq 1$.

This establishes the existence of a function $f : \mathbb{N} \rightarrow \mathbb{R}$ satisfying $f(1) = a$ and $f(n + 1) = af(n)$ for all $n \geq 1$. This function is usually denoted $f(n) = a^n$.

10. Let a be a nonzero real. By induction show that $a^n a^m = a^{n+m}$ and $(a^n)^m = a^{nm}$ for all integers n, m .

11. Let $p > 1$ be a natural. Show that for each nonzero $n \in \mathbb{Z}$ there is a unique $k \in \mathbb{N} \cup \{0\}$ and an integer m not divisible by p (i.e., m/p is not in \mathbb{Z}), such that $n = p^k m$.

12. Let $S \subset \mathbb{R}$ satisfy

(1) $1 \in S$ and

(2) $n \in S$ whenever $k \in S$ for all naturals $k < n$.

Show that $S \supset \mathbb{N}$. This is an alternate, and sometimes useful, form of induction.

13. ★ Fix $a > 0$ real, and let $S_a = \{n \in \mathbb{N} : na \in \mathbb{N}\}$. If S_a is nonempty, $m \in S_a$, and $p = \min S_a$, show that p divides m (Exercise 8).

14. ★ Let n, m be naturals and suppose that a prime p divides the product nm . Show that p divides n or m . (Consider $a = n/p$, and show that $\min S_a = 1$ or $\min S_a = p$.)

15. ★ (Fundamental Theorem of Arithmetic) By induction, show that every natural n either is 1 or is a product of primes, $n = p_1 \dots p_r$, with the p_j 's unique except, possibly, for the ordering. (Given n , either n is prime or $n = pm$ for some natural $1 < m < n$; use induction as in Exercise 12.)

16. ★ Given $0 < x < 1$, let $r_0 = x$. Define naturals q_n and remainders r_n by setting

$$\frac{1}{r_n} = q_{n+1} + r_{n+1}, \quad n \geq 0.$$

Thus, $q_{n+1} = [1/r_n]$ and $r_{n+1} = (1/r_n)$, and

$$x = \frac{1}{q_1 + \frac{1}{q_2 + \frac{1}{\ddots \frac{1}{q_{n-1} + \frac{1}{q_n + r_n}}}}}$$

is a *continued fraction*. This algorithm stops the first time $r_n = 0$. Then, the continued fraction is *finite*. If this never happens, this algorithm does not end, and the continued fraction is *infinite*. Show that the algorithm ends iff $x \in \mathbf{Q}$.

§1.4 The Completeness Property

We begin by showing \mathbf{N} has no upper bound. Indeed, if \mathbf{N} has an upper bound, then, \mathbf{N} has a (finite) sup, call it c . Then, c is an upper bound for \mathbf{N} whereas $c - 1$ is not an upper bound for \mathbf{N} , since c is the least such. Thus, there is an $n \geq 1$, satisfying $n > c - 1$, which gives $n + 1 > c$ and $n + 1 \in \mathbf{N}$. But this contradicts the fact that c is an upper bound. Hence, \mathbf{N} is not bounded above. In the notation of §1.2, $\sup \mathbf{N} = \infty$.

Let $S = \{1/n : n \in \mathbf{N}\}$ be the reciprocals of all naturals. Then, S is bounded below by 0, hence, S has an inf. We show that $\inf S = 0$. First, since 0 is a lower bound, by definition of inf, $\inf S \geq 0$. Second, let $c > 0$. Since $\sup \mathbf{N} = \infty$, there is some natural, call it k , satisfying $k > 1/c$. Multiplying this inequality by the positive c/k , we obtain $c > 1/k$. Since $1/k$ is an element of S , this shows that c is not a lower bound for S . Thus, any lower bound for S must be less or equal to 0. Hence, $\inf S = 0$.

The two results just derived are so important we state them again.

Theorem. $\sup \mathbf{N} = \infty$, and $\inf\{1/n : n \in \mathbf{N}\} = 0$. \square

As a consequence, since $\mathbf{Z} \supset \mathbf{N}$, it follows that $\sup \mathbf{Z} = \infty$. Since $\mathbf{Z} \supset (-\mathbf{N})$ and $\inf(A) = -\sup(-A)$, it follows that $\inf \mathbf{Z} \leq \inf(-\mathbf{N}) = -\sup \mathbf{N} = -\infty$, hence, $\inf \mathbf{Z} = -\infty$.

An *interval* is a subset of \mathbf{R} of the following form:

$$(a, b) = \{x : a < x < b\},$$

$$[a, b] = \{x : a \leq x \leq b\},$$

$$[a, b) = \{x : a \leq x < b\},$$

$$(a, b] = \{x : a < x \leq b\}.$$

Intervals of the form (a, b) are *open*, whereas those of the form $[a, b]$ are *compact*. When we use this notation, we are allowing b in (a, b) or $[a, b]$ to equal ∞ and a in (a, b) or $[a, b]$ to equal $-\infty$. Thus, $(a, \infty) = \{x : x > a\}$, $(-\infty, b] = \{x : x \leq b\}$, and $(-\infty, \infty) = \mathbb{R}$.

Let us go back to the open interval $J = (0, 1)$. We already know that 1 is an upper bound for J . If $0 < c < 1$, then, $1 - c > 0$. Since $\inf\{1/n : n \in \mathbb{N}\} = 0$, there is a natural n satisfying $0 < 1/n < 1 - c$ which implies $c < c + 1/n < 1$. Hence, $c + 1/n$ is in J and is greater than c , so, c is not an upper bound for J . We conclude that $\sup J = 1$ and $\max J$ does not exist (since, by definition, the max must lie in the set). Similarly, $\inf J = 0$, and $\min J$ does not exist.

For $x \in \mathbb{R}$, we define $|x|$, the *absolute value* of x , by

$$|x| = \max(x, -x).$$

Then, $x \leq |x|$ for all x , and, for $a > 0$, $\{x : -a < x < a\} = \{x : |x| < a\} = \{x : x < a\} \cap \{x : x > -a\}$, $\{x : x < -a\} \cup \{x : x > a\} = \{x : |x| > a\}$.

The absolute value satisfies the following properties:

- (1) $|x| > 0$ for all nonzero x , and $|0| = 0$,
- (2) $|x||y| = |xy|$ for all x, y ,
- (3) $|x + y| \leq |x| + |y|$ for all x, y .

We leave the first two as exercises. The third, the *triangle inequality*, is derived using $|x|^2 = x^2$ as follows:

$$\begin{aligned} |x + y|^2 &= (x + y)^2 = x^2 + 2xy + y^2 \\ &\leq |x|^2 + 2|xy| + |y|^2 = |x|^2 + 2|x||y| + |y|^2 = (|x| + |y|)^2. \end{aligned}$$

Since $a \leq b$ iff $a^2 \leq b^2$ for a, b nonnegative (Exercise 6 of §1.2), the triangle inequality is established.

Frequently, the triangle inequality is used in alternate forms, one of which is

$$|x - y| \geq |x| - |y|.$$

This follows by writing $|x| = |(x - y) + y| \leq |x - y| + |y|$ and transposing $|y|$ to the other side. Another form is

$$|a_1 + a_2 + \cdots + a_n| \leq |a_1| + |a_2| + \cdots + |a_n|, \quad n \geq 1.$$

We show how the completeness property can be used to derive the existence of $\sqrt{2}$ within \mathbb{R} .

Theorem. *There is a real a satisfying $a^2 = 2$.*

To see this, let $S = \{x : x \geq 1 \text{ and } x^2 < 2\}$. Since $1 \in S$, S is nonempty. Also, $x \in S$ implies $x = x1 \leq xx = x^2 < 2$, hence, S is bounded above by 2, hence, S has a sup, call it a . We claim that $a^2 = 2$. We establish this claim by ruling out the cases $a^2 < 2$ and $a^2 > 2$, leaving us with the desired conclusion (remember every real is positive or negative or zero).

So, suppose that $a^2 < 2$. If we find a natural n with

$$\left(a + \frac{1}{n}\right)^2 < 2,$$

then, $a + 1/n \in S$, hence, the real a could not have been an upper bound for S , much less the least such. To see how to find such an n , note that

$$\begin{aligned} \left(a + \frac{1}{n}\right)^2 &= a^2 + \frac{2a}{n} + \frac{1}{n^2} \\ &\leq a^2 + \frac{2a}{n} + \frac{1}{n} = a^2 + \frac{2a+1}{n} < 2 \end{aligned}$$

if $(2a+1)/n < 2 - a^2$, i.e., if $n > (2a+1)/(2 - a^2)$. Since $a^2 < 2$, $b = (2a+1)/(2 - a^2)$ is a perfectly well defined positive real. Since $\sup \mathbb{N} = \infty$, such a natural $n > b$ can always be found. This rules out $a^2 < 2$.

Before we rule out $a^2 > 2$, we note that S is bounded above by any positive b satisfying $b^2 > 2$ since, for b and x positive, $b^2 > x^2$ iff $b > x$.

Now suppose that $a^2 > 2$. Then, $b = (a^2 - 2)/2a$ is positive, hence, there is a natural n satisfying $1/n < b$ which implies $a^2 - 2a/n > 2$. Hence,

$$\left(a - \frac{1}{n}\right)^2 = a^2 - \frac{2a}{n} + \frac{1}{n^2} > 2,$$

so, $a - 1/n$ is an upper bound for S . This shows that a is not the *least* upper bound, contradicting the definition of a . Thus, we are forced to conclude that $a^2 = 2$. \square

A real a satisfying $a^2 = 2$ is called a *square root* of 2. Since $(-x)^2 = x^2$, there are two square roots of 2, one positive and one negative. From now on, the positive square root is denoted $\sqrt{2}$. Similarly, every positive a has a positive square root, which we denote \sqrt{a} .

More generally, for every $b > 0$ and $n \geq 1$, there is a unique $a > 0$ satisfying $a^n = b$, the *n th root* $a = b^{1/n}$ of b . Now, for $n \geq 1$, $k \geq 1$, and $m \in \mathbb{Z}$,

$$[(b^m)^{1/n}]^{nk} = \left\{ [(b^m)^{1/n}]^n \right\}^k = (b^m)^k = b^{mk},$$

hence, by uniqueness of roots, $(b^m)^{1/n} = (b^{mk})^{1/nk}$. Thus, for $r = m/n$ rational, we may set $b^r = (b^m)^{1/n}$, defining rational powers of positive reals.

Since $\sqrt{2} \notin \mathbb{Q}$, $\mathbb{R} \setminus \mathbb{Q}$ is not empty. The reals in $\mathbb{R} \setminus \mathbb{Q}$ are the *irrationals*. In fact, both the rationals and the irrationals have an interlacing or *density* property.

Theorem. *If $a < b$ are any two reals, there is a rational s between them, $a < s < b$, and there is an irrational t between them, $a < t < b$.*

To see this, first, choose a natural n satisfying $1/n < b - a$. Second, let $S = \{m \in \mathbb{N} : na < m\}$, and let $k = \inf S = \min S$. Since $k \in S$, $na < k$. Since

$k - 1 \notin S, k - 1 \leq na$. Hence, $s = k/n$ satisfies

$$a < s \leq a + \frac{1}{n} < b.$$

For the second assertion, choose a natural n satisfying $1/n\sqrt{2} < b - a$, let $T = \{m \in \mathbb{N} : \sqrt{2}na < m\}$, and let $k = \min T$. Since $k \in T, k > \sqrt{2}na$. Since $k - 1 \notin T, k - 1 \leq \sqrt{2}na$. Hence, $t = k/(n\sqrt{2})$ satisfies

$$a < t \leq a + \frac{1}{n\sqrt{2}} < b.$$

Moreover, t is necessarily irrational. \square

Approximation of reals by rationals is discussed further in the exercises.

Exercises 1.4

1. Show that $x \leq |x|$ for all x and, for $a > 0$, $\{x : -a < x < a\} = \{x : |x| < a\} = \{x : x < a\} \cap \{x : x > -a\}$, $\{x : x < -a\} \cup \{x : x > a\} = \{x : |x| > a\}$.
2. For all $x \in \mathbb{R}$, $|x| \geq 0$, $|x| > 0$ if $x \neq 0$, and $|x||y| = |xy|$ for all $x, y \in \mathbb{R}$.
3. By induction, show that $|a_1 + a_2 + \cdots + a_n| \leq |a_1| + |a_2| + \cdots + |a_n|$ for $n \geq 1$.
4. Show that every $a > 0$ has a unique positive square root.
5. Show that $ax^2 + bx + c = 0, a \neq 0$, has two, one, or no solutions in \mathbb{R} according to whether $b^2 - 4ac$ is positive, zero, or negative. When there are solutions, they are given by $x = (-b \pm \sqrt{b^2 - 4ac})/2a$.
6. By induction, show that $(1 + a)^n \leq 1 + (2^n - 1)a$ for $n \geq 1$ and $0 \leq a \leq 1$. Also show that $(1 + a)^n \geq 1 + na$ for $n \geq 1$ and $a \geq -1$.
7. For $a, b \geq 0$, show that $a^n \geq b^n$ iff $a \geq b$. Also show that every $b > 0$ has a unique positive n th root for all $n \geq 1$ (use Exercise 6 and modify the derivation for $\sqrt{2}$).
8. Show that the real t constructed in the derivation of the last theorem is irrational.
9. ★ Let a be any real. Show that, for each $\epsilon > 0$, no matter how small, there are integers $n \neq 0, m$ satisfying

$$\left|a - \frac{m}{n}\right| < \frac{\epsilon}{n}.$$

(Let (x) denote the fractional part of x , consider the sequence $(a), (2a), (3a), \dots$, and divide $[0, 1]$ into finitely many subintervals of length less than ϵ . Since there are infinitely many terms in the sequence, at least 2 of them must lie in the same subinterval.)

10. ★ Show that $a = \sqrt{2}$ satisfies

$$\left|a - \frac{m}{n}\right| \geq \frac{1}{(2\sqrt{2} + 1)n^2}, \quad n, m \geq 1.$$

(Consider the two cases $|a - m/n| \geq 1$ and $|a - m/n| \leq 1$, separately, and look at the minimum of $n^2|f(m/n)|$ with $f(x) = x^2 - 2$.)

11. ★ Let $a = \sqrt{1 + \sqrt{2}}$. Then, a is irrational, and there is a positive real c satisfying

$$\left|a - \frac{m}{n}\right| \geq \frac{c}{n^4}, \quad n, m \geq 1.$$

(Factor $f(a) = a^4 - 2a^2 - 1 = 0$, and proceed as in the previous exercise.)

12. ★ For $n \in \mathbb{Z} \setminus \{0\}$, define $|n|_2 = 1/2^k$ where k is the number of factors of 2 in n . Also define $|0|_2 = 0$. For $n/m \in \mathbb{Q}$ define $|n/m|_2 = |n|_2/|m|_2$. Show that $|\cdot|_2 : \mathbb{Q} \rightarrow \mathbb{R}$ is well defined and satisfies the absolute value properties (1), (2), and (3).

§1.5 Sequences and Limits

A *sequence*⁷ of real numbers is a function $f : \mathbb{N} \rightarrow \mathbb{R}$. Usually, we write a sequence as (a_n) where $a_n = f(n)$ is the n th term. For example, the formulas $a_n = n$, $b_n = 2n$, $c_n = 2^n$, and $d_n = 2^{-n} + 5n$ yield sequences (a_n) , (b_n) , (c_n) , and (d_n) . Later, we will consider sequences of sets (Q_n) and sequences of functions (f_n) , but now we discuss only sequences of reals.

It is important to distinguish between the *sequence* (a_n) (the function f) and the *set* $\{a_n\}$ (the range $f(\mathbb{N})$ of f). In fact, a sequence is an *ordered* set (a_1, a_2, a_3, \dots) and not just a set $\{a_1, a_2, a_3, \dots\}$. Sometimes it is more convenient to start sequences from the index $n = 0$, i.e., to consider a sequence as a function on $\mathbb{N} \cup \{0\}$. For example, the sequence $(1, 2, 4, 8, \dots)$ can be written $a_n = 2^n$, $n \geq 0$. Specific examples of sequences are usually constructed by induction as in Exercise 9 of §1.3. However, we will not repeat the construction carried out there for each sequence we encounter.

In this section, we are interested in the behavior of sequences as the index n increases without bound. Often this is referred to as the “limiting behavior” of sequences. For example, consider the sequences

$$(a_n) = (1/2, 2/3, 3/4, 4/5, \dots),$$

$$(b_n) = (1, -1, 1, -1, \dots),$$

$$(c_n) = \left(2, \sqrt{2}, \sqrt{\sqrt{2}}, \sqrt{\sqrt{\sqrt{2}}}, \dots\right),$$

$$(d_n) = (2, 3/2, 17/12, 577/408, \dots),$$

⁷This notion makes sense for finite sets also: A *finite sequence* (a_1, \dots, a_n) of reals is a function $f : \{1, \dots, n\} \rightarrow \mathbb{R}$.

where, in the last⁸ sequence, $d_1 = 2$, $d_2 = (d_1 + 2/d_1)/2$, $d_3 = (d_2 + 2/d_2)/2$, $d_4 = (d_3 + 2/d_3)/2$, and so on. What are the limiting behaviors of these sequences?

As n increases, the terms in (a_n) are arranged in increasing order, and $a_n \leq 1$ for all $n \geq 1$. However, if we increase n sufficiently, the terms $a_n = (n-1)/n = 1 - 1/n$ become arbitrarily close to 1, since $\sup\{1 - 1/n : n \geq 1\} = 1$ (§1.4). Thus, it seems reasonable to say that (a_n) approaches one or the limit of the sequence (a_n) equals one.

On the other hand, the sequence (b_n) does not seem to approach any single real, as it flips back and forth between 1 and -1 . Indeed, one is tempted to say that (b_n) has two limits, 1 and -1 .

The third sequence is more subtle. Since we have $\sqrt{x} < x$ for $x > 1$, the terms are arranged in decreasing order. Because of this it seems reasonable that (c_n) approaches its “bottom”, i.e., (c_n) approaches $L = \inf\{c_n : n \geq 1\}$. Although, in fact, this turns out to be so, it is not immediately clear just what L equals.

The limiting behavior of the fourth sequence is not at all clear. If one computes the first nine terms, it is clear that this sequence approaches something quickly. However, since such a computation is approximate, at the outset, we cannot be sure there is a single real number that qualifies as “the limit” of (d_n) . The sequence (d_n) is discussed in Exercise 12 of this section and in Exercise 5 of §1.6.

It is important to realize that

- What does “limit” mean?
- Does the limit exist?
- How do we compute the limit?

are very different questions. When the situation is sufficiently simple, say, as in (a_n) or (b_n) above, we may feel that the notion of “limit” is self-evident and needs no elaboration. Then, we may choose to deal with more complicated situations on a case-by-case basis and not worry about a “general” definition of limit. Historically, however, mathematicians have run into trouble using this ad hoc approach. Because of this, a more systematic approach was adopted in which a single definition of “limit” is applied. This approach was so successful that it is universally followed today.

Below, we define the concept of limit in two stages, first, for monotone sequences and, then, for general sequences. To deal with situations where sequences have more than one limit, the auxiliary concept of a “limit point” is introduced in Exercise 9. Now, we turn to the formal development.

Let (a_n) be any sequence. We say (a_n) is *decreasing* if $a_n \geq a_{n+1}$ for all natural n . If $L = \inf\{a_n : n \geq 1\}$, in this case, we say (a_n) *approaches* L as $n \nearrow \infty$, and we write $a_n \searrow L$ as $n \nearrow \infty$. Usually, we drop the phrase ‘as $n \nearrow \infty$ ’ and simply write $a_n \searrow L$. We say a sequence (a_n) is *increasing* if $a_n \leq a_{n+1}$ for all $n \geq 1$. If $L = \sup\{a_n : n \geq 1\}$, in this case, we say (a_n) *approaches* L as $n \nearrow \infty$, and we write $a_n \nearrow L$ as $n \nearrow \infty$. Usually, we drop the phrase ‘as $n \nearrow \infty$ ’ and simply write $a_n \nearrow L$. Alternatively, in either case, we say *the limit of (a_n) is L* , and we

⁸Decimal notation, e.g., $17 = 7 + (9 + 1)$, is reviewed in the next section.

write

$$\lim_{n \nearrow \infty} a_n = L.$$

Note that since sups and infs are uniquely determined, we say ‘the limit’ instead of ‘a limit’. Thus,

$$\lim_{n \nearrow \infty} \left(1 - \frac{1}{n}\right) = 1$$

since $\sup\{1 - 1/n : n \geq 1\} = 1$,

$$\lim_{n \nearrow \infty} \frac{1}{n} = 0$$

since $\inf\{1/n : n \geq 1\} = 0$, and

$$\lim_{n \nearrow \infty} n^2 = \infty$$

since $\sup\{n^2 : n \geq 1\} = \infty$.

We say a sequence is *monotone* if the sequence is either increasing or decreasing. Thus the concept of *limit* is now defined for every monotone sequence.

If (a_n) is a monotone sequence approaching a nonzero limit a , then, there is a natural N beyond which $a_n \neq 0$ for $n \geq N$. To see this, suppose that (a_n) is increasing and $a > 0$. Then, by definition $a = \sup\{a_n : n \geq 1\}$, hence, $a/2$ is not an upper bound for (a_n) . Thus, there is a natural N with $a_N > a/2 > 0$. Since the sequence is increasing, we conclude that $a_n \geq a_N > 0$ for $n \geq N$. If (a_n) is increasing and $a < 0$, then, $a_n \leq a < 0$ for all $n \geq 1$. If the sequence is decreasing, the reasoning is similar.

Before we define limits for arbitrary sequences, we show that every sequence (a_n) lies between a decreasing sequence (a_n^*) and an increasing sequence (a_{n*}) in a simple and systematic fashion.

Let (a_n) be any sequence. Let $a_1^* = \sup\{a_k : k \geq 1\}$, $a_2^* = \sup\{a_k : k \geq 2\}$, and, for each natural n , let $a_n^* = \sup\{a_k : k \geq n\}$. Thus, a_n^* is the sup of all the terms starting from the n th term. Since $\{a_k : k \geq n+1\} \subset \{a_k : k \geq n\}$ and the sup is monotone (§1.2), $a_{n+1}^* \leq a_n^*$. Thus, (a_n^*) is decreasing and $a_n \leq a_n^*$ since $a_n \in \{a_k : k \geq n\}$. Similarly, we set $a_{n*} = \inf\{a_k : k \geq n\}$ for each $n \geq 1$. Then, (a_{n*}) is increasing and $a_n \geq a_{n*}$. (a_n^*) is the *upper sequence*, and (a_{n*}) is the *lower sequence* of the sequence (a_n) (Figure 1.2).




FIGURE 1.2. Upper and lower sequences with $x_n = x_6$, $n \geq 6$.

If the sequence (a_n) is any sequence, then, the sequences (a_n^*) , (a_{n*}) are monotone, hence, they have limits,

$$a_n^* \searrow a^*, \quad a_{n*} \nearrow a_*.$$

In fact, $a_* \leq a^*$. To see this, fix a natural $N \geq 1$. Then,

$$a_{N*} \leq a_{n*} \leq a_n \leq a_n^* \leq a_N^*, \quad n \geq N.$$

But since (a_{n*}) is increasing, $a_{1*}, a_{2*}, \dots, a_{N*}$ are all $\leq a_{N*}$, hence,

$$a_{n*} \leq a_N^*, \quad n \geq 1.$$

Hence, a_N^* is an upper bound for the set $\{a_{n*} : n \geq 1\}$. Since a_* is the sup of this set, we must have $a_* \leq a_N^*$. But this is true for every natural N . Since a^* is the inf of the set $\{a_N^* : N \geq 1\}$, we conclude that $a_* \leq a^*$.

Theorem. Let (a_n) be a sequence, and let

$$a_n^* = \sup\{a_k : k \geq n\}, \quad a_{n*} = \inf\{a_k : k \geq n\}, \quad n \geq 1.$$

Then, (a_n^*) and (a_{n*}) are decreasing and increasing, respectively. Moreover, if a^* and a_* are their limits, then,

- $a_{n*} \leq a_n \leq a_n^*$ for all $n \geq 1$,
- $a_n^* \searrow a^*$,
- $a_{n*} \nearrow a_*$, and
- $-\infty \leq a_* \leq a^* \leq \infty$.

A sequence (a_n) is *bounded* if $\{a_k : k \geq 1\}$ is a bounded subset of \mathbf{R} . Otherwise, (a_n) is *unbounded*. We caution the reader that some of the terms a_n^*, a_{n*} as well as the limits a^*, a_* , may equal $\pm\infty$, when (a_n) is unbounded. Keeping this possibility in mind, the theorem is correct as it stands. \square

If the sequence (a_n) happens to be increasing, then, $a_n^* = a^*$ and $a_{n*} = a_n$ for all $n \geq 1$. If (a_n) happens to be decreasing, then, $a_n^* = a_n$ and $a_{n*} = a_*$ for all $n \geq 1$.

If N is a fixed natural and (a_n) is a sequence, let (a_{N+n}) be the sequence $(a_{N+1}, a_{N+2}, \dots)$. Then, $a_n \nearrow a_*$ iff $a_{N+n} \nearrow a_*$, and $a_n \searrow a^*$ iff $a_{N+n} \searrow a^*$. Also, by the sup reflection property (§1.2), $b_n = -a_n$ for all $n \geq 1$ implies $b_n^* = -a_{n*}$, $b_{n*} = -a_n^*$ for all $n \geq 1$. Hence, $b^* = -a_*$, $b_* = -a^*$.

Now we define the limit of an arbitrary sequence. Let (a_n) be any sequence, and let (a_n^*) , (a_{n*}) , a^* , a_* be the upper and lower sequences together with their limits. We call a^* the *upper limit* of the sequence (a_n) and a_* the *lower limit* of the sequence (a_n) . If they are equal, $a^* = a_*$, we say that $L = a^* = a_*$ is the *limit* of (a_n) , and we write

$$\lim_{n \nearrow \infty} a_n = L.$$

Alternatively, we say a_n *approaches* L , and we write $a_n \rightarrow L$ as $n \nearrow \infty$ or just $a_n \rightarrow L$. If they are not equal, $a^* \neq a_*$, we say that (a_n) *does not have a limit*.

If (a_n) is monotone, let L be its limit as a monotone sequence. Then, its upper and lower sequences are equal to itself and the constant sequence (L, L, \dots) . Thus, its upper limit is L , and its lower limit is L . Hence, L is its limit according to the second definition. In other words, *the two definitions are consistent*.

Let us look at an example. Take $a_n = (-1)^n/n$, $n \geq 1$, or

$$(a_n) = \left(-1, \frac{1}{2}, -\frac{1}{3}, \frac{1}{4}, \dots\right).$$

Then,

$$(a_n^*) = \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \dots\right),$$

$$(a_{n*}) = \left(-1, -\frac{1}{3}, -\frac{1}{3}, -\frac{1}{5}, -\frac{1}{5}, \dots\right).$$

Hence, $a^* = a_* = 0$, thus, $a_n \rightarrow 0$.

A *constant* sequence is a sequence of the form (a, a, \dots) where a is a fixed real. Here $a_n^* = a_{n*} = a$ for all $n \geq 1$. Hence, $a^* = a_* = a$, hence, $L = a$.

Not every sequence has a limit. For example $(1, 0, 1, 0, 1, 0, \dots)$ does not have a limit. Indeed, here, $a_n^* = 1$ and $a_{n*} = 0$ for all $n \geq 1$, hence, $a_* = 0 < 1 = a^*$.

Limits of sequences satisfy simple properties. For example, $a_n \rightarrow a$ implies $-a_n \rightarrow -a$, and $a_n \rightarrow L$ iff $a_{N+n} \rightarrow L$. Thus, in a very real sense, the limiting behaviour of a sequence does not depend on the first N terms of the sequence, for any $N \geq 1$. Here is the ordering property for sequences.

Theorem. Suppose that (a_n) , (b_n) , and (c_n) are sequences with $a_n \leq b_n \leq c_n$ for all $n \geq 1$. If $b_n \rightarrow K$ and $c_n \rightarrow L$, then, $K \leq L$. If $a_n \rightarrow L$ and $c_n \rightarrow L$, then, $b_n \rightarrow L$.

Note that, in the second assertion, the existence of the limit of (b_n) is not assumed, but is rather part of the conclusion. Why is this theorem true? Well, c_1^* is an upper bound for the set $\{c_k : k \geq 1\}$. Since $b_k \leq c_k$ for all k , c_1^* is an upper bound for $\{b_k : k \geq 1\}$. Since b_1^* is the least such, $b_1^* \leq c_1^*$. Repeating this argument with k starting at n , instead of at 1, yields $b_n^* \leq c_n^*$ for all $n \geq 1$. Repeating the same reasoning again yields $b^* \leq c^*$. If $b_n \rightarrow K$ and $c_n \rightarrow L$, then, $b^* = K$ and $c^* = L$, so, $K \leq L$, establishing the first assertion. To establish the second, we know that $b^* \leq c^*$. Now, set $C_n = -a_n$ and $B_n = -b_n$ for all $n \geq 1$. Then, $B_n \leq C_n$ for all $n \geq 1$, so, by what we just learned, $B^* \leq C^*$. But $B^* = -b_*$ and $C^* = -a_*$, so, $a_* \leq b_*$. We conclude that $a_* \leq b_* \leq b^* \leq c^*$. If $a_n \rightarrow L$ and $c_n \rightarrow L$, then, $a_* = L$ and $c^* = L$, hence, $b_* = b^* = L$. \square

As an application, $2^{-n} \rightarrow 0$ as $n \nearrow \infty$ since $0 < 2^{-n} < 1/n$ for all $n \geq 1$. Similarly, $\lim_{n \nearrow \infty} \left(\frac{1}{n} - \frac{1}{n^2}\right) = 0$ since

$$-\frac{1}{n} \leq -\frac{1}{n^2} \leq \frac{1}{n} - \frac{1}{n^2} \leq \frac{1}{n}$$

for all $n \geq 1$ and $\pm 1/n \rightarrow 0$ as $n \nearrow \infty$.

Let (a_n) be a sequence with nonnegative terms. Often the ordering property is used to show that $a_n \rightarrow 0$ by finding a sequence (e_n) satisfying $0 \leq a_n \leq e_n$ for all $n \geq 1$ and $e_n \rightarrow 0$.

Below and throughout the text, we will use the following easily checked fact: *If a and b are reals and $a < b + \epsilon$ for all real $\epsilon > 0$, then, $a \leq b$.* Indeed, either $a \leq b$ or $a > b$. If the latter case occurs, we may choose $\epsilon = (a - b)/2 > 0$, yielding the contradiction $a = b + (a - b) > b + \epsilon$. Thus, the former case must occur, or $a \leq b$. Moreover, *if a and b are reals and $b \leq a < b + \epsilon$ for all $\epsilon > 0$, then, $a = b$.*

Throughout the text, ϵ will denote a positive real number.

Theorem. *If $a_n \rightarrow a$ and $b_n \rightarrow b$ with a, b real, then, $\max(a_n, b_n) \rightarrow \max(a, b)$ and $\min(a_n, b_n) \rightarrow \min(a, b)$. Moreover, for any sequence (a_n) and L real, $a_n \rightarrow L$ iff $a_n - L \rightarrow 0$ iff $|a_n - L| \rightarrow 0$.*

Let $c_n = \max(a_n, b_n)$, $n \geq 1$, $c = \max(a, b)$, and let us assume, first, that the sequences (a_n) , (b_n) are decreasing. Then, their limits are their infs, and $c_n = \max(a_n, b_n) \geq \max(a, b) = c$. Hence, setting $c_* = \inf\{c_n : n \geq 1\}$, we conclude that $c_* \geq c$. On the other hand, given $\epsilon > 0$, there are n and m satisfying $a_n < a + \epsilon$ and $b_m < b + \epsilon$, so, $c_{n+m} = \max(a_{n+m}, b_{n+m}) \leq \max(a_n, b_m) < \max(a + \epsilon, b + \epsilon) = c + \epsilon$. Thus, $c_* < c + \epsilon$. Since $\epsilon > 0$ is arbitrary and we already know $c_* \geq c$, we conclude that $c_* = c$. Since (c_n) is decreasing, we have shown that $c_n \rightarrow c$.

Now, assume (a_n) , (b_n) are increasing. Then, their limits are their sups, and $c_n = \max(a_n, b_n) \leq \max(a, b) = c$. Hence, setting $c^* = \sup\{c_n : n \geq 1\}$, we conclude that $c^* \leq c$. On the other hand, given $\epsilon > 0$, there are n and m satisfying $a_n > a - \epsilon$ and $b_m > b - \epsilon$, so, $c_{n+m} = \max(a_{n+m}, b_{n+m}) \geq \max(a_n, b_m) > \max(a - \epsilon, b - \epsilon) = c - \epsilon$. Thus, $c^* > c - \epsilon$. Since $\epsilon > 0$ is arbitrary and we already know $c^* \leq c$, we conclude that $c^* = c$. Since (c_n) is increasing, we have shown that $c_n \rightarrow c$.

Now, for a general sequence (a_n) , we have (a_n^*) decreasing, (a_{n*}) increasing, and

$$\max(a_{n*}, b_{n*}) \leq c_n \leq \max(a_n^*, b_n^*), \quad n \geq 1.$$

Thus, (c_n) lies between two sequences converging to $c = \max(a, b)$. By the ordering property, we conclude that $c_n \rightarrow \max(a, b)$.

Since $\min(a, b) = -\max(-a, -b)$, the second assertion follows from the first.

For the third assertion, assume, first, $a_n \rightarrow L$, and set $b_n = a_n - L$. Since $\sup(A - a) = \sup A - a$ and $\inf(A - a) = \inf A - a$, $b_n^* = a_n^* - L$, and $b_{n*} = a_{n*} - L$. Hence, $b^* = a^* - L = 0$, and $b_* = a_* - L = 0$. Thus, $a_n - L \rightarrow 0$. If $a_n - L \rightarrow 0$, then, $L - a_n \rightarrow 0$. Hence, $|a_n - L| = \max(a_n - L, L - a_n) \rightarrow 0$ by the first assertion. Conversely, since

$$-|a_n - L| \leq a_n - L \leq |a_n - L|, \quad n \geq 1,$$

$|a_n - L| \rightarrow 0$ implies $a_n - L \rightarrow 0$, by the ordering property. Since $a_n = (a_n - L) + L$, this implies $a_n \rightarrow L$. \square

Often this theorem will be used to show that $a_n \rightarrow L$ by finding a sequence (e_n) satisfying $|a_n - L| \leq e_n$ and $e_n \rightarrow 0$. For example, let $A \subset \mathbb{R}$ be bounded above.

Then, $\sup A - 1/n$ is not an upper bound for A , hence, for each $n \geq 1$, there is a real $x_n \in A$ satisfying $\sup A - 1/n < x_n \leq \sup A$, hence, $|x_n - \sup A| < 1/n$. By the above, we conclude that $x_n \rightarrow \sup A$. When A is not bounded above, for each $n \geq 1$, there is a real $x_n \in A$ satisfying $x_n > n$. Then, $x_n \rightarrow \infty = \sup A$. In either case, we conclude, *if $A \subset \mathbb{R}$, there is a sequence $(x_n) \subset A$ with $x_n \rightarrow \sup A$* . Similarly, *if $A \subset \mathbb{R}$, there is a sequence $(x_n) \subset A$ with $x_n \rightarrow \inf A$* .

Now we derive the arithmetic properties of limits.

Theorem. *If $a_n \rightarrow a$ and c is real, then, $ca_n \rightarrow ca$. Let a, b be real. If $a_n \rightarrow a$, $b_n \rightarrow b$, then, $a_n + b_n \rightarrow a + b$ and $a_nb_n \rightarrow ab$. Moreover, if $b \neq 0$, then, $a_n/b_n \rightarrow a/b$.*

If $c = 0$, there is nothing to show. If $c > 0$, set $b_n = ca_n$. Since $\sup(cA) = c \sup A$ and $\inf(cA) = c \inf A$, $b_n^* = ca_n^*$, $b_{n*} = ca_{n*}$, $b^* = ca^*$, $b_* = ca_*$. Hence, $a_n \rightarrow a$ implies $ca_n \rightarrow ca$. Since $(-c)a_n = -(ca_n)$, the case with c negative follows.

To derive the additive property, assume, first, that $a = b = 0$. We have to show that $a_n + b_n \rightarrow 0$. Then,

$$2 \min(a_n, b_n) \leq a_n + b_n \leq 2 \max(a_n, b_n), \quad n \geq 1.$$

Thus, $a_n + b_n$ lies between two sequences approaching 0, so, $a_n + b_n \rightarrow 0$. For general a, b , apply the previous to $a'_n = a_n - a$, $b'_n = b_n - b$.

To derive the multiplicative property, first, note that $a_{1*} \leq a_n \leq a_1^*$, so, $|a_n| \leq k$ for some k , i.e., (a_n) is bounded. Use the triangle inequality to get

$$\begin{aligned} |a_nb_n - ab| &= |(a_n - a)b + (b_n - b)a_n| \leq |b| |a_n - a| + |a_n| |b_n - b| \\ &\leq |b| |a_n - a| + k |b_n - b|, \quad n \geq 1. \end{aligned}$$

Now, the result follows from the additive and ordering properties.

To obtain the division property, assume $b > 0$. From the above, $a_nb - ab_n \rightarrow 0$. Since $b_n \rightarrow b$, $b_{n*} \nearrow b$, so, there exists $N \geq 1$ beyond which $b_n \geq b_{N*} > 0$ for $n \geq N$. Thus,

$$0 \leq \left| \frac{a_n}{b_n} - \frac{a}{b} \right| = \frac{|a_nb - ab_n|}{|b_n| |b|} \leq \frac{|a_nb - ab_n|}{b_{N*} b}, \quad n \geq N.$$

Thus, $|a_n/b_n - a/b|$ lies between zero and a sequence approaching zero. The case $b < 0$ is entirely similar. \square

In fact, although we do not derive this, this theorem remains true when a or b are infinite, as long as we do not allow undefined expressions, such as $\infty - \infty$ (the allowable expressions are defined in §1.2).

As an application of this theorem,

$$\lim_{n \nearrow \infty} \frac{2n^2 + 1}{n^2 - 2n + 1} = \lim_{n \nearrow \infty} \frac{2 + \frac{1}{n^2}}{1 - \frac{2}{n} + \frac{1}{n^2}} = \frac{2 + 0}{1 - 2 \cdot 0 + 0} = 2.$$

If (a_n) is a sequence with positive terms and $b_n = 1/a_n$, then, $a_n \rightarrow 0$ iff $b_n \rightarrow \infty$ (Exercise 3). Now, let $a > 1$ and set $b = a - 1$. Then, $a^n = (1 + b)^n \geq 1 + nb$ for all $n \geq 1$ (Exercise 6 of §1.4). Hence, $a^n \nearrow \infty$. If $0 < a < 1$, then, $a = 1/b$ with $b > 1$, so, $a^n = 1/b^n \searrow 0$. Summarizing,

- if $a > 1$, then, $a^n \nearrow \infty$,
- if $a = 1$, then, $a^n = 1$ for all $n \geq 1$, and
- if $0 \leq a < 1$, then, $a^n \rightarrow 0$.

Sometimes we say that a sequence (a_n) *converges to* L if $a_n \rightarrow L$. If the specific limit is not relevant, we say that the sequence *converges* or is *convergent*. If a sequence has no limit, we say it *diverges*. More precisely, if the sequence (a_n) does not approach L , we say that it *diverges from* L , and we write $a_n \not\rightarrow L$. From the definition of $a_n \rightarrow L$, we see that $a_n \not\rightarrow L$ means either $a^* \neq L$ or $a_* \neq L$. This is so whether L is real or $\pm\infty$.

Typically, divergence is *oscillatory behavior*, e.g., $a_n = (-1)^n$. Here, the sequence goes back and forth never settling on anything, not even ∞ or $-\infty$. Nevertheless, this sequence is bounded. Of course a sequence may be oscillatory and unbounded, e.g., $a_n = (-1)^n n$.

Let (a_n) be a sequence, and suppose that $1 \leq k_1 < k_2 < k_3 < \dots$ is an increasing sequence of distinct *naturals*. Set $b_n = a_{k_n}$, $n \geq 1$. Then, $(b_n) = (a_{k_n})$ is a *subsequence* of (a_n) . If $a_n \rightarrow L$, then, $a_{k_n} \rightarrow L$. Conversely, if (a_n) is monotone, $a_{k_n} \rightarrow L$ implies $a_n \rightarrow L$ (Exercise 4).


Generally, if a sequence (x_n) has a subsequence (x_{k_n}) converging to L , we say that (x_n) *subconverges to* L .

Let (a_n) converge to a (finite) real limit L , and let $\epsilon > 0$ be given. Since (a_{n*}) is increasing to L , there must exist a natural N_* , such that $a_{n*} > L - \epsilon$ for $n \geq N_*$. Similarly, there must exist N^* beyond which we have $a_n^* < L + \epsilon$. Since $a_{n*} \leq a_n \leq a_n^*$ for all $n \geq 1$, we obtain $L - \epsilon < a_n < L + \epsilon$ for $n \geq N = \max(N^*, N_*)$. Thus, all but finitely many terms of the sequence lie in $(L - \epsilon, L + \epsilon)$ (Figure 1.3).

Note that choosing a smaller $\epsilon > 0$ is a more stringent condition on the terms. As such, it leads to (in general) a larger N , i.e., the number of terms that fall outside the interval $(L - \epsilon, L + \epsilon)$ depends on the choice of $\epsilon > 0$.

Conversely, suppose that $L - \epsilon < a_n < L + \epsilon$ for all but finitely many terms, for every $\epsilon > 0$. Then, for a given $\epsilon > 0$, by the ordering property, $L - \epsilon \leq a_{n*} \leq a_n^* \leq L + \epsilon$ for all but finitely terms. Hence, $L - \epsilon \leq a_* \leq a^* \leq L + \epsilon$. Since this holds for every $\epsilon > 0$, we conclude that $a_* = a^* = L$, i.e., $a_n \rightarrow L$. We have derived the following:

Theorem. *Let (a_n) be a sequence and let L be real. If $a_n \rightarrow L$, then, all but finitely many terms of the sequence lie within the interval $(L - \epsilon, L + \epsilon)$, for all $\epsilon > 0$. Conversely, if all but finitely many terms lie in the interval $(L - \epsilon, L + \epsilon)$, for all $\epsilon > 0$, then, $a_n \rightarrow L$. \square*

FIGURE 1.3. Convergence to L .

From this, we conclude that, if $a_n \rightarrow L$ and $L \neq 0$, then, $a_n \neq 0$ for all but finitely many n .

We end the section with an application. Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is injective, i.e., suppose that $(f(n)) = (a_n)$ is a sequence consisting of *distinct* naturals. Then, $f(n) \rightarrow \infty$. To see this, note that (since f is injective) for each n natural, there are only finitely many naturals k satisfying $f(k) < n$. Let $k_n = \max\{k : f(k) < n\}$. Then, $k > k_n$ implies $f(k) \geq n$ which implies $f_*(k_n + 1) = \inf\{f(k) : k > k_n\} \geq n$. Hence, $f_*(k_n + 1) \nearrow \infty$, as $n \nearrow \infty$. Since $(f_*(n))$ is monotone and $(f_*(k_n + 1))$ is a subsequence of $(f_*(n))$, it follows that $f_*(n) \nearrow \infty$, as $n \nearrow \infty$ (Exercise 4). Since $f(n) \geq f_*(n)$, we conclude that $f(n) \rightarrow \infty$.

Exercises 1.5

1. Fix $N \geq 1$ and (a_n) . Let (a_{N+n}) be the sequence $(a_{N+1}, a_{N+2}, \dots)$. Then, $a_n \nearrow L$ iff $a_{N+n} \nearrow L$, and $a_n \searrow L$ iff $a_{N+n} \searrow L$. Conclude that $a_n \rightarrow L$ iff $a_{n+N} \rightarrow L$.
2. If $a_n \rightarrow L$, then, $-a_n \rightarrow -L$.
3. If $A \subset \mathbb{R}^+$ is nonempty and $1/A = \{1/x : x \in A\}$, then, $\inf(1/A) = 1/\sup A$, where $1/\infty$ is interpreted here as 0. If (a_n) is a sequence with positive terms and $b_n = 1/a_n$, then, $a_n \rightarrow 0$ iff $b_n \rightarrow \infty$.
4. If $a_n \rightarrow L$ and (a_{k_n}) is a subsequence, then, $a_{k_n} \rightarrow L$. If (a_n) is monotone and $a_{k_n} \rightarrow L$, then, $a_n \rightarrow L$.
5. If $a_n \rightarrow L$ and $L \neq 0$, then, $a_n \neq 0$ for all but finitely many n .
6. Let $a_n = \sqrt{n+1} - \sqrt{n}$, $n \geq 1$. Compute (a_n^*) , (a_{n*}) , a^* , and a_* . Does (a_n) converge?
7. Let (a_n) be any sequence with upper and lower limits a^* and a_* . Show that (a_n) subconverges to a^* and subconverges to a_* , i.e., there are subsequences (a_{k_n}) and (a_{j_n}) satisfying $a_{k_n} \rightarrow a^*$ and $a_{j_n} \rightarrow a_*$.
8. Suppose that (a_n) diverges from $L \in \mathbb{R}$. Show that there is an $\epsilon > 0$ and a subsequence (a_{k_n}) satisfying $|a_{k_n} - L| \geq \epsilon$ for all $n \geq 1$.
9. Let (x_n) be a sequence. If (x_n) subconverges to L , we say that L is a *limit point* of (x_n) . Show that x_* and x^* are the least and the greatest limit points.
10. Show that a sequence (x_n) converges iff (x_n) has exactly one limit point.
11. Given $f : (a, b) \rightarrow \mathbb{R}$, let $M = \sup\{f(x) : a < x < b\}$. Show that there is a sequence (x_n) with $f(x_n) \rightarrow M$. (Consider the cases $M < \infty$ and $M = \infty$.)

12. ★ Define (d_n) by $d_1 = 2$ and

$$d_{n+1} = \frac{1}{2} \left(d_n + \frac{2}{d_n} \right), \quad n \geq 1,$$

and set $e_n = d_n - \sqrt{2}$, $n \geq 1$. By induction, show that $e_n \geq 0$ for $n \geq 1$. Also show that

$$e_{n+1} \leq \frac{e_n^2}{2\sqrt{2}}, \quad \text{for all } n \geq 1.$$

(First, check that, for any real $x > 0$, one has $(x + 2/x)/2 \geq \sqrt{2}$.)

13. ★ Let $0 < x < 1$ be irrational, and let (q_n) be as in Exercise 16 of §1.3. Let

$$x_n = \frac{1}{q_1 + \frac{1}{q_2 + \frac{1}{\ddots + \frac{1}{q_{n-1} + \frac{1}{q_n}}}}}.$$

Let x' and x'_n denote the continued fractions starting with $1/(q_2 + \dots)$, i.e., with the top layer “peeled off.” Then, $0 < x_n, x', x'_n < 1$.

- (1) Show that $|x - x_n| \leq x x_n |x' - x'_n|$, $n \geq 1$.
- (2) Iterate (1) to show that $|x - x_n| \leq 1/q_n$, $n \geq 1$.
- (3) Show that $x \leq (q_2 + 1)/(q_2 + 2)$, $x' \leq (q_3 + 1)/(q_3 + 2)$, etc.
- (4) If N of the q_k 's are bounded by c , iterate (1) and use (3) to obtain

$$|x - x_n| \leq \left(\frac{c+1}{c+2} \right)^N,$$

for all but finitely many n .

Conclude that $|x - x_n| \rightarrow 0$ as $n \nearrow \infty$. (Either $q_n \rightarrow \infty$ or $q_n \not\rightarrow \infty$.)

§1.6 Nonnegative Series and Decimal Expansions

Let (a_n) be a sequence of reals. The *series* formed from the sequence (a_n) is the sequence (s_n) with terms $s_1 = a_1$, $s_2 = a_1 + a_2$, and, for any $n \geq 1$, $s_n = a_1 + a_2 + \dots + a_n$. The sequence (s_n) is the sequence of *partial sums*. The terms a_n are called *summands*, and the series is *nonnegative* if $a_n \geq 0$ for all $n \geq 1$. We often use sigma notation, and write $s_n = \sum_{k=1}^n a_k$. Series are often written

$$a_1 + a_2 + \dots$$

In sigma notation, $\sum a_n$ or $\sum_{n=1}^{\infty} a_n$. If the sequence of partial sums (s_n) has a limit L , then, we say the series *sums* or *converges* to L , and we write

$$L = a_1 + a_2 + \cdots = \sum_{n=1}^{\infty} a_n.$$

Then, L is the *sum* of the series. *By convention, we do not allow $\pm\infty$ as limits for series, only reals.* Nevertheless, for nonnegative series, we write $\sum a_n = \infty$ to mean $\sum a_n$ diverges and $\sum a_n < \infty$ to mean $\sum a_n$ converges. As with sequences, sometimes it is more convenient to start a series from $n = 0$. In this case, we write $\sum_{n=0}^{\infty} a_n$.

Let $L = \sum_{n=1}^{\infty} a_n$ be a convergent series and let s_n denote its n th partial sum. The n th *tail* of the series is $L - s_n = \sum_{k=n+1}^{\infty} a_k$. Since the n th tail is the difference between the n th partial sum and the sum, we see that *the n th tail of a convergent series goes to zero*:

$$\lim_{n \nearrow \infty} \sum_{k=n+1}^{\infty} a_k = 0.$$

Let a be real. Our first series is the *geometric series*

$$1 + a + a^2 + \cdots = \sum_{n=0}^{\infty} a^n.$$

Here the n th partial sum $s_n = 1 + a + \cdots + a^n$ is computed as follows:

$$as_n = a(1 + a + \cdots + a^n) = a + a^2 + \cdots + a^{n+1} = s_n + a^{n+1} - 1.$$

Hence,

$$s_n = \frac{1 - a^{n+1}}{1 - a}, \quad a \neq 1.$$

If $a = 1$, then, $s_n = n$, so, $s_n \nearrow \infty$. If $|a| < 1$, $a^n \rightarrow 0$, so, $s_n \rightarrow 1/(1 - a)$. If $a > 1$, then, $a^n \nearrow \infty$, so, the series equals ∞ and hence, diverges. If $a < -1$, then, (a^n) diverges, so, the series diverges. If $a = -1$, s_n equals 0 or 1 (depending on n), hence, diverges, hence, the series diverges. We have shown

$$\sum_{n=0}^{\infty} a^n = \frac{1}{1 - a}, \quad \text{if } |a| < 1,$$

and $\sum_{n=0}^{\infty} a^n$ diverges if $|a| \geq 1$.

To study more general series, we need their arithmetic and ordering properties.

Theorem. If $\sum a_n = L$ and $\sum b_n = M$, then, $\sum (a_n + b_n) = L + M$. If $\sum a_n = L$, $c \in \mathbb{R}$, and $b_n = ca_n$, then, $\sum b_n = cL = c(\sum a_n)$. If $a_n \leq b_n \leq c_n$ and $\sum a_n = L = \sum c_n$, then, $\sum b_n = L$.

To see the first property, if s_n , t_n , and r_n denote the partial sums of $\sum a_n$, $\sum b_n$, and $\sum c_n$, then, $s_n + t_n$ equals the partial sum of $\sum (a_n + b_n)$. Hence, the result

follows from the corresponding arithmetic property of sequences. For the second property, note that $t_n = cs_n$. Hence, the result follows from the corresponding arithmetic property of sequences. The third property follows from the ordering property of sequences, since $s_n \leq t_n \leq r_n$, $s_n \rightarrow L$, and $r_n \rightarrow L$. \square

Now, we describe the comparison test which we use below to obtain the decimal expansions of reals.

Theorem (Comparison Test). *Let $\sum a_n$, $\sum b_n$ be nonnegative series with $a_n \leq b_n$ for all $n \geq 1$. If $\sum b_n < \infty$, then, $\sum a_n < \infty$. If $\sum a_n = \infty$, then, $\sum b_n = \infty$.*

Stated this way, the theorem follows from the ordering property for sequences and looks too simple to be of any serious use. \square In fact, we use it to express every real as a sequence of naturals. Recall that the *digits* are defined in §1.3. Define $d = 9 + 1$.

Theorem. *If d_1, d_2, \dots is a sequence of digits, then,*

$$\sum_{n=1}^{\infty} d_n d^{-n}$$

sums to a real x , $0 \leq x \leq 1$. Conversely, if $0 \leq x \leq 1$, there is a sequence of digits d_1, d_2, \dots , such that the series sums to x .

The first statement follows by comparison, since

$$\begin{aligned} \sum_{n=1}^{\infty} d_n d^{-n} &\leq \sum_{n=1}^{\infty} 9d^{-n} = \sum_{n=0}^{\infty} \frac{9}{d} d^{-n} \\ &= \frac{9}{d} \sum_{n=0}^{\infty} d^{-n} = \frac{9}{d} \cdot \frac{1}{1 - (1/d)} = 1. \end{aligned}$$

To establish the second statement, if $x = 1$, we simply take $d_n = 9$ for all $n \geq 1$. If $0 \leq x < 1$, let d_1 be the largest integer $\leq dx$. Then, $d_1 \geq 0$. This way we obtain a digit d_1 (since $x < 1$) satisfying $d_1 \leq dx < d_1 + 1$. Now, set $x_1 = dx - d_1$. Then, $0 \leq x_1 < 1$. Repeating the above process, we obtain a digit d_2 satisfying $d_2 \leq dx_1 < d_2 + 1$. Substituting yields $d_2 + dd_1 \leq d^2x < d_2 + dd_1 + 1$ or $d_2d^{-2} + d_1d^{-1} \leq x < d_2d^{-2} + d_1d^{-1} + d^{-2}$. Continuing in this manner yields a sequence of digits (d_n) satisfying

$$\left(\sum_{k=1}^n d_k d^{-k} \right) \leq x < \left(\sum_{k=1}^n d_k d^{-k} \right) + d^{-n}, \quad n \geq 1.$$

Thus, x lies between two sequences converging to the same limit. \square

In fact, two distinct sequences of digits yield the same real in $[0, 1]$ under only very special circumstances. The sequence (d_n) is the *decimal expansion* of the real $0 \leq x \leq 1$. As usual, we write

$$x = .d_1d_2d_3\dots$$

Thus $1/3 = .333\dots$

To extend the decimal notation to any nonnegative real, for each $x \geq 1$, there is a smallest natural N , such that $d^{-N}x < 1$. As usual, if $d^{-N}x = .d_1d_2\dots$, we write

$$x = d_1d_2\dots d_N.d_{N+1}d_{N+2}\dots,$$

the *decimal point* (.) moved N places. For example $1 = 1.00\dots$ and $d = 10.00\dots$. In fact, x is a natural iff $x = d_1d_2\dots d_N.00\dots$. Thus, for naturals, we drop the decimal point and the trailing zeros, e.g., $1 = 1$, $d = 10$. Note that we have two decimal representations of 1, $1 = .99\dots = 1.00\dots$. This is not an accident (Exercise 2).

The natural d , the *base* of the expansion, can be replaced by any natural > 1 . Then, the digits are $(0, 1, \dots, d-1)$, and we would obtain d -ary expansions. In §4.1, we use $d = 2$ with digits $(0, 1)$ leading to *binary expansions* and $d = 3$ with digits $(0, 1, 2)$ leading to *ternary expansions*. In §5.2, we discuss $d = 16$ with digits $(0, 1, \dots, 9, A, B, C, D, E, F)$ leading to *hexadecimal expansions*. This completes our discussion of decimal expansions.

How can one tell if a given series converges by inspecting the individual terms? Here is a necessary condition.

Theorem (*n*th Term Test). *If $\sum a_n = L \in \mathbf{R}$, then, $a_n \rightarrow 0$.*

To see this, we know that $s_n \rightarrow L$, and, so, $s_{n-1} \rightarrow L$. By the triangle inequality, $|a_n| = |s_n - s_{n-1}| = |(s_n - L) + (L - s_{n-1})| \leq |s_n - L| + |s_{n-1} - L| \rightarrow 0$. \square

However, a series whose n th term approaches zero need not converge. For example, the *harmonic series*

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \dots = \infty.$$

To see this, use comparison as follows,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n} &= 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \dots \\ &\geq 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{4} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \dots \\ &= 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \dots = \infty. \end{aligned}$$

On the other hand, let $0! = 1$ and let $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot n$ (n factorial) for $n \geq 1$. Then, the nonnegative series

$$\frac{1}{0!} + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots$$

converges. To see this, check that $2^{n-1} \leq n!$ by induction. Thus,

$$1 + \sum_{n=1}^{\infty} \frac{1}{n!} \leq 1 + \sum_{n=1}^{\infty} 2^{-n+1} = 3$$

and, hence, is convergent. Since the third partial sum is $s_3 = 2.5$, we see that the sum lies in the interval $(2.5, 3]$.

A series is *telescoping* if it is a sum of differences, i.e., of the form

$$\sum_{n=1}^{\infty} (a_n - a_{n+1}) = (a_1 - a_2) + (a_2 - a_3) + (a_3 - a_4) + \dots$$

In this case, the following is true.

Theorem. *If (a_n) is any sequence converging to zero, then, the corresponding telescoping series converges, and its sum is a_1 .*

This follows since the partial sums are

$$s_n = (a_1 - a_2) + (a_2 - a_3) + \dots + (a_n - a_{n+1}) = a_1 - a_{n+1}$$

and $a_{n+1} \rightarrow 0$. \square

As an application, note that

$$\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots = 1$$

since

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right) = 1.$$

Another application is to establish the convergence of

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^2} &= 1 + \sum_{n=2}^{\infty} \frac{1}{n^2} \\ &< 1 + \sum_{n=2}^{\infty} \frac{1}{n(n-1)} = 1 + \sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 2. \end{aligned}$$

Thus,

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots < 2.$$

Expressing this sum in terms of familiar quantities is a question of a totally different magnitude. Later (§5.6), we will see how this is done.

More generally,

$$\sum_{n=1}^{\infty} \frac{1}{n^{1+1/N}} < \infty$$

follows in a similar manner, for any $N \geq 1$.

Exercises 1.6

1. Let $0 < x < 1$ be real. Then, $x = .d_1d_2\dots$ is in \mathbb{Q} iff there are $n, m \geq 1$, such that $d_{j+n} = d_j$ for $j > m$, i.e., the sequence of digits repeats every n digits, from the $(m+1)$ st digit on.

2. Suppose that $.d_1d_2\dots = .e_1e_2\dots$ are distinct decimal expansions for the same real, and let N be the first natural with $d_N \neq e_N$. Then, either $d_N = e_N + 1$, $d_k = 0$, and $e_k = 9$ for $k > N$, or $e_N = d_N + 1$, $e_k = 0$, and $d_k = 9$ for $k > N$. Conclude that $x \in \mathbb{R}$ has more than one decimal expansion iff $10^N x \in \mathbb{Z}$ for some $N \in \mathbb{N} \cup \{0\}$.

3. Show that $2^{n-1} \leq n!$, $n \geq 1$.

4. Fix $N \geq 1$. Show that

$$(1) \left(\frac{n}{n+1}\right)^{1/N} \leq 1 - \frac{1}{N(n+1)}, n \geq 1,$$

$$(2) (n+1)^{1/N} - n^{1/N} \geq \frac{1}{N(n+1)^{(N-1)/N}}, n \geq 1, \text{ and}$$

$$(3) \sum_{n=2}^{\infty} \frac{1}{n^{1+1/N}} \leq N \sum_{n=1}^{\infty} (a_n - a_{n+1}) \text{ where } a_n = \frac{1}{n^{1/N}}, n \geq 1.$$

Conclude that $\sum_{n=1}^{\infty} \frac{1}{n^{1+1/N}} < \infty$. (Use Exercises 6 and 7 of §1.4 for (1)).

5. ★ Let (d_n) and (e_n) be as in Exercise 12 of §1.5. By induction, show that

$$e_{n+2} \leq 10^{-2^n}, \quad n \geq 1.$$

This shows that the decimal expansion of d_{n+2} agrees⁹ with that of $\sqrt{2}$ to 2^n places. For example, d_9 yields $\sqrt{2}$ to at least 128 decimal places. (First, show that $e_3 \leq 1/100$. Since the point here is to compute the decimal expansion of $\sqrt{2}$, do not use it in your derivation. Use only $1 < \sqrt{2} < 2$ and $(\sqrt{2})^2 = 2$.)

6. ★ Let $C \subset [0, 1]$ be the set of reals $x = .d_1d_2d_3\dots$ whose decimal digits d_n , $n \geq 1$, are zero or odd. Show that (§1.2) $C + C = [0, 2]$.

§1.7 Signed Series and Cauchy Sequences

A series is *signed* if its first term is positive, and at least one of its terms is negative. A series is *alternating* if it is of the form

$$\sum_{n=1}^{\infty} (-1)^{n-1} a_n = a_1 - a_2 + a_3 \cdots + (-1)^{n-1} a_n + \dots$$

with a_n positive for all $n \geq 1$. Alternating series are particularly tractable, but, first, we need a new concept.

⁹This algorithm was known to the Babylonians.

A sequence (not a series!) (a_n) is *Cauchy* if its terms approach each other, i.e., if there is a positive sequence (e_n) converging to zero, such that

$$|a_{n+m} - a_n| \leq e_n, \quad \text{for all } m, n \geq 1.$$

If a sequence is Cauchy, there are many choices for (e_n) . Any such sequence (e_n) is an *error sequence* for the Cauchy sequence (a_n) .

It follows from the definition that every Cauchy sequence is bounded, $|a_m| \leq |a_m - a_1| + |a_1| \leq e_1 + |a_1|$ for all $m \geq 1$.

It is easy to see that a convergent sequence is Cauchy. Indeed, if (a_n) converges to L , then, $b_n = |a_n - L| \rightarrow 0$, so (§1.5), $b_n^* \rightarrow 0$. Hence, by the triangle inequality

$$|a_{n+m} - a_n| \leq |a_{n+m} - L| + |a_n - L| \leq b_n^* + b_n^*, \quad m > 0, n \geq 1.$$

Since $2b_n^* \rightarrow 0$, $(2b_n^*)$ is an error sequence for (a_n) , so, (a_n) is Cauchy.

The following theorem shows that if the terms of a sequence “approach each other”, then, they “approach something”. To see that this is not a self-evident assertion, consider the following example. Let a_n be the rational given by the first n places in the decimal expansion of $\sqrt{2}$. Then, $|a_n - \sqrt{2}| \leq 10^{-n}$, hence, $a_n \rightarrow \sqrt{2}$, hence, (a_n) is Cauchy. But, as far as \mathbb{Q} is concerned, there is no limit, since $\sqrt{2} \notin \mathbb{Q}$. In other words, to actually establish the existence of the limit, one needs an additional property not enjoyed by \mathbb{Q} , the completeness property of \mathbb{R} .

Theorem. *A Cauchy sequence (a_n) is convergent.*

With the notation of §1.5, we need to show that $a_* = a^*$. But this follows since the sequence is Cauchy. Indeed, let (e_n) be any error sequence. Then, for all $n \geq 1, m \geq 0, j \geq 0$,

$$a_{n+m} - a_{n+j} \leq (a_{n+m} - a_n) + (a_n - a_{n+j}) \leq 2e_n.$$

For n and j fixed, this inequality is true for all $m \geq 0$. Taking the sup over all $m \geq 0$ yields

$$a_n^* - a_{n+j} \leq 2e_n$$

for all $j \geq 0, n \geq 1$. Now, for n fixed, this inequality is true for all $j \geq 0$. Taking the sup over all $j \geq 0$ and using $\sup(-A) = -\inf A$ yields

$$0 \leq a_n^* - a_{n*} \leq 2e_n, \quad n \geq 1.$$

Letting $n \nearrow \infty$ yields $0 \leq a^* - a_* \leq 0$, hence, $a^* = a_*$. \square

A series $\sum a_n$ is said to be *absolutely convergent* if $\sum |a_n|$ converges. For example, below, we will see that $\sum (-1)^{n-1}/n$ converges. Since $\sum 1/n$ diverges, however, $\sum (-1)^{n-1}/n$ does not converge absolutely. A convergent series that is not absolutely convergent is *conditionally convergent*.

If $\sum |a_n|$ is known to converge, one expects $\sum a_n$ to converge, because of the possibility of cancellation. In fact, this is the case.

Theorem. *If $\sum a_n$ converges absolutely, then, $\sum a_n$ converges.*

To see this, let (s_n) and (S_n) denote the sequences of partial sums corresponding to $\sum a_n$ and $\sum |a_n|$. Since $\sum |a_n|$ converges, we know that (S_n) is Cauchy. Let (e_n) be an error sequence for (S_n) . To show that (s_n) converges, it is enough to show that (s_n) is Cauchy. But, by the triangle inequality,

$$\begin{aligned} |s_{n+m} - s_n| &= |a_{n+1} + \cdots + a_{n+m}| \\ &\leq |a_{n+1}| + \cdots + |a_{n+m}| = S_{n+m} - S_n \leq e_n, \end{aligned}$$

so, (e_n) is an error sequence for (s_n) . \square

A typical application of this result is as follows. *If (a_n) is a sequence of positive reals decreasing to zero and (b_n) is bounded, then,*

$$(7.1) \quad \sum_{n=1}^{\infty} (a_n - a_{n+1})b_n$$

converges absolutely. Indeed, if $|b_n| \leq C, n \geq 1$, is a bound for (b_n) , then,

$$\sum_{n=1}^{\infty} |(a_n - a_{n+1})b_n| \leq \sum_{n=1}^{\infty} (a_n - a_{n+1})C = Ca_1 < \infty$$

since the last series is telescoping.

To extend the scope of this last result, we will need the following elementary formula:

$$(7.2) \quad a_1 b_1 + \sum_{n=2}^N a_n (b_n - b_{n-1}) = \sum_{n=1}^{N-1} (a_n - a_{n+1})b_n + a_N b_N.$$

This important identity, easily verified by decomposing the sums, is called *summation by parts*.

Theorem (Dirichlet Test). *If (a_n) is a positive sequence decreasing to zero and (c_n) is such that the sequence $b_n = c_1 + c_2 + \cdots + c_n, n \geq 1$, is bounded, then, $\sum_{n=1}^{\infty} a_n c_n$ converges and*

$$(7.3) \quad \sum_{n=1}^{\infty} a_n c_n = \sum_{n=1}^{\infty} (a_n - a_{n+1})b_n.$$

This is an immediate consequence of letting $N \nearrow \infty$ in (7.2) since $b_n - b_{n-1} = c_n$ for $n \geq 2$. \square An important aspect of the Dirichlet test is that the right side of (7.3) is, from above, absolutely convergent, whereas the left side is often only conditionally convergent. For example, taking $a_n = 1/n$ and $c_n = (-1)^{n-1}, n \geq 1$, yields $(b_n) = (1, 0, 1, 0, \dots)$. Hence, we conclude not only that

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots$$

converges but also that its sum equals the sum of the absolutely convergent series obtained by grouping the terms in pairs.

Now, we can state the situation with alternating series.

Theorem (Leibnitz Test). *If (a_n) is a positive decreasing sequence with $a_n \searrow 0$, then,*

$$(7.4) \quad a_1 - a_2 + a_3 - a_4 + \dots$$

converges to a limit L satisfying $0 \leq L \leq a_1$. If, in addition, (a_n) is strictly decreasing, then, $0 < L < a_1$. Moreover, if s_n denotes the n th partial sum, $n \geq 1$, then, the error $|L - s_n|$ at the n th stage is no greater than the $(n + 1)$ st term a_{n+1} , with $L \geq s_n$ or $L \leq s_n$ according to whether the $(n + 1)$ st term is added or subtracted.

For example,

$$(7.5) \quad L = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \dots$$

converges, and $0 < L < 1$. In fact, since $s_2 = 2/3$ and $s_3 = 13/15$, $2/3 < L < 13/15$.

In the previous section, estimating the sum of

$$1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots$$

to one decimal place involved estimating the entire series. Here, the situation is markedly different: The absolute error between the sum and the n th partial sum is no larger than the next term a_{n+1} .

To derive the Leibnitz test, clearly, the convergence of (7.4) follows by taking $c_n = (-1)^{n-1}$ and applying the Dirichlet test, as above. Now, the differences $a_n - a_{n+1}$, $n = 1, 3, 5, \dots$, are nonnegative. Grouping the terms in (7.4) in pairs, we obtain $L \geq 0$. Similarly, the differences $-a_n + a_{n+1}$, $n = 2, 4, 6, \dots$, are nonpositive. Grouping the terms in (7.4) in pairs, we obtain $L \leq a_1$. Thus, $0 \leq L \leq a_1$. But

$$(-1)^n(L - s_n) = a_{n+1} - a_{n+2} + a_{n+3} - a_{n+4} + \dots, \quad n \geq 1.$$

Repeating the above reasoning, we obtain $0 \leq (-1)^n(L - s_n) \leq a_{n+1}$, which implies the rest of the statement. If, in addition, (a_n) is strictly decreasing, this reasoning yields $0 < L < a_1$. \square

If $\sum a_n$ converges absolutely and $s_n = \sum_{k=1}^n |a_k|$, then, by the triangle inequality,

$$|s_m - s_n| = \sum_{k=n+1}^m |a_k| \leq \sum_{k=n+1}^{\infty} |a_k|, \quad m > n,$$

so, $e_n = \sum_{j=n+1}^{\infty} |a_j|$, $n \geq 1$, is an error sequence for (s_n) .

Our next topic is the rearrangement of series. A series $\sum A_n$ is a *rearrangement* of a series $\sum a_n$ if there is a bijection (§1.1) $f : \mathbb{N} \rightarrow \mathbb{N}$, such that $A_n = a_{f(n)}$ for all $n \geq 1$. Absolutely convergent series and conditionally convergent series behave very differently under rearrangements.

Theorem. *If $\sum a_n$ is absolutely convergent, any rearrangement $\sum A_n$ converges absolutely to the same limit. If $\sum a_n$ is conditionally convergent and c is any real number, then, there is a rearrangement $\sum A_n$ converging to c .*

To see this, first assume $\sum |a_n|$ is convergent, and let $e_n = \sum_{j=n+1}^{\infty} |a_j|$, $n \geq 1$. Then, (e_n) is an error sequence for the Cauchy sequence of partial sums of $\sum |a_n|$. Since (§1.5) $(f(n))$ is a sequence of distinct naturals, $f(n) \rightarrow \infty$. In fact (§1.5), if we let $f_*(n) = \inf\{f(k) : k \geq n\}$, then, $f_*(n) \nearrow \infty$. To show that $\sum |A_n|$ is convergent, it is enough to show that $\sum |A_n|$ is Cauchy.

To this end,

$$\begin{aligned} & |a_{f(n+1)}| + |a_{f(n+2)}| + \cdots + |a_{f(n+m)}| \\ & \leq |a_{f_*(n+1)}| + |a_{f_*(n+1)+1}| + |a_{f_*(n+1)+2}| + \cdots = e_{f_*(n+1)-1} \end{aligned}$$

which approaches zero, as $n \nearrow \infty$. Thus, $\sum |A_n|$ is Cauchy, hence, convergent. Hence, $\sum A_n$ is absolutely convergent.

Now, let s_n, S_n denote the partial sums of $\sum a_n$ and $\sum A_n$, respectively. Let $E_n = \sum_{k=n+1}^{\infty} |A_k|$, $n \geq 1$. Then, (E_n) is an error sequence for the Cauchy sequence of partial sums of $\sum |A_n|$. Now, in the difference $S_n - s_n$, there will be cancellation, the only terms remaining being of one of two forms, either $A_k = a_{f(k)}$ with $f(k) > n$ or a_k with $k = f(j)$ with $j > n$ (this is where surjectivity of f is used). Hence, in either case, the absolute values of the remaining terms in $S_n - s_n$ are summands in the series $e_n + E_n$, so,

$$|S_n - s_n| \leq e_n + E_n \rightarrow 0, \quad \text{as } n \nearrow \infty.$$

This completes the derivation of the absolute portion of the theorem.

Now, assume that $\sum a_n$ is conditionally convergent, and let $c \geq 0$ be any nonnegative real. Let (a_n^+) , (a_n^-) denote the positive and the negative terms in the series $\sum a_n$. Then, we must have $\sum a_n^+ = \infty$ and $\sum a_n^- = -\infty$. Otherwise, $\sum a_n$ would converge absolutely. Moreover, $a_n^+ \rightarrow 0$ and $a_n^- \rightarrow 0$ since $a_n \rightarrow 0$. We construct a rearrangement as follows: Take the minimum number of terms a_n^+ whose sum s_1^+ is greater than c , then, take the minimum number of terms a_n^- whose sum s_1^- with s_1^+ is less than c , then, take the minimum number of additional terms a_n^+ whose sum s_2^+ with s_1^- is greater than c , then, take the minimum number of additional terms a_n^- whose sum s_2^- with s_2^+ is less than c , etc. Because $a_n^+ \rightarrow 0$, $a_n^- \rightarrow 0$, $\sum a_n^+ = \infty$, and $\sum a_n^- = -\infty$, this rearrangement of the terms produces a series converging to c . Of course, if $c < 0$, one starts, instead, with the negative terms. \square

We can use the fact that the sum of a nonnegative series is unchanged under rearrangements to study series over other sets. For example, let $\mathbb{N}^2 = \mathbb{N} \times \mathbb{N}$ be

(§1.1) the set of ordered pairs of naturals (m, n) , and set

$$(7.6) \quad \sum_{(m,n) \in \mathbb{N}^2} \frac{1}{m^3 + n^3}.$$

What do we mean by such a series? To answer this, we begin with a definition.

A set A is *countable* if there is a bijection $f : \mathbb{N} \rightarrow A$, i.e., the elements of A form a sequence. If there is no such f , we say that A is *uncountable*. Let us show that \mathbb{N}^2 is countable:

$$(1, 1), (1, 2), (2, 1), (1, 3), (2, 2), (3, 1), (1, 4), (2, 3), (3, 2), (4, 1), \dots$$

Here, we are listing the pairs (m, n) according to the sum $m + n$ of their entries. It turns out that \mathbb{Q} is countable (Exercise 2), but \mathbb{R} is not countable (Exercise 4).

Every subset of \mathbb{N} is countable or finite (Exercise 1). Thus, if $f : A \rightarrow B$ is an injection and B is countable, then, A is finite or countable. Indeed, choosing a bijection $g : B \rightarrow \mathbb{N}$ yields a bijection $g \circ f$ of A with the subset $(g \circ f)(A) \subset \mathbb{N}$.

Similarly, if $f : A \rightarrow B$ is a surjection and A is countable, then, B is countable or finite. To see this, choose a bijection $g : \mathbb{N} \rightarrow A$. Then, $f \circ g : \mathbb{N} \rightarrow B$ is a surjection, so, we may define $h : B \rightarrow \mathbb{N}$ by setting $h(b)$ equal to the least n satisfying $f[g(n)] = b$, $h(b) = \min\{n \in \mathbb{N} : f[g(n)] = b\}$. Then, h is an injection, and, thus, B is finite or countable.

Let A be a countable set. Given a positive function $f : A \rightarrow \mathbb{R}$, we define the *sum of the series over A*

$$\sum_{a \in A} f(a)$$

as the sum of $\sum_{n=1}^{\infty} f(a_n)$ obtained by taking any bijection of A with \mathbb{N} . Since the sum of a positive series is unchanged by rearrangement, *this is well defined*. As an exercise, we leave it to be shown that (7.6) converges.

Series over \mathbb{N}^2 are called *double series*. A useful arrangement of a double series follows the order of \mathbb{N}^2 displayed above,

$$\sum_{n=1}^{\infty} \left(\sum_{i+j=n+1} a_{ij} \right).$$

This is the *Cauchy order*.

Theorem. For (a_{mn}) positive,

$$(7.7) \quad \sum_{(m,n) \in \mathbb{N}^2} a_{mn} = \sum_{k=1}^{\infty} \left(\sum_{i+j=k+1} a_{ij} \right) = \sum_{n=1}^{\infty} \left(\sum_{m=1}^{\infty} a_{mn} \right) = \sum_{m=1}^{\infty} \left(\sum_{n=1}^{\infty} a_{mn} \right).$$

To see this, recall that the first equality is due to the fact that a positive double series may be summed in any order. Since the third and fourth sums are similar, it is enough to derive the second equality. To this end, note that for any natural K , the set $A_K \subset \mathbb{N}^2$ of pairs (i, j) with $i + j \leq K + 1$ is contained in the set

$B_{MN} \subset \mathbb{N}^2$ of pairs (m, n) with $m \leq M, n \leq N$, for N, M large enough (Figure 1.4). Hence,

$$\sum_{k=1}^K \left(\sum_{i+j=k+1} a_{ij} \right) \leq \sum_{n=1}^N \left(\sum_{m=1}^M a_{mn} \right) \leq \sum_{n=1}^{\infty} \left(\sum_{m=1}^{\infty} a_{mn} \right).$$

Letting $K \nearrow \infty$, we obtain

$$\sum_{k=1}^{\infty} \left(\sum_{i+j=k+1} a_{ij} \right) \leq \sum_{n=1}^{\infty} \left(\sum_{m=1}^{\infty} a_{mn} \right).$$

Conversely, for any $N, M, B_{MN} \subset A_K$ for K large enough, hence,

$$\sum_{n=1}^N \left(\sum_{m=1}^M a_{mn} \right) \leq \sum_{k=1}^K \left(\sum_{i+j=k+1} a_{ij} \right) \leq \sum_{k=1}^{\infty} \left(\sum_{i+j=k+1} a_{ij} \right).$$

Letting $M \nearrow \infty$,

$$\sum_{k=1}^{\infty} \left(\sum_{i+j=k+1} a_{ij} \right) \geq \sum_{n=1}^N \left(\sum_{m=1}^{\infty} a_{mn} \right).$$

Letting $N \nearrow \infty$,

$$\sum_{k=1}^{\infty} \left(\sum_{i+j=k+1} a_{ij} \right) \geq \sum_{n=1}^{\infty} \left(\sum_{m=1}^{\infty} a_{mn} \right).$$

This yields (7.7). \square




FIGURE 1.4. The sets B_{MN} and A_K .

To give an application of this, note that, since $\sum 1/n^{1+1/N}$ converges, by comparison, so does

$$Z(s) = \sum_{n=2}^{\infty} \frac{1}{n^s}, \quad s > 1.$$

(In the next chapter, we will know what n^s means for s real. Now, think of s as rational.) Then, (7.7) can be used to show that

$$(7.8) \quad \sum_{n=2}^{\infty} \frac{1}{n^s - 1} = Z(s) + Z(2s) + Z(3s) + \dots$$

As another application, we describe the product of two series $\sum a_n$ and $\sum b_n$. Given (7.7), it is reasonable to define their *product* or *Cauchy product* as the series

$$\sum_{n=1}^{\infty} c_n = \sum_{n=1}^{\infty} \left(\sum_{i+j=n+1} a_i b_j \right).$$

For example, $c_1 = a_1 b_1$, $c_2 = a_1 b_2 + a_2 b_1$, $c_3 = a_1 b_3 + a_2 b_2 + a_3 b_1$, and

$$c_n = a_1 b_n + a_2 b_{n-1} + \dots + a_n b_1, \quad n \geq 1.$$

Then, (7.7) shows that the Cauchy product sums to ab if both series are nonnegative and $\sum a_n = a$ and $\sum b_n = b$. It turns out this is also true for absolutely convergent signed series: If $\sum a_n$ and $\sum b_n$ converge absolutely, then, their Cauchy product converges absolutely to the product of their sums (Exercise 7).

If $a_1 + a_2 + a_3 + \dots$ is absolutely convergent, its *alternating version* is $a_1 - a_2 + a_3 - \dots$. For example the alternating version of

$$\frac{1}{1-x} = 1 + x + x^2 + \dots$$

equals

$$\frac{1}{1+x} = 1 - x + x^2 - \dots$$

Clearly, the alternating version is also absolutely convergent and the alternating version of the alternating version of a series is itself. Note that the alternating version of a series $\sum a_n$ need not be an alternating series. This happens iff $\sum a_n$ is positive.

Exercises 1.7

1. If $A \subset B$ and B is countable, then, A is countable or finite. (If $B = \mathbb{N}$, look at the smallest element in A , then, the next smallest, and so on.)
2. Show that \mathbb{Q} is countable.
3. If A_1, A_2, \dots is a sequence of countable sets, then, $\bigcup_{n=1}^{\infty} A_n$ is countable. Conclude that $\mathbb{Q} \times \mathbb{Q}$ is countable.
4. Show that $[0, 1]$ and \mathbb{R} are not countable. (Assume $[0, 1]$ is countable. List the elements as a_1, a_2, \dots . Using the decimal expansions of a_1, a_2, \dots , construct a decimal expansion not in the list.)
5. Show that (7.6) converges.

6. Derive (7.8).

7. If $\sum a_n$ and $\sum b_n$ converge absolutely, then, the Cauchy product of $\sum a_n$ and $\sum b_n$ converges absolutely to the product $(\sum a_n)(\sum b_n)$.

8. Let $\sum a_n$ and $\sum b_n$ be absolutely convergent. Then, the product of the alternating versions of $\sum a_n$ and $\sum b_n$ is the alternating version of the product of $\sum a_n$ and $\sum b_n$.




FIGURE 1.5. The golden mean.

9. ★ Given a sequence (q_n) of naturals, let x_n be as in Exercise 13 of §1.5. Show that (x_n) is Cauchy, hence, convergent to an irrational x . Thus, continued fractions yield a bijection between sequences of naturals and irrationals in $(0, 1)$. From this point of view, the continued fraction (Figure 1.5)

$$\frac{1 + \sqrt{5}}{2} = 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}}}}$$

is special. This is the *golden mean*.

Continuity

§2.1 Compactness in \mathbb{R} and \mathbb{R}^2

An *open interval* is a set of reals of the form $(a, b) = \{x : a < x < b\}$. As in §1.4, we are allowing $a = -\infty$ or $b = \infty$ or both. A *compact interval* is a set of reals of the form $[a, b] = \{x : a \leq x \leq b\}$, where a, b are real. The *length* of $[a, b]$ is $b - a$. Recall (§1.5) that a sequence *subconverges* to L if it has a subsequence converging to L .

Theorem. *Let $[a, b]$ be a compact interval and let (x_n) be any sequence in $[a, b]$. Then, (x_n) subconverges to some x in $[a, b]$.*

To derive this result, assume, first, that $a = 0$ and $b = 1$. Divide the interval $I = [a, b] = [0, 1]$ into 10 subintervals (of the same length), and call them I_0, \dots, I_9 , ordering them from left to right (Figure 2.1). Pick one of them, say I_{d_1} , containing infinitely many terms of (x_n) , i.e., $\{n : x_n \in I_{d_1}\}$ is infinite, and pick one of the terms of the sequence in I_{d_1} and call it x'_1 . Then, the length of I_{d_1} is $1/10$. Now, divide I_{d_1} into 10 subintervals again ordered left to right and called $I_{d_1 0}, \dots, I_{d_1 9}$. Pick one of them, say $I_{d_1 d_2}$, containing infinitely many terms of the sequence, and pick one of the terms (beyond x'_1) in the sequence in $I_{d_1 d_2}$ and call it x'_2 . The length of $I_{d_1 d_2}$ is $1/100$. Continuing in this manner yields $I \supset I_{d_1} \supset I_{d_1 d_2} \supset I_{d_1 d_2 d_3} \supset \dots$ and a subsequence (x'_n) where the length of $I_{d_1 d_2 \dots d_n}$ is 10^{-n} and $x'_n \in I_{d_1 d_2 \dots d_n}$ for all $n \geq 1$. But, by construction, the real

$$x = .d_1 d_2 d_3 \dots$$

lies in all the intervals $I_{d_1 d_2 \dots d_n}$, $n \geq 1$. Hence, $|x'_n - x| \leq 10^{-n} \rightarrow 0$. Since (x'_n) is a subsequence of (x_n) , this derives the result if $[a, b] = [0, 1]$. If this is not so, the same argument works. The only difference is that the limiting point now obtained is $a + x(b - a)$. \square



FIGURE 2.1. The intervals $I_{d_1 d_2 \dots d_n}$.

Thus, this theorem is equivalent to, more or less, the existence of decimal expansions. If $[a, b]$ is replaced by an open interval (a, b) , the theorem is false as it stands, since the limiting point x may be one of the endpoints, and, hence, the theorem needs to be modified. A useful modification is the following.

Theorem. *If (x_n) is any sequence of reals, then, (x_n) subconverges to some real x or to ∞ or to $-\infty$.*

To see this, consider the two cases (x_n) bounded and (x_n) unbounded. In the first case, we can find reals a, b with $(x_n) \subset [a, b]$. Hence, by the previous theorem, we obtain the subconvergence to some real $x \in [a, b]$. In the second case, if (x_n) is not bounded above, for each $n \geq 1$, choose x'_n satisfying $x'_n > n$. If (x_n) is not bounded below, for each $n \geq 1$, choose x'_n satisfying $x'_n < -n$. Then, (x'_n) converges to ∞ or to $-\infty$, yielding the subconvergence of (x_n) to ∞ or to $-\infty$. \square

We will need a related result in \mathbf{R}^2 in §4.2.

Recall that \mathbf{R}^2 is the product $\mathbf{R}^2 = \mathbf{R} \times \mathbf{R} = \{(x, y) : x, y \in \mathbf{R}\}$. A *compact rectangle* is a set in \mathbf{R}^2 of the form $[a, b] \times [c, d] = \{(x, y) : a \leq x \leq b, c \leq y \leq d\}$, where a, b, c , and d are reals. An *open rectangle* is a set of the form $(a, b) \times (c, d) = \{(x, y) : a < x < b, c < y < d\}$ (Figure 2.2). As usual, with open rectangles, either a or c may equal $-\infty$, and either b or d may equal ∞ . The *area* of a compact rectangle $[a, b] \times [c, d]$ is $(b - a)(d - c)$.





FIGURE 2.2. A and B are rectangles, but C is not.

Theorem. *Let R be a compact rectangle and let Q_1, Q_2, Q_3, \dots be any sequence of open rectangles covering R , i.e.,*

$$R \subset \bigcup_{n=1}^{\infty} Q_n.$$

Then, there is a natural N , such that $R \subset Q_1 \cup Q_2 \cup \dots \cup Q_N$.

This theorem says that no matter what their shape or size, one never really needs infinitely many open rectangles to cover (Figure 2.3) a given compact rectangle.

FIGURE 2.3. A finite cover of R by rectangles.

A finite number always suffices. Note that the rectangles Q_i may overlap, i.e., may not be disjoint.

Here the derivation is similar to the previous. We assume, first, that $R = [0, 1] \times [0, 1]$ is a square, and we argue by contradiction: Suppose that there is no finite subcollection of (Q_n) covering R . Divide R into $100 = 10 \cdot 10$ subsquares (of the same area) R_{00}, \dots, R_{99} . Here, the subsquares are ordered from left to right and bottom to top, i.e., $d < d'$ implies R_{de} is to the left of $R_{d'e}$, and $e < e'$ implies R_{de} is below $R_{de'}$. Since there are finitely many subsquares, there is at least one subsquare, call it $R_{d_1e_1}$, that is not covered by any finite subcollection of (Q_n) . Divide $R_{d_1e_1}$ into 100 subsquares, ordered as above. At least one of them, call it $R_{d_1e_1d_2e_2}$, is not covered by any finite subcollection of (Q_n) . Continuing in this manner, we obtain $R \supset R_{d_1e_1} \supset R_{d_1e_1d_2e_2} \supset \dots$ where, for each $m \geq 1$, $R_{d_1e_1d_2e_2\dots d_me_m}$ has area 100^{-m} and is not covered by any finite subcollection of (Q_n) . Let x and y be the reals

$$x = .d_1d_2\dots, \quad y = .e_1e_2\dots$$

Then, (x, y) lies in all the squares $R_{d_1e_1d_2e_2\dots d_me_m}$, $m \geq 1$. Since, in particular, (x, y) lies in R , there is at least one rectangle Q_i containing (x, y) . Since Q_i is open, (x, y) lies in the interior of Q_i and not on any of its sides. Since the dimensions of $R_{d_1e_1d_2e_2\dots d_me_m}$ approach zero as $m \nearrow \infty$, we conclude that for m large enough we have $R_{d_1e_1d_2e_2\dots d_me_m} \subset Q_i$. But this shows that $R_{d_1e_1d_2e_2\dots d_me_m}$ can be covered by one, hence, a finite subcollection of (Q_n) , contradicting the choice of $R_{d_1e_1d_2e_2\dots d_me_m}$. Thus, our initial assumption must be false, i.e., we conclude that there is a finite subcollection Q_1, \dots, Q_N covering R . If R is not a square, the same argument works. The limiting point now obtained is $(a + (b - a)x, c + (d - c)y)$. \square

If R is replaced by an open rectangle, the argument breaks down. If any of the Q_n 's are replaced by compact rectangles, again the argument breaks down.

A rectangle should be thought of as the two-dimensional analog of an interval. The first two theorems above are stated for intervals, and the third is stated for rectangles. In fact, the results above extend to any dimension $n \geq 1$, and, then, the criteria in these theorems are equivalent manifestations of the property of *compactness*.

Exercises 2.1

1. Find a sequence of open rectangles Q_1, Q_2, \dots , covering $R = (0, 1) \times (0, 1)$, such that no finite subcollection Q_1, \dots, Q_N covers R .
2. Find a compact rectangle Q_1 and a sequence of open rectangles Q_2, Q_3, \dots , covering $R = [0, 1] \times [0, 1]$, such that no finite subcollection Q_1, \dots, Q_N covers R .
3. In the derivation of the first theorem, suppose that the intervals are chosen, at each stage, to be the leftmost interval containing infinitely many terms. In other words, suppose that I_{d_1} is the leftmost of the intervals I_j containing infinitely many terms, $I_{d_1 d_2}$ is the leftmost of the intervals $I_{d_1 j}$ containing infinitely many terms, etc. In this case, show that the limiting point obtained is x_* .

§2.2 Continuous Limits

Let (a, b) be an open interval, and let $a < c < b$. The interval (a, b) , *punctured* at c , is the set $(a, b) \setminus \{c\} = \{x : a < x < b, x \neq c\}$.

Let f be a function defined on an interval (a, b) punctured at c , $a < c < b$. We say L is the *limit* of $f(x)$ as x *approaches* c , and we write

$$\lim_{x \rightarrow c} f(x) = L$$

or $f(x) \rightarrow L$ as $x \rightarrow c$, if, for *every* sequence $(x_n) \subset (a, b)$ satisfying $x_n \neq c$ for all $n \geq 1$ and converging to c , $f(x_n) \rightarrow L$.

For example, let $f(x) = x^2$, and let $(a, b) = \mathbb{R}$. If $x_n \rightarrow c$, then (§1.5), $x_n^2 \rightarrow c^2$. This holds true no matter what sequence (x_n) is chosen, as long as $x_n \rightarrow c$. Hence, in this case, $\lim_{x \rightarrow c} f(x) = c^2$.

Going back to the general definition, suppose that f is also defined at c . Then *the value $f(c)$ has no bearing on $\lim_{x \rightarrow c} f(x)$* (Figure 2.4). For example, if $f(x) = 0$ for $x \neq 0$ and $f(0)$ is defined arbitrarily, then, $\lim_{x \rightarrow 0} f(x) = 0$. For a more dramatic example of this phenomenon, see Exercise 1.




FIGURE 2.4. The value $f(c)$ has no bearing on the limit at c .

Of course, not every function has limits. For example, set $f(x) = 1$ if $x \in \mathbb{Q}$ and $f(x) = 0$ if $x \in \mathbb{R} \setminus \mathbb{Q}$. Choose any c in $(a, b) = \mathbb{R}$. Since (§1.4) there is a rational and an irrational between any two reals, for each $n \geq 1$ we can find $r_n \in \mathbb{Q}$ and $i_n \in \mathbb{R} \setminus \mathbb{Q}$ with $c < r_n < c + 1/n$ and $c < i_n < c + 1/n$. Thus, $r_n \rightarrow c$ and $i_n \rightarrow c$, but $f(r_n) = 1$ and $f(i_n) = 0$ for all $n \geq 1$. Hence, f has no limit anywhere on \mathbb{R} .

Let (x_n) be a sequence approaching b . If $x_n < b$ for all $n \geq 1$, we write $x_n \rightarrow b-$. Let f be defined on (a, b) . We say L is the *limit* of $f(x)$ as x approaches b from the left, and we write

$$\lim_{x \rightarrow b-} f(x) = L,$$

if $x_n \rightarrow b-$ implies $f(x_n) \rightarrow L$. In this case, we also write $f(b-) = L$. If $b = \infty$, we write, instead, $\lim_{x \rightarrow \infty} f(x) = L$, $f(\infty) = L$, i.e., we drop the minus.

Let (x_n) be a sequence approaching a . If $x_n > a$ for all $n \geq 1$, we write $x_n \rightarrow a+$. Let f be defined on (a, b) . We say L is the *limit* of $f(x)$ as x approaches a from the right, and we write

$$\lim_{x \rightarrow a+} f(x) = L,$$

if $x_n \rightarrow a+$ implies $f(x_n) \rightarrow L$. In this case, we also write $f(a+) = L$. If $a = -\infty$, we write, instead, $\lim_{x \rightarrow -\infty} f(x) = L$, $f(-\infty) = L$, i.e., we drop the plus.

Of course, L above is either a real or $\pm\infty$.

Theorem. Let f be defined on an interval (a, b) punctured at c , $a < c < b$. Then, $\lim_{x \rightarrow c} f(x)$ exists and equals L iff $f(c+)$ and $f(c-)$ both exist and equal L .

If $\lim_{x \rightarrow c} f(x) = L$, then, $f(x_n) \rightarrow L$ for any sequence $x_n \rightarrow c$, whether the sequence is to the right, the left, or neither. Hence, $f(c-) = L$ and $f(c+) = L$.

Conversely, suppose that $f(c-) = f(c+) = L$ and $x_n \rightarrow c$ with $x_n \neq c$ for all $n \geq 1$. We have to show that $f(x_n) \rightarrow L$. Let f^* and f_* denote the upper and lower limits of the sequence $(f(x_n))$, and set $f_n^* = \sup\{f(x_k) : k \geq n\}$. Then, $f_n^* \searrow f^*$. Hence, for any subsequence $(f_{k_n}^*)$, we have $f_{k_n}^* \searrow f^*$. Now, we have to show that $f^* = L = f_*$. Break up the sequence (x_n) as the union of two subsequences. Let (y_n) denote the terms x_k that are greater than c , and let (z_n) denote the terms x_k that are less than c , arranged in their given order. Since $f(c+) = L$ and $y_n \rightarrow c+$, we conclude that $f(y_n) \rightarrow L$, hence, its upper sequence converges to L , $\sup_{i \geq n} f(y_i) \searrow L$. Since $f(c-) = L$ and $z_n \rightarrow c-$, we conclude that $f(z_n) \rightarrow L$, hence, its upper sequence converges to L , $\sup_{i \geq n} f(z_i) \searrow L$.

For each $m \geq 1$, let x_{k_m} denote the term in (x_n) corresponding to y_m , if the term y_m appears after the term z_m in (x_n) . Otherwise, if z_m appears after y_m , let x_{k_m} denote the term in (x_n) corresponding to z_m . Thus, for each $n \geq 1$, if $j \geq k_n$, we must have x_j equal y_i or z_i with $i \geq n$. Hence,

$$f_{k_n}^* = \sup_{j \geq k_n} f(x_j) \leq \max \left[\sup_{i \geq n} f(y_i), \sup_{i \geq n} f(z_i) \right], \quad n \geq 1.$$

Now, both sequences on the right are decreasing in $n \geq 1$ to L , and the sequence on the left decreases to f^* as $n \nearrow \infty$. Thus, $f^* \leq L$. Now, let $g = -f$. Since

$g(c+) = g(c-) = -L$, by what we have just learned, we conclude that the upper limit of $(g(x_n))$ is $\leq -L$. But the upper limit of $(g(x_n))$ equals minus the lower limit f_* of $(f(x_n))$. Hence, $f_* \geq L$, so, $f^* = f_* = L$. \square

Since continuous limits are defined in terms of limits of sequences, they enjoy the same arithmetic and ordering properties. For example,

$$\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x),$$

$$\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x).$$

These properties will be used without comment.

A function f is *increasing* (*decreasing*) if $x \leq x'$ implies $f(x) \leq f(x')$ ($f(x) \geq f(x')$, respectively), for all x, x' in the domain of f . The function f is *strictly increasing* (*strictly decreasing*) if $x < x'$ implies $f(x) < f(x')$ ($f(x) > f(x')$, respectively), for all x, x' in the domain of f . If f is increasing or decreasing, we say f is *monotone*. If f is strictly increasing or strictly decreasing, we say f is *strictly monotone*.

In the exercises, the concept of a partition (Figure 2.5) is needed. If (a, b) is an open interval, a *partition of (a, b)* is a choice of $n \geq 0$ points (x_1, x_2, \dots, x_n) in (a, b) , arranged in increasing order. When choosing a partition, we write $a = x_0 < x_1 < \dots < x_n < x_{n+1} = b$, always denoting the endpoints a and b by x_0 and x_{n+1} , respectively (even when they are infinite). When $n = 0$, we obtain the empty partition consisting of no points, i.e., just the interval (a, b) . We use the same notation for compact intervals, i.e., a *partition of $[a, b]$* , by definition, is a partition of (a, b) .




FIGURE 2.5. A partition of (a, b) .

Exercises 2.2

1. Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by setting $f(m/n) = 1/n$, for $m/n \in \mathbb{Q}$ with no common factor in m and $n > 0$, and $f(x) = 0$, $x \notin \mathbb{Q}$. Show that $\lim_{x \rightarrow c} f(x) = 0$ for all $c \in \mathbb{R}$.
2. Let f be increasing on (a, b) . Then, $f(a+)$ (exists and) equals $\inf\{f(x) : a < x < b\}$, and $f(b-)$ equals $\sup\{f(x) : a < x < b\}$.
3. If f is monotone on (a, b) , then, $f(c+)$ and $f(c-)$ exist, and $f(c)$ is between $f(c-)$ and $f(c+)$, for all $c \in (a, b)$. Show also that, for each $\delta > 0$, there are, at most, countably many points $c \in (a, b)$ where $|f(c+) - f(c-)| \geq \delta$. Conclude that there are, at most, countably many points c in (a, b) at which $f(c+) \neq f(c-)$.
4. If $f : (a, b) \rightarrow \mathbb{R}$ let I_n be the sup of the sums

$$(2.1) \quad |f(x_2) - f(x_1)| + |f(x_3) - f(x_2)| + \dots + |f(x_n) - f(x_{n-1})|$$

over all partitions $a = x_0 < x_1 < x_2 < \cdots < x_n < x_{n+1} = b$ of (a, b) consisting of n points, and let $I = \sup\{I_n : n \geq 2\}$. We say that f is of *bounded variation* on (a, b) if I is finite. Show that bounded variation on (a, b) implies bounded on (a, b) . The sum in (2.1) is the *variation* of f corresponding to the partition $a < x_1 < x_2 < \cdots < x_n < b$, whereas I , the sup of all such sums over all partitions consisting of arbitrarily many points, is the *total variation* of f over (a, b) .

5. If f is bounded increasing on an interval (a, b) , then, f is of bounded variation on (a, b) . If $f = g - h$ with g, h bounded increasing on (a, b) , then, f is of bounded variation on (a, b) .

6. Let f be of bounded variation on (a, b) , and, for $a < x < b$, let $v(x)$ denote the sup of the sums (2.1) over all partitions $a = x_0 < x_1 < x_2 < \cdots < x_n < x_{n+1} = b$ with $x_n = x$ fixed. Show that $a < x < y < b$ implies $v(x) + |f(y) - f(x)| \leq v(y)$, hence, $v : (a, b) \rightarrow \mathbb{R}$ and $v - f : (a, b) \rightarrow \mathbb{R}$ are bounded increasing. Conclude that f is of bounded variation iff f is the difference of two, bounded increasing functions.

7. Show that the f in Exercise 1 is not of bounded variation on $(0, 2)$ (remember that $\sum 1/n = \infty$).

§2.3 Continuous Functions

Let f be defined on (a, b) , and choose $a < c < b$. We say that f is *continuous at c* if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

If f is continuous at every real c in (a, b) , then, we say that f is *continuous on (a, b)* or, if (a, b) is understood from the context, f is *continuous*.

Recalling the definition of $\lim_{x \rightarrow c}$, we see that f is continuous at c iff, for all sequences (x_n) satisfying $x_n \rightarrow c$ and $x_n \neq c$, $n \geq 1$, $f(x_n) \rightarrow f(c)$. In fact, f is continuous at c iff $x_n \rightarrow c$ implies $f(x_n) \rightarrow f(c)$, i.e., the condition $x_n \neq c$, $n \geq 1$, is superfluous. To see this, suppose that f is continuous at c , and suppose that $x_n \rightarrow c$, but $f(x_n) \not\rightarrow f(c)$. Since $f(x_n) \not\rightarrow f(c)$, by Exercise 8 of §1.5, there is an $\epsilon > 0$ and a subsequence (x'_n) , such that $|f(x'_n) - f(c)| \geq \epsilon$ and $x'_n \rightarrow c$, for $n \geq 1$. But, then, $f(x'_n) \neq f(c)$ for all $n \geq 1$, hence, $x'_n \neq c$ for all $n \geq 1$. Since $x'_n \rightarrow c$, by the continuity at c , we obtain $f(x'_n) \rightarrow f(c)$, contradicting $|f(x'_n) - f(c)| \geq \epsilon$. Thus, f is continuous at c iff $x_n \rightarrow c$ implies $f(x_n) \rightarrow f(c)$.

In the previous section we saw that $f(x) = x^2$ is continuous at c . Since this works for any c , f is continuous. Repeating this argument, one can show that $f(x) = x^4$ is continuous, since $x^4 = x^2 x^2$. A simpler example is to choose a real k and to set $f(x) = k$ for all x . Here, $f(x_n) = k$, and $f(c) = k$ for all sequences (x_n) and all c , so, f is continuous. Another example is $f : (0, \infty) \rightarrow \mathbb{R}$ given by

$f(x) = 1/x$. By the division property of sequences, $x_n \rightarrow c$ implies $1/x_n \rightarrow 1/c$ for $c > 0$, so, f is continuous.

Functions can be continuous at various points and not continuous at other points. For example, the function f in Exercise 1 of §2.2 is continuous at every irrational c and not continuous at every rational c . On the other hand, the function $f : \mathbf{R} \rightarrow \mathbf{R}$, given by (§2.2),

$$f(x) = \begin{cases} 1, & x \in \mathbf{Q} \\ 0, & x \notin \mathbf{Q}, \end{cases}$$

is continuous at no point.

Continuous functions have very simple arithmetic and ordering properties. If f and g are defined on (a, b) and k is real, we have functions $f + g$, kf , fg , $\max(f, g)$, $\min(f, g)$ defined on (a, b) by setting, for $a < x < b$,

$$(f + g)(x) = f(x) + g(x),$$

$$(kf)(x) = kf(x),$$

$$(fg)(x) = f(x)g(x),$$

$$\max(f, g)(x) = \max[f(x), g(x)],$$

$$\min(f, g)(x) = \min[f(x), g(x)].$$

If g is *nonzero* on (a, b) , i.e., $g(x) \neq 0$ for all $a < x < b$, define f/g by setting

$$(f/g)(x) = \frac{f(x)}{g(x)}, \quad a < x < b.$$

Theorem. *If f and g are continuous, then, so are $f + g$, kf , fg , $\max(f, g)$, and $\min(f, g)$. Moreover, if g is nonzero, then, f/g is continuous.*

This is an immediate consequence of the arithmetic and ordering properties of sequences: If $a < c < b$ and $x_n \rightarrow c$, then, $f(x_n) \rightarrow f(c)$, and $g(x_n) \rightarrow g(c)$. Hence, $f(x_n) + g(x_n) \rightarrow f(c) + g(c)$, $kf(x_n) \rightarrow kf(c)$, $f(x_n)g(x_n) \rightarrow f(c)g(c)$, $\max[f(x_n), g(x_n)] \rightarrow \max[f(c), g(c)]$, and $\min[f(x_n), g(x_n)] \rightarrow \min[f(c), g(c)]$. If $g(c) \neq 0$, then, $f(x_n)/g(x_n) \rightarrow f(c)/g(c)$. \square

For example, we see immediately that $f(x) = |x|$ is continuous on \mathbf{R} since $|x| = \max(x, -x)$.

Let us prove, by induction, that, for all $k \geq 1$, the *monomials* $f_k(x) = x^k$ are continuous (on \mathbf{R}). For $k = 1$, this is so since $x_n \rightarrow c$ implies $f_1(x_n) = x_n \rightarrow c = f_1(c)$. Assuming that this is true for k , $f_{k+1} = f_k f_1$ since $x^{k+1} = x^k x$. Hence, the result follows from the arithmetic properties of continuous functions.

A *polynomial* $f : \mathbf{R} \rightarrow \mathbf{R}$ is a linear combination of monomials, i.e., a polynomial has the form

$$f(x) = a_0 x^d + a_1 x^{d-1} + a_2 x^{d-2} + \cdots + a_{d-1} x + a_d.$$

If $a_0 \neq 0$, we call d the *degree* of f . The reals a_0, a_1, \dots, a_d , are the *coefficients* of the polynomial.

Let f be a polynomial of degree $d > 0$, and let $a \in \mathbb{R}$. Then, there is a polynomial g of degree $d - 1$ satisfying¹⁰

$$(3.1) \quad \frac{f(x) - f(a)}{x - a} = g(x), \quad x \neq a.$$

To see this, since every polynomial is a linear combination of monomials, it is enough to check (3.1) on monomials. But, for $f(x) = x^n$,

$$(3.2) \quad \frac{x^n - a^n}{x - a} = x^{n-1} + x^{n-2}a + \cdots + xa^{n-2} + a^{n-1}, \quad x \neq a,$$

which can be checked¹¹ by cross-multiplying. This establishes (3.1).

Since a monomial is continuous and a polynomial is a linear combination of monomials, by induction on the degree, we obtain the following.

Theorem. *Every polynomial f is continuous on \mathbb{R} . Moreover, if d is its degree, there are, at most, d real numbers x satisfying $f(x) = 0$.*

A real x satisfying $f(x) = 0$ is called a *zero* or a *root* of f . Thus, every polynomial f has, at most, d roots. To see this, proceed by induction on the degree of f . If $d = 1$, $f(x) = a_0x + a_1$, so, f has one root $x = -a_1/a_0$. Now, suppose that every d th degree polynomial has, at most, d roots, and let f be a polynomial of degree $d + 1$. We have to show that the number of roots of f is at most $d + 1$. If f has no roots, we are done. Otherwise, let a be a root, $f(a) = 0$. Then, by (3.1) there is a polynomial g of degree d such that $f(x) = (x - a)g(x)$. Thus, any root $b \neq a$ of f must satisfy $g(b) = 0$. Since by the inductive hypothesis, g has, at most, d roots, we see that f has, at most, $d + 1$ roots. \square

A polynomial may have no roots, e.g., $f(x) = x^2 + 1$. However, every polynomial of odd degree has at least one root (Exercise 1).

A *rational function* is a quotient $f = p/q$ of two polynomials. The natural domain of f is $\mathbb{R} \setminus Z(q)$, where $Z(q)$ denotes the set of roots of q . Since $Z(q)$ is a finite set, the natural domain of f is a finite union of open intervals. We conclude that *every rational function is continuous where it is defined*.

Let $f : (a, b) \rightarrow \mathbb{R}$. If f is not continuous at $c \in (a, b)$, we say that f is *discontinuous* at c . There are “mild” discontinuities, and there are “wild” discontinuities. The mildest situation (Figure 2.4) is when the limits $f(c+)$ and $f(c-)$ exist and are equal, but not equal to $f(c)$. This can be easily remedied by modifying the value of $f(c)$ to equal $f(c+) = f(c-)$. With this modification, the resulting function, then, is continuous at c . Because of this, such a point c is called a *removable discontinuity*. For example, the function f in Exercise 1 of §2.2 has removable discontinuities at every rational.

The next level of complexity is when $f(c+)$ and $f(c-)$ exist but may or may not be equal. In this case, we say that f has a *jump discontinuity* (Figure 2.6) or

¹⁰ g depends on a .

¹¹(3.2) with $x = 1$ was used to sum the geometric series in §1.6.

a *mild discontinuity* at c . For example, every monotone function has (at worst) jump discontinuities. In fact, every function of bounded variation has (at worst) jump discontinuities (Exercise 18). The (amount of) *jump* at c , a real number, is $f(c+) - f(c-)$. In particular, a jump discontinuity of jump zero is nothing more than a removable discontinuity.




FIGURE 2.6. A jump of 1 at each integer.

Any discontinuity that is not a jump is called a *wild discontinuity* (Figure 2.7). If f has a wild discontinuity at c , then, from above, f cannot be of bounded variation on any open interval surrounding c . The converse of this statement is false. It is possible for f to have mild discontinuities but not be of bounded variation (Exercise 7 of §2.2).




FIGURE 2.7. A wild discontinuity.

An alternate and useful description of continuity is in terms of a modulus of continuity. Let $f : (a, b) \rightarrow \mathbb{R}$, and fix $a < c < b$. For $\delta > 0$, let

$$\mu_c(\delta) = \sup\{|f(x) - f(c)| : |x - c| < \delta, a < x < b\}.$$

Since the sup, here, is possibly that of an unbounded set, we may have $\mu_c(\delta) = \infty$. The function $\mu_c : (0, \infty) \rightarrow [0, \infty) \cup \{\infty\}$ is the *modulus of continuity of f at c* (Figure 2.8).

For example, let $f : (1, 10) \rightarrow \mathbb{R}$ be given by $f(x) = x^2$ and pick $c = 9$. Since x^2 is monotone over any interval not containing zero, the maximum value of $|x^2 - 81|$ over any interval not containing zero is obtained by plugging in the endpoints. Hence, $\mu_9(\delta)$ is obtained by plugging in $x = 9 \pm \delta$, leading to $\mu_9(\delta) = \delta(\delta + 18)$. In fact, this is correct only if $0 < \delta \leq 1$. If $1 \leq \delta \leq 8$, the interval under consideration is $(9-\delta, 9+\delta) \cap (1, 10) = (9-\delta, 10)$. Here, plugging

in the endpoints leads to $\mu_9(\delta) = \max(19, 18\delta - \delta^2)$. If $\delta \geq 8$, then, $(9 - \delta, 9 + \delta)$ contains $(1, 10)$ and, hence, $\mu_9(\delta) = 80$. Summarizing, for $f(x) = x^2$, $c = 9$, and $(a, b) = (1, 10)$,

$$\mu_c(\delta) = \begin{cases} \delta(\delta + 18), & 0 < \delta \leq 1, \\ \max(19, 18\delta - \delta^2), & 1 \leq \delta \leq 8, \\ 80, & \delta \geq 8. \end{cases}$$




FIGURE 2.8. Computing the modulus of continuity.

Going back to the general definition, note that $\mu_c(\delta)$ is an increasing function of δ , and, hence, $\mu_c(0+)$ exists (Exercise 2 of §2.2).

Theorem. Let $f : (a, b) \rightarrow \mathbb{R}$, and choose $c \in (a, b)$. The following are equivalent:

- (1) f is continuous at c .
- (2) $\mu_c(0+) = 0$.
- (3) For all $\epsilon > 0$, there exists $\delta > 0$, such that $|x - c| < \delta$ implies $|f(x) - f(c)| < \epsilon$.

That (1) implies (2) is left as Exercise 2. Now, assume (2), and suppose that $\epsilon > 0$ is given. Since $\mu_c(0+) = 0$, there exists a $\delta > 0$ with $\mu_c(\delta) < \epsilon$. Then, by definition of μ_c , $|x - c| < \delta$ implies $|f(x) - f(c)| \leq \mu_c(\delta) < \epsilon$, which establishes (3). Now, assume the ϵ - δ criterion (3), and let $x_n \rightarrow c$. Then, for all but a finite number of terms, $|x_n - c| < \delta$. Hence, for all but a finite number of terms, $f(c) - \epsilon < f(x_n) < f(c) + \epsilon$. Let $y_n = f(x_n)$, $n \geq 1$. By the ordering properties of sup and inf, $f(c) - \epsilon \leq y_{n*} \leq y_n^* \leq f(c) + \epsilon$. By the ordering properties of sequences, $f(c) - \epsilon \leq y_* \leq y^* \leq f(c) + \epsilon$. Since $\epsilon > 0$ is arbitrary, $y^* = y_* = f(c)$. Thus, $y_n = f(x_n) \rightarrow f(c)$. Since (x_n) was any sequence converging to c , $\lim_{x \rightarrow c} f(x) = f(c)$, i.e., (1). \square

Thus, in practice, one needs to compute $\mu_c(\delta)$ only for δ small enough, since it is the behavior of μ_c near zero that counts. For example, to check continuity of $f(x) = x^2$ at $c = 9$, it is enough to note that $\mu_9(\delta) = \delta(\delta + 18)$ for small enough δ , which clearly approaches zero as $\delta \rightarrow 0+$.

To check the continuity of $f(x) = x^2$ at $c = 9$ using the ϵ - δ criterion (3), given $\epsilon > 0$, it is enough to exhibit a $\delta > 0$ with $\mu_9(\delta) < \epsilon$. Such a δ is the lesser of

$\epsilon/20$ and 1 , $\delta = \min(\epsilon/20, 1)$. To see this, first, note that $\delta(\delta + 18) \leq 19$ for this δ . Then, $\epsilon \leq 19$ implies $\delta(\delta + 18) \leq (\epsilon/20)(1 + 18) = (19/20)\epsilon < \epsilon$, whereas $\epsilon > 19$ implies $\delta(\delta + 18) < \epsilon$. Hence, in either case, $\mu_9(\delta) < \epsilon$, establishing (3).

We never use the ϵ - δ criterion (3) in this text. It is included here merely for historical reasons. In practice, the definition (1) or the criterion (2) are more efficient.

Now, we turn to the mapping properties of a continuous function. First, we define one-sided continuity. Let f be defined on $(a, b]$. We say that f is *continuous at b from the left* if $f(b-) = f(b)$. In addition, if f is continuous on (a, b) , we say that f is *continuous on (a, b)* . Let f be defined on $[a, b)$. We say that f is *continuous at a from the right* if $f(a+) = f(a)$. In addition, if f is continuous on (a, b) , we say that f is *continuous on $[a, b)$* .

Note that a function f is continuous at a particular point c iff f is continuous at c from the right and continuous at c from the left.

Let f be defined on $[a, b]$. We say that f is *continuous on $[a, b]$* if f is continuous on $[a, b)$ and $(a, b]$. Checking the definitions, we see f is continuous on A if, for every $c \in A$ and every sequence $(x_n) \subset A$ converging to c , $f(x_n) \rightarrow f(c)$, whether A is (a, b) , $(a, b]$, $[a, b)$, or $[a, b]$.

Theorem. *Let f be continuous on a compact interval $[a, b]$. Then, $f([a, b])$ is a compact interval $[m, M]$.*

Thus, a continuous function maps compact intervals to compact intervals. Of course, $f([a, b])$ may not equal $[f(a), f(b)]$. For example, if $f(x) = x^2$, $f([-2, 2]) = [0, 4]$ and $[f(-2), f(2)] = \{4\}$. We derive two consequences of this theorem.

Let $f([a, b]) = [m, M]$. Then, we have two reals c and d in $[a, b]$, such that $f(c) = m$ and $f(d) = M$. In other words, M is a max, and m is a min for the set $f([a, b])$. Thus, *a continuous function on a compact interval admits a max and a min*. Of course, this is not generally true on noncompact intervals since $f(x) = 1/x$ has no max on $(0, 1]$.

A second consequence is: Suppose that L is an intermediate value between $f(a)$ and $f(b)$. Then, there must be a c , $a < c < b$, satisfying $f(c) = L$. This follows since $f(a)$ and $f(b)$ are two reals in $f([a, b])$, and $f([a, b])$ is an interval. Thus, *a continuous function on a compact interval attains every intermediate value*. This is the *intermediate value property*.

On the other hand, the two consequences, the existence of the max and the min and the intermediate value property, combine to yield the theorem. To see this, let $m = f(c)$ and $M = f(d)$ denote the max and the min, with $c, d \in [a, b]$. If $m = M$, f is constant, hence, $f([a, b]) = [m, M]$. If $m < M$ and $m < L < M$, apply the intermediate value property to conclude that there is an x between c and d with $f(x) = L$. Hence, $f([a, b]) = [m, M]$. Thus, to derive the theorem, it is enough to derive the two consequences.

For the first, let $M = \sup\{f(x) : a \leq x \leq b\}$. If $M < \infty$, for all $n \geq 1$, we choose $x_n \in [a, b]$ satisfying $f(x_n) > M - 1/n$. If $M = \infty$, for all $n \geq 1$, we

choose $x_n \in [a, b]$ satisfying $f(x_n) > n$. In either case, we obtain a sequence (x_n) with $f(x_n) \rightarrow M$. But (§2.1) (x_n) subconverges to some $c \in [a, b]$. By continuity, $(f(x_n))$ subconverges to $f(c)$. Since $(f(x_n))$ also converges to M , $M = f(c)$, so, f has a max. Applying this to $g = -f$, we see that g has a max which implies f has a min.

For the second, suppose that $f(a) < f(b)$, and let L be an intermediate value, $f(a) < L < f(b)$. We proceed as in the construction of $\sqrt{2}$ in §1.4. Let $S = \{x \in [a, b] : f(x) < L\}$, and let $c = \sup S$. S is nonempty since $a \in S$, and S is clearly bounded. For all $n \geq 1$, $c - 1/n$ is not an upper bound for S . Hence, for each $n \geq 1$, there is a real $x_n \in S$ with $c \geq x_n > c - 1/n$, which gives $x_n \rightarrow c$. By continuity, $f(x_n) \rightarrow f(c)$. Since $f(x_n) < L$ for all $n \geq 1$, we obtain $f(c) \leq L$. On the other hand, $c + 1/n$ is not in S , hence, $f(c + 1/n) \geq L$. Since $c + 1/n \rightarrow c$, we obtain $f(c) \geq L$. Thus, $f(c) = L$. The case $f(a) > f(b)$ is similar or is established by applying the previous to $-f$. \square

From this theorem, it follows that a continuous function maps open intervals to intervals. However, they need not be open. For example, with $f(x) = x^2$, $f((-2, 2)) = [0, 4)$. However, a function that is continuous and strictly monotone maps open intervals to open intervals (Exercise 3).

The above theorem is the result of compactness mixed with continuity. This mixture yields other surprises. Let $f : (a, b) \rightarrow \mathbb{R}$ be given, and fix a subset $A \subset (a, b)$. For $\delta > 0$, set

$$\mu_A(\delta) = \sup\{\mu_c(\delta) : c \in A\}.$$

This is the *uniform modulus of continuity of f on A* . Since $\mu_c(\delta)$ is a decreasing function of δ for each $c \in A$, it follows that $\mu_A(\delta)$ is a decreasing function of δ , and hence, $\mu_A(0+)$ exists. We say f is *uniformly continuous on A* if $\mu_A(0+) = 0$. When $A = (a, b)$ equals the whole domain of the function, we delete the subscript A and write $\mu(\delta)$ for the uniform modulus of continuity of f on its domain.

Whereas continuity is a property pertaining to the behavior of a function at (or near) a given point c , uniform continuity is a property pertaining to the behavior of f near a given set A . Moreover, since $\mu_c(\delta) \leq \mu_A(\delta)$, uniform continuity on A implies continuity at every point $c \in A$.

Inserting the definition of $\mu_c(\delta)$ in $\mu_A(\delta)$ yields

$$\mu_A(\delta) = \sup\{|f(x) - f(c)| : |x - c| < \delta, a < x < b, c \in A\},$$

where, now, the sup is over both x and c .

For example, for $f(x) = x^2$, the uniform modulus $\mu_A(\delta)$ over $A = (1, 10)$ equals the sup of $|x^2 - y^2|$ over all $1 < x < y < 10$ with $y - x < \delta$. But this is largest when $y = x + \delta$, hence, $\mu_A(\delta)$ is the sup of $\delta^2 + 2x\delta$ over $1 < x < 10 - \delta$ which yields $\mu_A(\delta) = 20\delta - \delta^2$. In fact, this is correct only if $0 < \delta \leq 9$. For $\delta = 9$, the sup is already over all of $(1, 10)$, hence, cannot get any bigger. Hence, $\mu_A(\delta) = 99$ for $\delta \geq 9$. Summarizing, for $f(x) = x^2$ and $A = (1, 10)$,

$$\mu_A(\delta) = \begin{cases} 20\delta - \delta^2, & 0 < \delta \leq 9, \\ 99, & \delta \geq 9. \end{cases}$$

Since f is uniformly continuous on A if $\mu_A(0+) = 0$, in practice one needs to compute $\mu_A(\delta)$ only for δ small enough. For example, to check uniform continuity of $f(x) = x^2$ over $A = (1, 10)$, it is enough to note that $\mu_A(\delta) = 20\delta - \delta^2$ for small enough δ , which clearly approaches zero as $\delta \rightarrow 0+$.

Now, let $f : (a, b) \rightarrow \mathbf{R}$ be continuous, and fix $A \subset (a, b)$. What additional conditions on f are needed to guarantee uniform continuity on A ? When A is a finite set $\{c_1, \dots, c_N\}$,

$$\mu_A(\delta) = \max [\mu_{c_1}(\delta), \dots, \mu_{c_N}(\delta)],$$

and, hence, f is necessarily uniformly continuous on A .

When A is an infinite set, this need not be so. For example, with $f(x) = x^2$ and $B = (0, \infty)$, $\mu_B(\delta)$ equals the sup of $\mu_c(\delta) = 2c\delta + \delta^2$ over $0 < c < \infty$, or $\mu_B(\delta) = \infty$, for each $\delta > 0$. Hence, f is not uniformly continuous on B .

It turns out that continuity on a compact interval is sufficient for uniform continuity.

Theorem. *If $f : [a, b] \rightarrow \mathbf{R}$ is continuous, then, f is uniformly continuous on (a, b) .*

To see this, suppose that $\mu(0+) = \mu_{(a,b)}(0+) > 0$, and set $\epsilon = \mu(0+)/2$. Since μ is increasing, $\mu(1/n) \geq 2\epsilon$, $n \geq 1$. Hence, for each $n \geq 1$, by the definition of the sup in the definition of $\mu(1/n)$, there is a $c_n \in (a, b)$ with $\mu_{c_n}(1/n) > \epsilon$. Now, by the definition of the sup in $\mu_{c_n}(1/n)$, for each $n \geq 1$, there is an $x_n \in (a, b)$ with $|x_n - c_n| < 1/n$ and $|f(x_n) - f(c_n)| > \epsilon$. By compactness, (x_n) subconverges to some $x \in [a, b]$. Since $|x_n - c_n| < 1/n$ for all $n \geq 1$, (c_n) subconverges to the same x . Hence, by continuity, $(|f(x_n) - f(c_n)|)$ subconverges to $|f(x) - f(x)| = 0$, which contradicts the fact that this last sequence is bounded below by $\epsilon > 0$. \square

The conclusion may be false if f is continuous on (a, b) but not on $[a, b]$ (see Exercise 23). One way to understand the difference between continuity and uniform continuity is as follows.

Let f be a continuous function defined on an interval (a, b) , and pick $c \in (a, b)$. Then, by definition of μ_c , $|f(x) - f(c)| \leq \mu_c(\delta)$ whenever x lies in the interval $(c - \delta, c + \delta)$. Setting $g(x) = f(c)$ for $x \in (c - \delta, c + \delta)$, we see that, for any error tolerance ϵ , by choosing δ satisfying $\mu_c(\delta) < \epsilon$, we obtain a constant function g approximating f to within ϵ , at least in the interval $(c - \delta, c + \delta)$. Of course, in general, we do not expect to approximate f closely by one and the same constant function over the whole interval (a, b) . Instead, we use piecewise constant functions.

We say $g : (a, b) \rightarrow \mathbf{R}$ is *piecewise constant* if there is a partition $a = x_0 < x_1 < \dots < x_n < x_{n+1} = b$, such that g restricted to (x_{i-1}, x_i) is constant for $i = 1, \dots, n+1$ (in this definition, the values of g at the points x_i are not restricted in any way). The *mesh* δ of the partition $a = x_0 < x_1 < \dots < x_{n+1} = b$, by definition, is the largest length of the subintervals, $\delta = \max_{1 \leq i \leq n+1} |x_i - x_{i-1}|$. Note that an interval has partitions of arbitrarily small mesh iff the interval is bounded.

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, from above, f is uniformly continuous on (a, b) . Given a partition $a = x_0 < x_1 < \cdots < x_{n+1} = b$ with mesh δ , choose $x_i^\#$ in (x_{i-1}, x_i) arbitrarily, $i = 1, \dots, n+1$. Then, by definition of μ , $|f(x) - f(x_i^\#)| \leq \mu(\delta)$ for $x \in (x_{i-1}, x_i)$. If we set $g(x) = f(x_i^\#)$ for $x \in (x_{i-1}, x_i)$, $i = 1, \dots, n+1$, and $g(x_i) = f(x_i)$, $i = 0, 1, \dots, n+1$, we obtain a piecewise constant function $g : [a, b] \rightarrow \mathbb{R}$ satisfying $|f(x) - g(x)| \leq \mu(\delta)$ for every $x \in [a, b]$. Since f is uniformly continuous, $\mu(0+) = 0$. Hence, for any error tolerance $\epsilon > 0$, we can find a mesh δ , such that $\mu(\delta) < \epsilon$. We have derived the following (Figure 2.9).

Theorem. *If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then, for each $\epsilon > 0$, there is a piecewise constant function $f_\epsilon : [a, b] \rightarrow \mathbb{R}$, such that*

$$|f(x) - f_\epsilon(x)| \leq \epsilon, \quad a \leq x \leq b. \quad \square$$




FIGURE 2.9. Piecewise constant approximation.

If f is continuous on an open interval, this result may be false. For example $f(x) = 1/x$, $0 < x < 1$, cannot be approximated as above by a piecewise constant function (unless infinitely many subintervals are used), precisely because f “shoots up to ∞ ” near 0.

Let us turn to the continuity of compositions (§1.1). Suppose that $f : (a, b) \rightarrow \mathbb{R}$ and $g : (c, d) \rightarrow \mathbb{R}$ are given with the range of f lying in the domain of g , $f[(a, b)] \subset (c, d)$. Then, the composition $g \circ f : (a, b) \rightarrow \mathbb{R}$ is given by $(g \circ f)(x) = g[f(x)]$, $a < x < b$.

Theorem. *If f and g are continuous, so is $g \circ f$.*

Since f is continuous, $x_n \rightarrow c$ implies $f(x_n) \rightarrow f(c)$. Since g is continuous, $(g \circ f)(x_n) = g[f(x_n)] \rightarrow g[f(c)] = (g \circ f)(c)$. \square

This result can be written

$$\lim_{x \rightarrow c} g[f(x)] = g \left[\lim_{x \rightarrow c} f(x) \right].$$

Since $g(x) = |x|$ is continuous, this implies

$$\lim_{x \rightarrow c} |f(x)| = \left| \lim_{x \rightarrow c} f(x) \right|.$$

The final issue is the invertibility of continuous functions. Let $f : [a, b] \rightarrow [m, M]$ be a continuous function. When is there an inverse (§1.1) $g : [m, M] \rightarrow [a, b]$? If it exists, is the inverse g necessarily continuous? It turns out that the answers to these questions are related to the monotonicity properties (§2.2) of the continuous function. For example, if f is continuous and increasing on $[a, b]$ and $A \subset [a, b]$, $\sup f(A) = f(\sup A)$, and $\inf f(A) = f(\inf A)$ (Exercise 4). It follows that the upper and lower limits of $(f(x_n))$ are $f(x^*)$ and $f(x_*)$, respectively, where x^*, x_* are the upper and lower limits of (x_n) (Exercise 5).

Inverse Function Theorem. *Let $f : [a, b] \rightarrow \mathbf{R}$ be continuous. Then, f is injective iff f is strictly monotone. In this case, let $[m, M] = f([a, b])$. Then, the inverse $g : [m, M] \rightarrow [a, b]$ is continuous and strictly monotone.*

If f is strictly monotone and $x \neq x'$, then, $x < x'$ or $x > x'$ which implies $f(x) < f(x')$ or $f(x) > f(x')$, hence, f is injective.

Conversely, suppose that f is injective and $f(a) < f(b)$. We claim that f is strictly increasing (Figure 2.10). To see this, suppose not and choose $a \leq x < x' \leq b$ with $f(x) > f(x')$. There are two possibilities: Either $f(a) < f(x)$ or $f(a) \geq f(x)$. In the first case, we can choose L in $(f(a), f(x)) \cap (f(x'), f(x))$. By the intermediate value property there are c, d with $a < c < x < d < x'$ with $f(c) = L = f(d)$. Since f is injective, this cannot happen, ruling out the first case. In the second case we must have $f(x') < f(b)$, hence, $x' < b$, so, we choose L in $(f(x'), f(x)) \cap (f(x'), f(b))$. By the intermediate value property, there are c, d with $x < c < x' < d < b$ with $f(c) = L = f(d)$. Since f is injective, this cannot happen, ruling out the second case. Thus, f is strictly increasing. If $f(a) > f(b)$, applying what we just learned to $-f$ yields $-f$ strictly increasing or f strictly decreasing. Thus, in either case, f is strictly monotone.




FIGURE 2.10. Derivation of the IFT when $f(a) < f(b)$.

Clearly strict monotonicity of f implies that of g . Now, assume that f is strictly increasing, the case with f strictly decreasing being entirely similar. We have to show that g is continuous. Suppose that $(y_n) \subset [m, M]$ with $y_n \rightarrow y$. Let $x = g(y)$, let $x_n = g(y_n)$, $n \geq 1$, and let x^* and x_* denote the upper and lower limits of (x_n) . We have to show $g(y_n) = x_n \rightarrow x = g(y)$. Since f is continuous and increasing, $f(x^*)$ and $f(x_*)$ are the upper and lower limits of $y_n = f(x_n)$. Hence, $f(x^*) = y = f(x_*)$. Hence, by injectivity, $x^* = x = x_*$. \square

As an application, note that $f(x) = x^2$ is strictly increasing on $[0, n]$, hence, has an inverse $g_n(x) = \sqrt{x}$ on $[0, n^2]$, for each $n \geq 1$. By uniqueness of inverses (Exercise 4 of §1.1), the functions g_n , $n \geq 1$, agree wherever their domains overlap, hence, yield a single, continuous, strictly monotone $g : [0, \infty) \rightarrow [0, \infty)$ satisfying $g(x) = \sqrt{x}$, $x \geq 0$. Similarly, for each $n \geq 1$, $f(x) = x^n$ is strictly increasing on $[0, \infty)$. Thus, every positive real x has a unique positive n th root $x^{1/n}$, and, moreover, the function $g(x) = x^{1/n}$ is continuous on $[0, \infty)$. By composition, it follows that $f(x) = x^{m/n} = (x^m)^{1/n}$ is continuous and strictly monotone on $[0, \infty)$ for all naturals m, n . Since $x^{-a} = 1/x^a$ for $a \in \mathbb{Q}$, we see that the *power functions* $f(x) = x^r$ are defined, strictly increasing, and continuous on $(0, \infty)$ for all rationals r . Moreover, $x^{r+s} = x^r x^s$, $(x^r)^s = x^{rs}$ for r, s rational, and, for $r > 0$ rational, $x^r \rightarrow 0$ as $x \rightarrow 0$ and $x^r \rightarrow \infty$ as $x \rightarrow \infty$. The following limit is important: For $x > 0$,

$$(3.3) \quad \lim_{n \nearrow \infty} x^{1/n} = 1.$$

To derive this, assume $x \geq 1$. Then, $x \leq x x^{1/n} = x^{(n+1)/n}$, so, $x^{1/(n+1)} \leq x^{1/n}$, so, the sequence $(x^{1/n})$ is decreasing and bounded below by 1, hence, its limit $L \geq 1$ exists. Since $L \leq x^{1/2n}$, $L^2 \leq x^{2/2n} = x^{1/n}$, hence, $L^2 \leq L$ or $L \leq 1$. We conclude that $L = 1$. If $0 < x < 1$, then, $1/x > 1$, so, $x^{1/n} = 1/(1/x)^{1/n} \rightarrow 1$ as $n \nearrow \infty$.

Any function that can be obtained from polynomials or rational functions by arithmetic operations and/or the taking of roots is called a (*constructible*) *algebraic function*. For example,

$$f(x) = \frac{1}{\sqrt{x(1-x)}}, \quad 0 < x < 1,$$

is an algebraic function.

We now know what a^b means for any $a > 0$ and $b \in \mathbb{Q}$. But what if $b \notin \mathbb{Q}$? What does $2^{\sqrt{2}}$ mean? To answer this, fix $a > 1$ and $b > 0$, and let

$$c = \sup\{a^r : 0 < r < b, r \in \mathbb{Q}\}.$$

Let us check that when b is rational, $c = a^b$. Since $r < s$ implies $a^r < a^s$, $a^r \leq a^b$ when $r < b$. Hence, $c \leq a^b$. Similarly, $c \geq a^{b-1/n} = a^b/a^{1/n}$ for all $n \geq 1$. Let $n \nearrow \infty$ and use (3.3) to get $c \geq a^b$. Hence, $c = a^b$ when b is rational. Thus, it is consistent to *define*, for any $a > 1$ and *real* $b > 0$,

$$a^b = \sup\{a^r : 0 < r < b, r \in \mathbb{Q}\},$$

$a^0 = 1$, and $a^{-b} = 1/a^b$. For all b real, we define $1^b = 1$, whereas for $0 < a < 1$, we define $a^b = 1/(1/a)^b$. This defines $a^b > 0$ for all positive real a and all real b . Moreover (Exercise 7),

$$a^b = \inf\{a^s : s > b, s \in \mathbb{Q}\}.$$

Now, we show that a^b satisfies the usual rules.

Theorem.

- For $a > 1$ and $0 < b < c$ real, $1 < a^b < a^c$.
- For $0 < a < 1$ and $0 < b < c$ real, $a^b > a^c$.
- For $0 < a < b$ and $c > 0$ real, $a^c b^c = (ab)^c$, $(b/a)^c = b^c/a^c$, and $a^c < b^c$.
- For $a > 0$ and b, c real, $a^{b+c} = a^b a^c$.
- For $a > 0$, b, c real, $a^{bc} = (a^b)^c$.

Since $A \subset B$ implies $\sup A \leq \sup B$, $a^b \leq a^c$ when $a > 1$ and $b < c$. Since, for any $b < c$, there is an $r \in \mathbb{Q} \cap (b, c)$, $a^b < a^r < a^c$, thus, the first assertion. Since, for $0 < a < 1$, $a^b = 1/(1/a)^b$, applying the first assertion to $1/a$ yields $(1/a)^b < (1/a)^c$ or $a^b > a^c$, yielding the second assertion. For the third, assume $a > 1$. If $0 < r < c$ is in \mathbb{Q} , then, $a^r < a^c$ and $b^r < b^c$ yields $(ab)^r = a^r b^r < a^c b^c$. Taking the sup over $r < c$ yields $(ab)^c \leq a^c b^c$. If $r < c$ and $s < c$ are positive rationals, let t denote their max. Then, $a^r b^s \leq a^t b^t = (ab)^t < (ab)^c$. Taking the sup of this last inequality over all $0 < r < c$, first, then, over all $0 < s < c$ yields $a^c b^c \leq (ab)^c$. Hence $(ab)^c = a^c b^c$ for $b > a > 1$. Using this, we obtain $(b/a)^c a^c = b^c$ or $(b/a)^c = b^c/a^c$. Since $b/a > 1$ implies $(b/a)^c > 1$, we also obtain $a^c < b^c$. The cases $a < b < 1$ and $a < 1 < b$ follow from the case $b > a > 1$. This establishes the third. For the fourth, the case $0 < a < 1$ follows from the case $a > 1$, so, assume $a > 1$, $b > 0$, and $c > 0$. If $r < b$ and $s < c$ are positive rationals, then, $a^{b+c} \geq a^{r+s} = a^r a^s$. Taking the sups over r and s yields $a^{b+c} \geq a^b a^c$. If $r < b + c$ is rational, let $d = (b + c - r)/3 > 0$. Pick rationals t and s with $b > t > b - d$, $c > s > c - d$. Then, $t + s > b + c - 2d > r$, so, $a^r < a^{t+s} = a^t a^s \leq a^b a^c$. Taking the sup over all such r , we obtain $a^{b+c} \leq a^b a^c$. This establishes the fourth when b and c are positive. The cases $b \leq 0$ or $c \leq 0$ follow from the positive case. The fifth involves approximating b and c by rationals, and we leave it to the reader. \square

As an application, we define the power function with an irrational exponent. This is a nonalgebraic or *transcendental* function. The transcendental functions in this book are the power function x^a (when a is irrational), the exponential function a^x , the logarithm $\log_a x$, the trigonometric functions and their inverses, and the gamma function. The trigonometric functions are discussed in §3.5, the gamma function in §5.1, whereas the power, exponential, and logarithm functions are discussed below.

Theorem. *Let a be real, and let $f(x) = x^a$ on $(0, \infty)$. For $a > 0$, f is strictly increasing and continuous with $f(0+) = 0$ and $f(\infty) = \infty$. For $a < 0$, f is strictly decreasing and continuous with $f(0+) = \infty$ and $f(\infty) = 0$.*

Since $x^{-a} = 1/x^a$, the second part follows from the first, so, assume $a > 0$. Let r, s be positive rationals with $r < a < s$, and let $x_n \rightarrow c$. We have to show that $x_n^a \rightarrow c^a$. But the sequence (x_n^a) lies between (x_n^r) and (x_n^s) . Since we already know that the rational power function is continuous, we conclude that the upper and lower limits L^*, L_* , of (x_n^a) satisfy $c^r \leq L_* \leq L^* \leq c^s$. Taking the sup over all r rational and the inf over all s rational, with $r < a < s$, gives $L^* = L_* = c^a$. Thus, f is continuous. Also, since $x^r \rightarrow \infty$ as $x \rightarrow \infty$ and $x^r \leq x^a$ for $r < a$, $f(\infty) = \infty$. Since $x^a \leq x^s$ for $s > a$ and $x^s \rightarrow 0$ as $x \rightarrow 0+$, $f(0+) = 0$. \square

Now we vary b and fix a in a^b .

Theorem. *Fix $a > 1$. Then, the function $f(x) = a^x$, $x \in \mathbb{R}$, is strictly increasing and continuous. Moreover,*

$$(3.4) \quad f(x + x') = f(x)f(x'),$$

$$f(-\infty) = 0, f(0) = 1, \text{ and } f(\infty) = \infty.$$

From the previous section, we know that f is strictly increasing. Since $a^n \nearrow \infty$ as $n \nearrow \infty$, $f(\infty) = \infty$. Since $f(-x) = 1/f(x)$, $f(-\infty) = 0$. Continuity remains to be shown. If $x_n \searrow c$, then, (a^{x_n}) is decreasing and $a^{x_n} \geq a^c$, so, its limit L is $\geq a^c$. On the other hand, for $d > 0$, the sequence is eventually below $a^{c+d} = a^c a^d$, hence, $L \leq a^c a^d$. Choosing $d = 1/n$, we obtain $a^c \leq L \leq a^c a^{1/n}$. Let $n \nearrow \infty$ to get $L = a^c$. Thus, $a^{x_n} \searrow a^c$. If $x_n \rightarrow c+$ is not necessarily decreasing, then, $x_n^* \searrow c$, hence, $a^{x_n^*} \rightarrow a^c$. But $x_n^* \geq x_n$ for all $n \geq 1$, hence, $a^{x_n^*} \geq a^{x_n} \geq a^c$, so, $a^{x_n} \rightarrow a^c$. Similarly, from the left. \square

The function $f(x) = a^x$ is the *exponential function with base $a > 1$* . In fact, the exponential is the unique continuous function f on \mathbb{R} satisfying the functional equation (3.4) and $f(1) = a$.

By the inverse function theorem, f has an inverse g on any compact interval, hence, on \mathbb{R} . We call g the *logarithm with base $a > 1$* , and write $g(x) = \log_a x$. By definition of inverse, $a^{\log_a x} = x$, for $x > 0$, and $\log_a(a^x) = x$, for $x \in \mathbb{R}$.

Theorem. *The inverse of the exponential $f(x) = a^x$ with base $a > 1$ is the logarithm with base $a > 1$, $g(x) = \log_a x$. The logarithm is continuous and strictly increasing on $(0, \infty)$. The domain of \log_a is $(0, \infty)$, the range is \mathbb{R} , $\log_a(0+) = -\infty$, $\log_a 1 = 0$, $\log_a \infty = \infty$, and*

$$\log_a(bc) = \log_a b + \log_a c,$$

$$\log_a(b^c) = c \log_a b,$$

for $b > 0, c > 0$.

This follows immediately from the properties of the exponential function with base $a > 1$. \square

Exercises 2.3

1. If f is a polynomial of odd degree, then, $f(\pm\infty) = \pm\infty$ or $f(\pm\infty) = \mp\infty$, and there is at least one real c with $f(c) = 0$.
2. Show that f is continuous at c iff $\mu_c(0+) = 0$.
3. If $f : (a, b) \rightarrow \mathbb{R}$ is continuous, then, $f((a, b))$ is an interval. In addition, if f is strictly monotone, $f((a, b))$ is an open interval.
4. If f is continuous and increasing on $[a, b]$ and $A \subset [a, b]$, then, $\sup f(A) = f(\sup A)$, and $\inf f(A) = f(\inf A)$.
5. With f as in Exercise 4, let x^* and x_* be the upper and lower limits of a sequence (x_n) . Then, $f(x^*)$ and $f(x_*)$ are the upper and lower limits of $(f(x_n))$.
6. With $r, s \in \mathbb{Q}$ and $x > 0$, show that $(x^r)^s = x^{rs}$ and $x^{r+s} = x^r x^s$.
7. Show that $a^b = \inf\{a^s : s > b, s \in \mathbb{Q}\}$.
8. With b and c real and $a > 0$, show that $(a^b)^c = a^{bc}$.
9. Fix $a > 0$. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, $f(1) = a$, and $f(x+x') = f(x)f(x')$ for $x, x' \in \mathbb{R}$, then, $f(x) = a^x$.
10. Use the ϵ - δ criterion to show that $f(x) = 1/x$ is continuous at $x = 1$.
11. ★ A real x is *algebraic* if x is a root of a polynomial of degree $d \geq 1$,

$$a_0x^d + a_1x^{d-1} + \cdots + a_{d-1}x + a_d = 0,$$

with rational coefficients a_0, a_1, \dots, a_d . A real is *transcendental* if it is not algebraic. For example, every rational is algebraic. Show that the set of algebraic numbers is countable (§1.7). Conclude that the set of transcendental numbers is uncountable.

12. ★ Let a be an algebraic number. If $f(a) = 0$ for some polynomial f with rational coefficients, but $g(a) \neq 0$ for any polynomial g with rational coefficients of lesser degree, then, f is a *minimal polynomial* for a , and the degree of f is the *algebraic order* of a . Now, suppose that a is algebraic of order $d \geq 2$. Show that all the roots of a minimal polynomial f are irrational.
13. ★ Suppose that the algebraic order of a is $d \geq 2$. Then, there is a $c > 0$, such that

$$\left| a - \frac{m}{n} \right| \geq \frac{c}{n^d}, \quad n, m \geq 1.$$

(See Exercise 10 of §1.4. Here, you will need the modulus of continuity μ_a at a of $g(x) = f(x)/(x - a)$, where f is a minimal polynomial of a .)

14. ★ Use the previous exercise to show that

$$.1100010\dots 010\dots = \frac{1}{10} + \frac{1}{10^2} + \frac{1}{10^6} + \dots = \sum_{n=1}^{\infty} \frac{1}{10^{n!}}$$

is transcendental.

15. For $s > 1$ real, $\sum_{n=1}^{\infty} n^{-s}$ converges.

16. If $a > 1$, $b > 0$, and $c > 0$, then, $b^{\log_a c} = c^{\log_a b}$, and

$$\sum_{n=1}^{\infty} \frac{1}{5^{\log_3 n}}$$

converges.

17. Give an example of an $f : [0, 1] \rightarrow [0, 1]$ that is invertible but not monotone.

18. Let f be of bounded variation (Exercise 6 of §2.2) on (a, b) . Then, the set of points at which f is not continuous is at most countable. Moreover, every discontinuity, at worst, is a jump.

19. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and $f(\infty) = f(-\infty) = -\infty$, then, $\max\{f(x) : x \in \mathbb{R}\}$ exists.

20. ★ If $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies

$$\lim_{x \rightarrow \pm\infty} \frac{f(x)}{|x|} = +\infty,$$

we say that f is *superlinear*. If f is superlinear and continuous, then

$$g(y) = \max_{-\infty < x < \infty} [xy - f(x)], \quad y \in \mathbb{R},$$

is well defined (the max exists), and g is superlinear. (For y fixed, take a sequence (x_n) , such that $x_n y - f(x_n) \nearrow \sup\{xy - f(x) : x \in \mathbb{R}\}$, and use superlinearity to show that (x_n) is bounded, hence, subconverges to some x attaining the sup.)

21. ★ If $f : \mathbb{R} \rightarrow \mathbb{R}$ is superlinear and continuous, then, g is also continuous. (Modify the logic of the previous solution.)

22. Let $f(x) = 1 + [x] - x$, $x \in \mathbb{R}$, where $[x]$ denotes the greatest integer $\leq x$ (Figure 2.6). Compute

$$\lim_{n \nearrow \infty} \left(\lim_{m \nearrow \infty} [f(n!x)]^m \right)$$

for $x \in \mathbb{Q}$ and for $x \notin \mathbb{Q}$.

23. Let $f(x) = 1/x$, $0 < x < 1$. Compute $\mu_c(\delta)$ explicitly for $0 < c < 1$ and $\delta > 0$. Show that $\mu(\delta) = \infty$ for all $\delta > 0$. Conclude that f is not uniformly continuous on $(0, 1)$. (There are two cases, $c \leq \delta$ and $c > \delta$.)

24. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous, and suppose that $f(\infty)$ and $f(-\infty)$ exist and are finite. Show that f is uniformly continuous on \mathbb{R} .
25. ★ Use $\sqrt{2}^{\sqrt{2}}$ to show that there are irrationals a, b , such that a^b is rational.

3

Differentiation

§3.1 Derivatives

Let f be defined on (a, b) , and choose $c \in (a, b)$. We say that f is *differentiable at c* if

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

exists as a real, i.e., exists and is not $\pm\infty$. If it exists, we denote this limit $f'(c)$ or $\frac{df}{dx}(c)$, and we say that $f'(c)$ is the *derivative* of f at c . If f is differentiable at c for all $a < c < b$, we say that f is *differentiable* on (a, b) or, if it is clear from the context, *differentiable*. In this case, the derivative $f' : (a, b) \rightarrow \mathbb{R}$ is a function defined on all of (a, b) .

For example, the function $f(x) = mx + b$ is differentiable on \mathbb{R} with derivative $f'(c) = m$ for all c since

$$\lim_{x \rightarrow c} \frac{(mx + b) - (mc + b)}{x - c} = \lim_{x \rightarrow c} m = m.$$

Since its graph is a line, the derivative of $f(x) = mx + b$ (at any real) is the slope of its graph. In particular, the derivative of a constant function $f(x) = b$ for all x is zero.

If $f(x) = x^2$, then, f is differentiable with derivative

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} \frac{(x - c)(x + c)}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

If f is differentiable at c , then,

$$\begin{aligned} \lim_{x \rightarrow c} f(x) &= \lim_{x \rightarrow c} \left[\left(\frac{f(x) - f(c)}{x - c} \right) (x - c) + f(c) \right] \\ &= \lim_{x \rightarrow c} \left(\frac{f(x) - f(c)}{x - c} \right) \lim_{x \rightarrow c} (x - c) + f(c) \\ &= f'(c) \cdot 0 + f(c) = f(c). \end{aligned}$$

So, f is continuous at c . Hence, *a differentiable function is continuous.*

However, $f(x) = |x|$ is continuous at 0 but not differentiable there since

$$\lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = 1,$$

whereas

$$\lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0} = -1.$$

However,

$$(|x|)' = \frac{x}{|x|}, \quad x \neq 0,$$

since $|x| = x$, hence, $(|x|)' = 1$ on $(0, \infty)$, and $|x| = -x$, hence, $(|x|)' = -1$ on $(-\infty, 0)$.

Derivatives are computed using their arithmetic properties.

Theorem. If f and g are differentiable on (a, b) , and k is real, so are $f + g$, kf , fg , and, for $a < x < b$,

$$(f + g)'(x) = f'(x) + g'(x),$$

$$(kf)'(x) = kf'(x), \quad a < x < b.$$

$$(fg)'(x) = f'(x)g(x) + f(x)g'(x),$$

Moreover, if g is nonzero on (a, b) , then, f/g is differentiable and

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}, \quad a < x < b.$$

The first two identities together are called the *linearity* of the derivative, the third is the *product rule*, whereas the last is the *quotient rule*. To derive these rules, let $a < c < b$. For sums,

$$\begin{aligned} (f + g)'(c) &= \lim_{x \rightarrow c} \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c} \\ &= \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} + \lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c} \\ &= f'(c) + g'(c). \end{aligned}$$

For scalar multiplication,

$$\begin{aligned} (kf)'(c) &= \lim_{x \rightarrow c} \frac{kf(x) - kf(c)}{x - c} \\ &= k \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = kf'(c). \end{aligned}$$

For products,

$$(fg)'(c) = \lim_{x \rightarrow c} \frac{f(x)g(x) - f(c)g(c)}{x - c}$$

$$\begin{aligned}
&= \lim_{x \rightarrow c} \frac{f(x)g(x) - f(c)g(x) + f(c)g(x) - f(c)g(c)}{x - c} \\
&= \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \lim_{x \rightarrow c} g(x) + f(c) \lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c} \\
&= f'(c)g(c) + f(c)g'(c).
\end{aligned}$$

For quotients,

$$\begin{aligned}
\left(\frac{f}{g}\right)'(c) &= \lim_{x \rightarrow c} \frac{f(x)/g(x) - f(c)/g(c)}{x - c} \\
&= \lim_{x \rightarrow c} \frac{f(x)g(c) - f(c)g(x)}{(x - c)g(x)g(c)} \\
&= \lim_{x \rightarrow c} \frac{\frac{f(x) - f(c)}{x - c}g(c) - f(c)\frac{g(x) - g(c)}{x - c}}{g(x)g(c)} \\
&= \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}. \quad \square
\end{aligned}$$

Above, we saw that the derivative of $f(x) = x$ is $f'(x) = 1$. By induction, we show that the derivative of the monomial $f(x) = x^n$ is nx^{n-1} . Since this is true for $n = 1$, assume it is true for $n \geq 1$. Then, by the product rule if $f(x) = x^{n+1}$,

$$f'(x) = (x^{n+1})' = (x^n x)' = (x^n)'x + x^n(x)' = nx^{n-1}x + x^n(1) = (n+1)x^n.$$

This establishes that $(x^n)' = nx^{n-1}$ for all $n \geq 1$. Since polynomials are linear combinations of monomials, they are differentiable everywhere. For example,

$$(x^3 + 5x + 1)' = (x^3)' + (5x)' + (1)' = 3x^2 + 5.$$

Moreover,

$$(1.1) \quad (x^n)' = nx^{n-1}, \quad n \in \mathbb{Z}, x \neq 0.$$

This is clear for $n = 0$ whereas, for $n \geq 1$, using the quotient rule, we find that

$$\begin{aligned}
(x^{-n})' &= \left(\frac{1}{x^n}\right)' = \frac{(1)'x^n - 1(x^n)'}{(x^n)^2} \\
&= \frac{0 \cdot x^n - nx^{n-1}}{x^{2n}} = -\frac{n}{x^{n+1}} = -nx^{-n-1}.
\end{aligned}$$

This establishes (1.1). Another consequence of the quotient rule is that a rational function is differentiable wherever it is defined. For example,

$$\left(\frac{x^2 - 1}{x^2 + 1}\right)' = \frac{(2x)(x^2 + 1) - (x^2 - 1)(2x)}{(x^2 + 1)^2} = \frac{4x}{(x^2 + 1)^2}.$$

We say that a function g is *tangent to f at c* if the difference $f(x) - g(x)$ vanishes faster than first order in $x - c$, i.e., if

$$\lim_{x \rightarrow c} \frac{f(x) - g(x)}{x - c} = 0.$$

Suppose that $g(x) = mx + b$ is tangent to f at c . Since the graph of g is a line, it is reasonable to call it the *line tangent to f at $(c, f(c))$* or, more simply, the *tangent line at c* . Note two lines are tangent to each other iff they coincide. Thus, a function f can have, at most, one tangent line at a given real c .




FIGURE 3.1. The derivative is the slope of the tangent line.

If f is differentiable at c , then, $g(x) = f'(c)(x - c) + f(c)$ is tangent to f at c , since

$$\begin{aligned}\lim_{x \rightarrow c} \frac{f(x) - g(x)}{x - c} &= \lim_{x \rightarrow c} \frac{f(x) - f(c) - f'(c)(x - c)}{x - c} \\ &= \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} - f'(c) = 0.\end{aligned}$$

Hence, the derivative $f'(c)$ of f at c is the slope of the tangent line at c (Figure 3.1).

If f is differentiable at c , there is a positive k and some interval $(c - d, c + d)$ about c on which

$$(1.2) \quad |f(x) - f(c)| \leq k|x - c|, \quad c - d < x < c + d.$$

Indeed, if this were not so, for each $n \geq 1$, we would find a real $x_n \in (c - 1/n, c + 1/n)$ contradicting this claim, i.e., satisfying

$$\left| \frac{f(x_n) - f(c)}{x_n - c} \right| > n.$$

But, then, $x_n \rightarrow c$, and, hence, this inequality would contradict differentiability at c .

The following describes the behavior of derivatives under composition.

Theorem (Chain Rule). *Let f, g be differentiable on $(a, b), (c, d)$, respectively. If $f((a, b)) \subset (c, d)$, then, $g \circ f$ is differentiable on (a, b) with*

$$(g \circ f)'(x) = g'(f(x))f'(x), \quad a < x < b.$$

To see this, let $a < c < b$, and assume, first, $f'(c) \neq 0$. Then, $x_n \rightarrow c$ and $x_n \neq c$ for all $n \geq 1$ implies $f(x_n) \rightarrow f(c)$ and $(f(x_n) - f(c))/(x_n - c) \rightarrow f'(c) \neq 0$. Hence, there is an $N \geq 1$, such that $f(x_n) - f(c) \neq 0$ for $n \geq N$. Thus,

$$\lim_{n \rightarrow \infty} \frac{g(f(x_n)) - g(f(c))}{x_n - c} = \lim_{n \rightarrow \infty} \frac{g(f(x_n)) - g(f(c))}{f(x_n) - f(c)} \cdot \frac{f(x_n) - f(c)}{x_n - c}$$

$$\begin{aligned}
&= \lim_{n \nearrow \infty} \frac{g(f(x_n)) - g(f(c))}{f(x_n) - f(c)} \lim_{n \nearrow \infty} \frac{f(x_n) - f(c)}{x_n - c} \\
&= g'(f(c)) f'(c).
\end{aligned}$$

Since $x_n \rightarrow c$ and $x_n \neq c$ for all $n \geq 1$, by definition of $\lim_{x \rightarrow c}$ (§2.2),

$$(g \circ f)'(c) = \lim_{x \rightarrow c} \frac{g(f(x)) - g(f(c))}{x - c} = g'(f(c)) f'(c).$$

This establishes the result when $f'(c) \neq 0$. If $f'(c) = 0$, by (1.2) there is a k with

$$|g(y) - g(f(c))| \leq k|y - f(c)|$$

for y near $f(c)$. Since $x \rightarrow c$ implies $f(x) \rightarrow f(c)$, in this case, we obtain

$$\begin{aligned}
|(g \circ f)'(c)| &= \lim_{x \rightarrow c} \left| \frac{g(f(x)) - g(f(c))}{x - c} \right| \\
&\leq \lim_{x \rightarrow c} \frac{k|f(x) - f(c)|}{|x - c|} \\
&= k|f'(c)| = 0.
\end{aligned}$$

Hence, $(g \circ f)'(c) = 0 = g'(f(c)) f'(c)$. \square

For example,

$$\left(\left(1 - \frac{x}{n} \right)^n \right)' = n \left(1 - \frac{x}{n} \right)^{n-1} \cdot \left(-\frac{1}{n} \right) = - \left(1 - \frac{x}{n} \right)^{n-1}$$

follows by choosing $g(x) = x^n$ and $f(x) = 1 - x/n$, $0 < x < n$.

If we set $u = f(x)$ and $y = g(u) = g(f(x))$, then, the chain rule takes the easily remembered form

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}.$$

We say that $f : (a, b) \rightarrow \mathbf{R}$ has a *local* maximum at $c \in (a, b)$ if, for some $\delta > 0$, $f(x) \leq f(c)$ on $(c - \delta, c + \delta)$. Similarly, we say that f has a local minimum at $c \in (a, b)$ if, for some $\delta > 0$, $f(x) \geq f(c)$ on $(c - \delta, c + \delta)$. Alternatively, we say that c is a local max or a local min of f . If, instead, these inequalities hold for all x in (a, b) , then, we say that c is a (*global*) maximum or a (*global*) minimum of f on (a, b) . It is possible for a function to have a local maximum at every rational (see Exercise 9).

A *critical point* of a differentiable f is a real c with $f'(c) = 0$. A *critical value* of f is a real d , such that $d = f(c)$ for some critical point c .

Let f defined on (a, b) . Suppose that f has a local minimum at c , and is differentiable there. Then, for $x > c$ near c , $f(x) \geq f(c)$, so,

$$f'(c) = \lim_{x \rightarrow c+} \frac{f(x) - f(c)}{x - c} \geq 0.$$

For $x < c$ near c , $f(x) \geq f(c)$, so,

$$f'(c) = \lim_{x \rightarrow c-} \frac{f(x) - f(c)}{x - c} \leq 0.$$

Hence, $f'(c) = 0$. Applying this result to $g = -f$, we see that if f has a local maximum at c , then, $f'(c) = 0$. We conclude that *a local maximum or a local minimum is a critical point*. The converse is not, generally, true since $c = 0$ is a critical point of $f(x) = x^3$ but is neither a local maximum nor a local minimum.

Using critical points, one can maximize and minimize functions over their domains. For example, to compute

$$\min_{a < x < b} f(x)$$

when f is differentiable, it is enough to compute the critical values of f and compare them with $f(a+)$ and $f(b-)$, assuming these limits exist. If the least of these values is $f(c)$ for some critical point $c \in (a, b)$, then, f is minimized at c . If the least of these values is $f(b-)$ or $f(a+)$, then, f has an inf but no minimum over (a, b) . Similarly, for computing max. For example,

$$\max_{-\infty < x < \infty} (6x - x^2) = 9$$

since the only critical point of $f(x) = 6x - x^2$ is at $x = 3$ and $f(\infty) = f(-\infty) = -\infty$.

Mean Value Theorem. *If f is continuous on $[a, b]$ and differentiable on (a, b) , then, there is a c in (a, b) with*

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

To see this (Figure 3.2), first we subtract a line from f by setting

$$g(x) = f(x) - \left\{ \left[\frac{f(b) - f(a)}{b - a} \right] (x - a) + f(a) \right\}, \quad a < x < b.$$

Then, g is continuous on $[a, b]$, differentiable on (a, b) , and $g(a) = g(b) = 0$. If $g(x) = 0$ everywhere on $[a, b]$, let $a < c < b$ be any real. If $g(x) > 0$ somewhere in (a, b) , let c be a real at which g is maximized. If $g(x) < 0$ somewhere in (a, b) , let c be a real at which g is minimized. In all three cases, we obtain $g'(c) = 0$. Since

$$g'(c) = f'(c) - \frac{f(b) - f(a)}{b - a},$$

we are done. \square




FIGURE 3.2. The mean value theorem.

For example, choose $f(x) = (1 - x/n)^n$, $a = 0$, $b > 0$. Then, $f'(x) = -(1 - x/n)^{n-1}$ is between -1 and 0 when $0 < x < n$. By the mean value theorem, we conclude that

$$0 \leq \frac{1 - (1 - b/n)^n}{b} \leq 1, \quad 0 < b < n, n \geq 1,$$

since the ratio equals the negative of $(f(b) - f(0))/(b - 0)$. The point of this inequality is that, when $b > 0$ is small, the numerator is small enough to compensate for the smallness of the denominator, yielding a quotient bounded between 0 and 1 .

As a consequence of the mean value theorem, if f and g are differentiable on (a, b) and $f'(x) = g'(x)$ for all x , then, f and g differ by a constant, $f(x) = g(x) + C$. To see this, note that $h(x) = f(x) - g(x)$ satisfies $h'(x) = 0$, so, by the mean value theorem $(h(c) - h(d))/(c - d)$ equals h' at some intermediate real. Hence, $h(c) = h(d)$, hence, h is a constant function.

Let $(-b, b)$ be an interval symmetric about 0 . Given a function $f : (-b, b) \rightarrow \mathbb{R}$, its *even part* f^e is the function

$$f^e(x) = \frac{f(x) + f(-x)}{2},$$

and its *odd part* f^o is

$$f^o(x) = \frac{f(x) - f(-x)}{2}.$$

Clearly, $f = f^e + f^o$.

A function f is *even* over $(-b, b)$ if $f = f^e$ on $(-b, b)$ and *odd* over $(-b, b)$ if $f = f^o$ on $(-b, b)$. Thus, an even function satisfies $f(-x) = f(x)$ on $(-b, b)$, whereas an odd function satisfies $f(-x) = -f(x)$ on $(-b, b)$. For example, x^n is even or odd on \mathbb{R} according to whether n is even or odd.

Exercises 3.1

1. Let $a > 0$ and define $f(x) = |x|^a$. Show that f is differentiable at 0 iff $a > 1$.
2. Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by setting $f(x) = 0$, when x is irrational, and setting $f(m/n) = 1/n^3$ when $n > 0$ and m have no common factor. Use Exercise 10 of §1.4 to show that f is differentiable at $\sqrt{2}$. What is $f'(\sqrt{2})$?
3. Let $f(x) = ax^2/2$ with $a > 0$, and set

$$(1.3) \quad g(y) = \max_{-\infty < x < \infty} (xy - f(x)), \quad y \in \mathbb{R}.$$

By direct computation, show that $g(y) = y^2/2a$ and f' and g' are inverses.

4. ★ If $g : \mathbb{R} \rightarrow \mathbb{R}$ is superlinear (Exercise 20 of §2.3) and differentiable, then, $g'(\mathbb{R})$ is unbounded above and below, $\sup g'(\mathbb{R}) = \infty$ and $\inf g'(\mathbb{R}) = -\infty$. (Argue by contradiction, and use the mean value theorem.)

5. Suppose that f is continuous on (a, b) , differentiable on (a, b) punctured at c , $a < c < b$, and $\lim_{x \rightarrow c} f'(x) = L$ exists. Show that $f'(c)$ exists and equals L .
6. Suppose that $f : (a, b) \rightarrow \mathbb{R}$ is differentiable, $a < c < b$, and $f'(c+)$ and $f'(c-)$ exist. Show that $f'(c+) = f'(c) = f'(c-)$. (As opposed to the previous exercise, here, we assume that $f'(c)$ exists.)
7. Suppose that f is differentiable on a bounded interval (a, b) with $|f'| \leq I$. Show that f is of bounded variation (Exercise 4 of §2.2) over (a, b) with total variation $\leq I(b - a)$.
8. Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ in Exercise 1 of §2.2 has a local maximum at every $c \in \mathbb{Q}$.
9. Suppose that $f : (-b, b) \rightarrow \mathbb{R}$ is differentiable. Then, f' is even or odd if f is odd or even, respectively. ★ Moreover, if $f : (-\infty, \infty) \rightarrow \mathbb{R}$ is even and superlinear (Exercise 20 of §2.3), then, the function g , given by (1.3) above, is even.

§3.2 Mapping Properties

To differentiate roots, we need to know how derivatives of inverses behave. But continuous functions are invertible iff they are strictly monotone (§2.3), so, we begin by using the derivative to identify monotonicity.

Theorem. *Let $f : (a, b) \rightarrow \mathbb{R}$ be differentiable. If $f'(x) \neq 0$ for $a < x < b$, then, f is strictly monotone on (a, b) and $f'(x) > 0$ on (a, b) or $f'(x) < 0$ on (a, b) . Moreover, $f'(x) \geq 0$ on (a, b) iff f is increasing, and $f'(x) \leq 0$ on (a, b) iff f is decreasing.*

By the mean value theorem, given $a < x < y < b$, there is a c in (x, y) satisfying

$$f(y) - f(x) = f'(c)(y - x).$$

If f' is never zero, this shows that f is injective, hence, strictly monotone by the inverse function theorem (§2.3). This also shows that $f' \geq 0$ on (a, b) implies f is increasing and $f' \leq 0$ on (a, b) implies f is decreasing. Conversely, f increasing implies $f(x) \geq f(c)$ for $x > c$, so,

$$f'(c) = \lim_{x \rightarrow c+} \frac{f(x) - f(c)}{x - c} \geq 0,$$

for all $a < c < b$. Similarly, if f is decreasing. In particular, we conclude that if f' is never zero and f is monotone, we must have $f' > 0$ on (a, b) or $f' < 0$ on (a, b) . □

It is not, generally, true that strict monotonicity implies the nonvanishing of f' . For example, $f(x) = x^3$ is strictly increasing on \mathbb{R} but $f'(0) = 0$.

Since its derivative was computed in the previous section, the function

$$f(x) = \frac{x^2 - 1}{x^2 + 1}$$

is strictly increasing on $(0, \infty)$ and strictly decreasing on $(-\infty, 0)$. Thus, the critical point $x = 0$ is a minimum of f on \mathbb{R} .

A useful consequence of this theorem is the following: *If f and g are differentiable on (a, b) , continuous on $[a, b]$, $f(a) = g(a)$, and $f'(x) \geq g'(x)$ on (a, b) , then, $f(x) \geq g(x)$ on $[a, b]$.* This follows by applying the theorem to $h = f - g$.

Another consequence is that derivatives, although not necessarily continuous, satisfy the intermediate value property (Exercise 8).

Now we can state the inverse function theorem for differentiable functions.

Inverse Function Theorem. *Let f be continuous on $[a, b]$, differentiable on (a, b) , and suppose that $f'(x) \neq 0$ on (a, b) . Let $[m, M] = f([a, b])$. Then, $f : [a, b] \rightarrow [m, M]$ is invertible and its inverse g is continuous on $[m, M]$, differentiable on (m, M) , and $g'(y) \neq 0$ on (m, M) . Moreover,*

$$g'(y) = \frac{1}{f'(g(y))}, \quad m < y < M.$$

Note, first, that $f' > 0$ on (a, b) or $f' < 0$ on (a, b) by the previous theorem. Suppose that $f' > 0$ on (a, b) , the case $f' < 0$ being entirely similar. Then, f is strictly increasing, hence, the range $[m, M]$ must equal $[f(a), f(b)]$, f is invertible, and its inverse g is strictly increasing and continuous. If $a < c < b$ and $y_n \rightarrow f(c)$, $y_n \neq f(c)$ for all $n \geq 1$, then, $x_n = g(y_n) \rightarrow g(f(c)) = c$ and $x_n \neq c$ for all $n \geq 1$, so, $y_n = f(x_n)$, $n \geq 1$, and

$$\lim_{n \rightarrow \infty} \frac{g(y_n) - g(f(c))}{y_n - f(c)} = \lim_{n \rightarrow \infty} \frac{x_n - c}{f(x_n) - f(c)} = \frac{1}{f'(c)}.$$

Since (y_n) is any sequence converging to $f(c)$, this implies

$$g'(f(c)) = \lim_{y \rightarrow f(c)} \frac{g(y) - g(f(c))}{y - f(c)} = \frac{1}{f'(c)}.$$

Since $y = f(c)$ iff $c = g(y)$, the result follows. \square

As an application, let $b > 0$. Since for $n > 0$, the function $f(x) = x^n$ is continuous on $[0, b]$ and $f'(x) = nx^{n-1} \neq 0$ on $(0, b)$, its inverse $g(y) = y^{1/n}$ is continuous on $[0, b^n]$ and differentiable on $(0, b^n)$ with

$$g'(y) = \frac{1}{f'(g(y))} = \frac{1}{n(g(y))^{n-1}} = \frac{1}{ny^{(n-1)/n}} = \frac{1}{n}y^{(1/n)-1}.$$

Since $b > 0$ is arbitrary, this is valid on $(0, \infty)$. Similarly, this holds on $(0, \infty)$ for $n < 0$.

By applying the chain rule, for all rationals $r = m/n$, the power functions $f(x) = x^r = x^{m/n} = (x^m)^{1/n}$ are differentiable on $(0, \infty)$ with derivative $f'(x) =$

rx^{r-1} , since

$$\begin{aligned} f'(x) &= \left((x^m)^{1/n} \right)' = \frac{1}{n} (x^m)^{1/n-1} (x^m)' \\ &= \frac{1}{n} x^{(m/n)-m} m x^{m-1} = \frac{m}{n} x^{(m/n)-1} = rx^{r-1}. \end{aligned}$$

Thus, the derivative of $f(x) = x^r$ is $f'(x) = rx^{r-1}$ for $x > 0$, for all $r \in \mathbb{Q}$.

Using the chain rule, we now know how to differentiate any algebraic function. For example, the derivative of

$$f(x) = \sqrt{\frac{1-x^2}{1+x^2}}, \quad 0 < x < 1,$$

is

$$f'(x) = \frac{1}{2} \left(\frac{1-x^2}{1+x^2} \right)^{-1/2} \cdot \left(\frac{-4x}{(1+x^2)^2} \right) = \frac{-2x}{(1+x^2)\sqrt{1-x^4}}, \quad 0 < x < 1.$$

Now let $a > 0$ and let $r < s$ be rationals with $r < a < s$. We wish to compute

$$(2.1) \quad \lim_{x \rightarrow 1+} \frac{x^a - 1}{x - 1}$$

for any $a \in \mathbb{R}$. Recall that, for $a \in \mathbb{Q}$, this limit is $a1^{a-1} = a$. Since for any $x_n \rightarrow 1+$, the sequence $B_n = (x_n^a - 1)/(x_n - 1)$ lies between the sequences $A_n = (x_n^r - 1)/(x_n - 1)$ and $C_n = (x_n^s - 1)/(x_n - 1)$, the upper and lower limits of (B_n) lie between $\lim_{n \nearrow \infty} A_n = r$ and $\lim_{n \nearrow \infty} C_n = s$. Since $r < a < s$ are arbitrary, the upper and lower limits both equal a , hence, $B_n = (x_n^a - 1)/(x_n - 1) \rightarrow a$, hence, the limit (2.1) equals a . Since $f(x) = 1/x$ is continuous at $x = 1$, $x_n \rightarrow 1-$ implies $y_n = 1/x_n \rightarrow 1+$, so,

$$\lim_{n \nearrow \infty} \frac{x_n^a - 1}{x_n - 1} = \lim_{n \nearrow \infty} \frac{1/y_n^a - 1}{1/y_n - 1} = \lim_{n \nearrow \infty} y_n^{1-a} \cdot \frac{y_n^a - 1}{y_n - 1} = 1 \cdot a = a.$$

Thus,

$$\lim_{x \rightarrow 1-} \frac{x^a - 1}{x - 1} = a.$$

Hence, $f(x) = x^a$ is differentiable at $x = 1$ with $f'(1) = a$. Since

$$\lim_{x \rightarrow c} \frac{x^a - c^a}{x - c} = c^{a-1} \lim_{x/c \rightarrow 1} \frac{(x/c)^a - 1}{(x/c) - 1} = ac^{a-1},$$

f is differentiable on $(0, \infty)$ with $f'(c) = ac^{a-1}$. Thus, for all real $a > 0$, the derivative of $f(x) = x^a$ at $x > 0$ is $f'(x) = ax^{a-1}$. Using the quotient rule, the same result holds for real $a \leq 0$.

As an application, let v be any real greater than 1. Then, by the chain rule, the derivative of $f(x) = (1+x)^v - 1 - vx$ is $f'(x) = v(1+x)^{v-1} - v$, hence, the only critical point is $x = 0$. Since $f(-1) = -1 + v > 0 = f(0)$ and $f(\infty) = \infty$,

the minimum of f over $(-1, \infty)$ is $f(0) = 0$. Hence,

$$(2.2) \quad (1 + b)^v \geq 1 + vb, \quad b \geq -1.$$

We already knew this for v a natural (Exercise 6 of §1.4), but now we know this for any real $v \geq 1$.

Now, we compute the derivative of the exponential function $f(x) = a^x$ with base $a > 1$. We begin with finding $f'(0)$.

If $0 < x \leq y$ and $a > 1$, then, insert $v = y/x \geq 1$ and $b = a^x - 1 > 0$ in (2.2), and rearrange to get

$$\frac{a^x - 1}{x} \leq \frac{a^y - 1}{y}, \quad 0 < x \leq y.$$

Thus,

$$m_+ = \lim_{x \rightarrow 0+} \frac{a^x - 1}{x}$$

exists since it equals $\inf\{(a^x - 1)/x : x > 0\}$ (Exercise 2 of §2.2). Moreover, $m_+ \geq 0$ since $a^x > 1$ for $x > 0$. Also,

$$m_- = \lim_{x \rightarrow 0-} \frac{a^x - 1}{x} = \lim_{x \rightarrow 0+} \frac{a^{-x} - 1}{-x} = \lim_{x \rightarrow 0+} a^{-x} \cdot \frac{a^x - 1}{x} = 1 \cdot m_+ = m_+.$$

Hence, the exponential with base $a > 1$ is differentiable at $x = 0$, and we denote its derivative there by $m(a)$. Since $a^x = a^c a^{x-c}$,

$$\lim_{x \rightarrow c} \frac{a^x - a^c}{x - c} = a^c \lim_{x \rightarrow c} \frac{a^{x-c} - 1}{x - c} = a^c m(a).$$

Hence, f is differentiable on \mathbb{R} , and $f'(x) = a^x m(a)$ with $m(a) \geq 0$. If $m(a) = 0$, then, $f'(x) = 0$ for all x , hence, f is constant, a contradiction. Hence, $m(a) > 0$. Also, for $b > 1$ and $a > 1$,

$$\begin{aligned} m(b) &= \lim_{x \rightarrow 0} \frac{b^x - 1}{x} = \lim_{x \rightarrow 0} \frac{(a^{\log_a b})^x - 1}{x} \\ &= \lim_{x \rightarrow 0} \frac{a^{x \log_a b} - 1}{x} = m(a) \log_a b, \end{aligned}$$

by the chain rule. By fixing a and varying b , we see that m is a continuous, strictly increasing function with $m(\infty) = \infty$ and $m(1+) = 0$. By the intermediate value property §2.2, we conclude that $m((1, \infty)) = (0, \infty)$.

Thus, and this is very important, there is a unique real $e > 1$ with $m(e) = 1$. The exponential and logarithm functions with base e are called *natural*. Throughout the book, e denotes this particular number. The decimal expansion of e is computed in §3.4. We summarize the results.

Theorem. For all $a > 0$, the exponential $f(x) = a^x$ is differentiable on \mathbb{R} . There is a unique real $e > 1$, such that $f(x) = e^x$ implies $f'(x) = e^x$. Moreover, $f(x) = a^x$ implies $f'(x) = a^x \log_e a$.

For $a > 1$, this was derived above. To derive the theorem for $0 < a < 1$, use $a^x = (1/a)^{-x}$ and the chain rule. \square

In the sequel, $\log x$ will denote $\log_e x$, i.e., we drop the e when writing the natural logarithm. Then,

$$e^{\log x} = x, \quad \log e^x = x.$$

We end with the derivative of $f(x) = \log_a x$. Since this is the inverse of the exponential,

$$f'(x) = \frac{1}{a^{f(x)} \log a} = \frac{1}{x \log a}.$$

Thus, $f(x) = \log_a x$ implies $f'(x) = 1/x \log a$, $x > 0$. In particular $\log e = 1$, so, $f(x) = \log x$ implies $f'(x) = 1/x$, $x > 0$.

For example, combining the above with the chain rule,

$$(\log |x|)' = \frac{1}{x}, \quad x \neq 0.$$

Another example is ($x \neq \pm 1$)

$$\left[\log \left(\frac{x-1}{x+1} \right) \right]' = \frac{1}{\left(\frac{x-1}{x+1} \right)} \cdot \left(\frac{x-1}{x+1} \right)' = \left(\frac{x+1}{x-1} \right) \cdot \frac{2}{(x+1)^2} = \frac{2}{x^2-1}.$$

We will need the following in §3.4.

Generalized Mean Value Theorem. *If f and g are continuous on $[a, b]$, differentiable on (a, b) , and $g'(x) \neq 0$ on (a, b) , there exists a c in (a, b) , such that¹²*

$$\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}.$$

Either $g' > 0$ on (a, b) or $g' < 0$ on (a, b) . Assume $g' > 0$ on (a, b) . To see the theorem, let h denote the inverse function of g , so, $h(g(x)) = x$, and set $F(x) = f(h(x))$. Then $F(g(x)) = f(x)$, F is continuous on $[g(a), g(b)]$ and differentiable on $(g(a), g(b))$. So, applying the mean value theorem, the chain rule, and the inverse function theorem, there is a d in $(g(a), g(b))$, such that

$$\begin{aligned} \frac{f(b) - f(a)}{g(b) - g(a)} &= \frac{F(g(b)) - F(g(a))}{g(b) - g(a)} \\ &= F'(d) = f'(h(d))h'(d) = \frac{f'(h(d))}{g'(h(d))}. \end{aligned}$$

Now, let $c = h(d)$. Then, c is in (a, b) . The case $g' < 0$ on (a, b) is similar. \square

We end the section with l'Hopital's rule.

¹² $g(b) - g(a)$ is not zero because it equals $g'(d)(b - a)$ for some $a < d < b$.

L'Hopital's Rule. Let f and g be differentiable on an open interval (a, b) punctured at c , $a < c < b$. Suppose that $\lim_{x \rightarrow c} f(x) = 0$ and $\lim_{x \rightarrow c} g(x) = 0$. Then, $g'(x) \neq 0$ for $x \neq c$ and

$$\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = L$$

imply¹³

$$(2.3) \quad \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L.$$

To obtain this, define f and g at c by setting $f(c) = g(c) = 0$. Then, f and g are continuous on (a, b) . Now, let $x_n \rightarrow c+$. Apply the generalized mean value theorem on (c, x_n) for each $n \geq 1$. Then,

$$\frac{f(x_n)}{g(x_n)} = \frac{f(x_n) - f(c)}{g(x_n) - g(c)} = \frac{f'(d_n)}{g'(d_n)} \rightarrow L,$$

since $c < d_n < x_n$. Similarly, this also holds when $x_n \rightarrow c-$, and, thus, this holds for $x_n \rightarrow c$, which establishes (2.3). \square

The above deals with the “indeterminate form” $f(x)/g(x) \rightarrow 0/0$. The case $f(x)/g(x) \rightarrow \infty/\infty$ can be handled by turning the fraction $f(x)/g(x)$ upside down and applying the above. We do not state this case as we do not use it.

Exercises 3.2

1. Show that $1 + x \leq e^x$ for $x \geq 0$.
2. Let $f(x) = e^{|x|} - 1$. With $g(y)$ as in Exercise 3 of §3.1, by direct computation, show that $g(y) = |y| \log |y| - |y| + 1$ for $|y| \geq 1$ and $g(y) = 0$ for $|y| \leq 1$ (Exercise 9 of §3.1).
3. Show that $\lim_{x \rightarrow 0} \log(1 + x)/x = 1$ and $\lim_{n \nearrow \infty} (1 + a/n)^n = e^a$ for $a \in \mathbf{R}$ (take the log of both sides). If $a_n \rightarrow a$, show also that $\lim_{n \nearrow \infty} (1 + a_n/n)^n = e^a$.
4. Let $x \in \mathbf{R}$. Show that the sequence $(1 + x/n)^n$, $n \geq |x|$, increases to e^x as $n \nearrow \infty$ (use (2.2)).
5. Let $f(x) = |x|^p/p$ with $p > 1$. With $g(y)$ as in Exercise 3 of §3.1, by direct computation, show that $g(y) = |y|^q/q$ where $(1/p) + (1/q) = 1$ and f' and g' are inverses.
6. Use the mean value theorem to show that

$$1 - \frac{1}{\sqrt{1 + x^2}} \leq x^2, \quad x > 0.$$

¹³ $g'(x) \neq 0$ for $x \neq c$ since $g(x) = g(x) - g(c) = g'(d)(x - c)$.

7. Use the mean value theorem to show that

$$\frac{1}{(2j-1)^x} - \frac{1}{(2j)^x} \leq \frac{x}{(2j-1)^{x+1}}, \quad j \geq 1, x > 0.$$

8. If $f : (a, b) \rightarrow \mathbb{R}$ is differentiable, then, $f' : (a, b) \rightarrow \mathbb{R}$ satisfies the *intermediate value property*: If $a < c < d < b$ and $f'(c) < L < f'(d)$, then, $L = f'(x)$ for some $c < x < d$. (Start with $L = 0$. Here the point is that f' need not be continuous.)

9. ★ If $g : \mathbb{R} \rightarrow \mathbb{R}$ is superlinear (Exercise 4 of §3.1) and differentiable, then, $g'(\mathbb{R}) = \mathbb{R}$, i.e., g' is surjective.

§3.3 Graphing Techniques

Let f be differentiable on (a, b) . If $f' = df/dx$ is differentiable on (a, b) , we denote its derivative by $f'' = (f')'$; f'' is the *second derivative* of f . If f'' is differentiable on (a, b) , $f''' = (f'')'$ is the *third derivative* of f . In general, we let $f^{(n)}$ denote the *n th derivative* or the *derivative of order n* , where, by convention, we take $f^{(0)} = f$. If f has all derivatives, f', f'', f''', \dots , we say f is *smooth* on (a, b) .

An alternate and useful notation for higher derivatives is obtained by thinking of $f' = df/dx$ as the result of applying d/dx to f , i.e., $df/dx = (d/dx)f$. From this point of view, d/dx signifies the operation of differentiation. Thus, applying d/dx twice, we obtain

$$f'' = \left(\frac{d}{dx}\right) \left(\frac{d}{dx}\right) f = \left(\frac{d^2}{dx^2}\right) f = \frac{d^2 f}{dx^2}.$$

Similarly, third derivatives may be denoted

$$f''' = \left(\frac{d}{dx}\right) \left(\frac{d^2 f}{dx^2}\right) = \frac{d^3 f}{dx^3}.$$

For example, $f(x) = x^2$ has $f'(x) = 2x$, $f''(x) = 2$, and $f^{(n)}(x) = 0$ for $n \geq 3$. More generally, by induction, $f(x) = x^n$, $n \geq 0$, has derivatives

$$(3.1) \quad f^{(k)}(x) = \frac{d^k f}{dx^k} = \begin{cases} \frac{n!}{(n-k)!} x^{n-k}, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

so, $f(x) = x^n$ is smooth. By the arithmetic properties of derivatives, it follows that rational functions are smooth wherever they are defined.

Not all functions are smooth. The function $f(x) = |x|$ is not differentiable at zero. Using this, one can show that $f(x) = x^n |x|$ is n times differentiable on \mathbb{R} , but $f^{(n)}$ is not differentiable at zero. More generally, for f, g differentiable, we do not expect $\max(f, g)$ to be differentiable. However, since $f(x) = x^{1/n}$ is smooth on $(0, \infty)$, algebraic functions are smooth on any open interval of definition. Also, the functions x^a , a^x , and $\log_a x$ are smooth on $(0, \infty)$.

We know the sign of f' determines the monotonicity of f , in the sense that $f' \geq 0$ iff f is increasing and $f' \leq 0$ iff f is decreasing. How is the sign of f'' reflected in the graph of f ? Since $f'' = (f')'$, we see that $f'' \geq 0$ iff f' is increasing and $f'' \leq 0$ iff f' is decreasing.

More precisely, we say f is *convex* on (a, b) if, for all $a < x < y < b$,

$$f((1-t)x + ty) \leq (1-t)f(x) + tf(y), \quad 0 \leq t \leq 1.$$

Take any two points on the graph of f and join them by a chord or line segment. Then, f is convex if the chord lies on or above the graph (Figure 3.3). We say f is *concave* on (a, b) if, for all $a < x < y < b$,

$$f((1-t)x + ty) \geq (1-t)f(x) + tf(y), \quad 0 \leq t \leq 1.$$

Take any two points on the graph of f and join them by a chord. Then, f is concave if the chord lies on or below the graph.




FIGURE 3.3. Examples of convex and strictly convex functions.

For example $f(x) = x^2$ is convex and $f(x) = -x^2$ is concave. A function $f : (a, b) \rightarrow \mathbb{R}$ that is both convex and concave is called *affine*. It is easy to see that $f : (a, b) \rightarrow \mathbb{R}$ is affine iff $f' = m$ is constant on (a, b) .

Similarly, we say that f is *strictly convex* on (a, b) if, for all $a < x < y < b$,

$$f((1-t)x + ty) < (1-t)f(x) + tf(y), \quad 0 < t < 1.$$

Take any two points on the graph of f and join them by a chord. Then, f is strictly convex if the chord lies strictly above the graph. Similarly, we define *strictly concave*.

Note that a strictly convex $f : (a, b) \rightarrow \mathbb{R}$ cannot attain its minimum m at more than one point in (a, b) . Indeed, if f had two minima at x and x' and $x'' = (x + x')/2$, then, $f(x'') < [f(x) + f(x')]/2 = (m + m)/2 = m$, contradicting the fact that m is a minimum.

The negative of a (strictly) convex function is (strictly) concave.

Theorem. Suppose that f is differentiable on (a, b) . Then, f is convex iff f' is increasing and f is concave iff f' is decreasing. Moreover, f is strictly convex iff f' is strictly increasing, and f is strictly concave iff f' is strictly decreasing. If f is twice differentiable on (a, b) , then, f is convex iff $f'' \geq 0$ and f is concave iff $f'' \leq 0$. Moreover, f is strictly convex if $f'' > 0$, and f is strictly concave if $f'' < 0$.

Since $-f$ is convex iff f is concave, we derive only the convex part. First, suppose that f' is increasing. If $a < x < y < b$ and $0 < t < 1$, let $z = (1-t)x + ty$. Then,

$$\frac{f(z) - f(x)}{z - x} = f'(c)$$

for some $x < c < z$. Also

$$\frac{f(y) - f(z)}{y - z} = f'(d)$$

for some $z < d < y$. Since $f'(c) \leq f'(d)$,

$$(3.2) \quad \frac{f(z) - f(x)}{z - x} \leq \frac{f(y) - f(z)}{y - z}.$$

Clearing denominators in this last inequality, we obtain convexity. Conversely, suppose that f is convex and let $a < x < z < y < b$. Then, we have (3.2). If $a < x < z < y < w < b$, apply (3.2) to $a < x < z < y < b$, then, apply (3.2) to $a < z < y < w < b$. Combining the resulting inequalities yields

$$\frac{f(z) - f(x)}{z - x} \leq \frac{f(w) - f(y)}{w - y}.$$

Fixing x , y , and w and letting $z \rightarrow x$ yields

$$f'(x) \leq \frac{f(w) - f(y)}{w - y}.$$

Let $w \rightarrow y$ to obtain $f'(x) \leq f'(y)$, hence, f' is increasing. If f' is strictly increasing, then, the inequality (3.2) is strict, hence, f is strictly convex. Conversely, if f' is increasing but $f'(c) = f'(d)$ for some $c < d$, then, f' is constant on $[c, d]$. Hence, f is affine on $[c, d]$ contradicting strict convexity. This shows that f is strictly convex iff f' is strictly increasing.

When f is twice differentiable, $f'' \geq 0$ iff f' is increasing, hence, the third statement. Since $f'' > 0$ implies f' strictly increasing, we also have the fourth statement. \square

A key feature of convexity (Figure 3.4) is that the graph of a convex function lies above any of its tangent lines (Exercise 3).




FIGURE 3.4. A convex function lies above any of its tangents.

A real c is an *inflection point* of f if f is convex on one side of c and concave on the other. For example $c = 0$ is an inflection point of $f(x) = x^3$ since f is convex on $x > 0$ and concave on $x < 0$. From the theorem we see that $f''(c) = 0$ at any inflection point c where f is twice differentiable.

If c is a critical point and $f''(c) > 0$, then, f' is strictly increasing near c , hence, $f'(x) < 0$ for $x < c$ near c and $f'(x) > 0$ for $x > c$ near c . Thus, $f'(c) = 0$ and $f''(c) > 0$ implies c is a local minimum. Similarly, $f'(c) = 0$ and $f''(c) < 0$ implies c is a local maximum. The converses are not, generally, true since $c = 0$ is a minimum of $f(x) = x^4$, but $f'(0) = f''(0) = 0$.

For example, $f(x) = a^x$, $a > 1$, satisfies $f'(x) = a^x \log a$ and $f''(x) = a^x (\log a)^2$. Since $\log a > 0$, a^x is increasing and strictly convex everywhere. Also $f(x) = \log_a x$ has $f'(x) = 1/x \log a$ and $f''(x) = -1/x^2 \log a$, so, $\log_a x$ is increasing and strictly concave everywhere. The graphs are as shown in Figure 3.5.




FIGURE 3.5. The exponential and logarithm functions.

In the following, we sketch the graphs of some twice differentiable functions on an interval (a, b) , using knowledge of the critical points, the inflection points, the signs of f' and f'' , and $f(a+)$, $f(b-)$.


If $f(x) = 1/(x^2 + 1)$, $-\infty < x < \infty$, then, $f'(x) = -2x/(x^2 + 1)^2$. Hence, $f'(x) < 0$ for $x > 0$ and $f'(x) > 0$ for $x < 0$, so, f is increasing for $x < 0$ and decreasing for $x > 0$. Hence, 0 is a global maximum. Moreover,

$$f''(x) = \left(\frac{-2x}{(x^2 + 1)^2} \right)' = \frac{6x^2 - 2}{(x^2 + 1)^3},$$

so, $f''(0) < 0$ which is consistent with 0 being a maximum. Now, $f''(x) < 0$ on $|x| < 1/\sqrt{3}$ and $f''(x) > 0$ on $|x| > 1/\sqrt{3}$. Hence, $x = \pm 1/\sqrt{3}$ are inflection points. Since $f(0) = 1$ and $f(\infty) = f(-\infty) = 0$, we obtain the graph in Figure 3.6.

Let $f(x) = 1/\sqrt{x(1-x)}$, $0 < x < 1$. Then,


$$f'(x) = \frac{2x - 1}{2(x(1-x))^{3/2}}$$

FIGURE 3.6. $f(x) = 1/(1+x^2)$.

and

$$f''(x) = \frac{3 - 8x(1-x)}{4(x(1-x))^{5/2}}.$$

Thus, $x = 1/2$ is a critical point. Since $f''(1/2) > 0$, $x = 1/2$ is a local minimum. In fact, f is decreasing to the left of $1/2$ and increasing to the right of $1/2$, hence, $1/2$ is a global minimum. Since $3 - 8x(1-x) > 0$ on $(0, 1)$, $f''(x) > 0$, hence, f is convex. Since $f(0+) = \infty$ and $f(1-) = \infty$, the graph is as shown in Figure 3.7.

FIGURE 3.7. $f(x) = 1/\sqrt{x(1-x)}$.

Let $f(x) = \frac{3x+1}{x(1-x)}$. This rational function is defined away from $x = 0$ and $x = 1$. Thus, we graph f on the intervals $(-\infty, 0)$, $(0, 1)$, $(1, \infty)$. Computing,

$$f'(x) = \frac{3x^2 + 2x - 1}{x^2(1-x)^2}.$$

Solving $3x^2 + 2x - 1 = 0$, $x = -1, 1/3$ are the critical points. Moreover, $f(\infty) = 0$, $f(1+) = -\infty$, $f(1-) = \infty$, $f(0+) = \infty$, $f(0-) = -\infty$, and $f(-\infty) = 0$. Since there are no critical points in $(1, \infty)$, f is increasing on $(1, \infty)$. Moreover,

$$f''(x) = \frac{6x^3 - 6x(1-x) + 2}{x^3(1-x)^3},$$

so, f is concave on $(1, \infty)$. Moreover, the numerator in $f''(x)$ is $\geq 1/2$ on $(0, 1)$. Hence, $f''(x) > 0$ on $(0, 1)$. Hence, f is convex in $(0, 1)$ and $x = 1/3$ is a minimum of f . Since $f''(-1) = -1 < 0$, $x = -1$ is a maximum. Since




FIGURE 3.8. $f(x) = \frac{3x + 1}{x(1 - x)}$.

$f''(x) \rightarrow 0+$ as $x \rightarrow -\infty$, there is an inflection point in $(-\infty, -1)$. Thus, the graph is as shown in Figure 3.8.

To graph $x^n e^{-x}$, $x \geq 0$, we will need to show that

$$(3.3) \quad \lim_{x \rightarrow \infty} x^n e^{-x} = 0.$$

To establish the limit, we first show that

$$(3.4) \quad e^x \geq 1 + x + \frac{x^2}{2!} + \cdots + \frac{x^n}{n!}, \quad x \geq 0,$$

for all $n \geq 1$. We do this by induction. For $n = 1$, let $f(x) = e^x$, $g(x) = 1 + x$. Then, $f'(x) = e^x \geq 1 = g'(x)$ on $x > 0$ and $f(0) = g(0)$, hence, $f(x) \geq g(x)$ establishing (3.4) for $n = 1$. Now, let $g_n(x)$ denote the right side of (3.4), and suppose that (3.4) is true for some $n \geq 1$. Since $f'(x) = f(x) \geq g_n(x) = g'_{n+1}(x)$ and $f(0) = g_{n+1}(0)$, we conclude that $f(x) \geq g_{n+1}(x)$, establishing (3.4) for $n + 1$. By induction, (3.4) is true for all $n \geq 1$. Now, (3.4) with $n + 1$ replacing n implies $e^x \geq x^{n+1}/(n + 1)!$ which implies

$$x^n e^{-x} \leq \frac{(n + 1)!}{x}, \quad x > 0,$$

which implies (3.3).


Setting $f_n(x) = x^n e^{-x}$, $n \geq 1$, $f_n(0) = 0$ and $f_n(\infty) = 0$. Moreover,

$$f'_n(x) = x^{n-1}(n - x)e^{-x}$$

and

$$f''_n(x) = x^{n-2} [x^2 - 2nx + n(n - 1)] e^{-x}.$$

Thus, the critical point is $x = n$ and f_n is increasing on $(0, n)$ and decreasing on (n, ∞) . Hence, $x = n$ is a max. The reals $x = n \pm \sqrt{n}$ are inflection points. Between them, f_n is concave and elsewhere convex. The graph is as shown in Figure 3.9.

FIGURE 3.9. $f_n(x) = x^n e^{-x}$.

If we let $n \nearrow \infty$ in (3.4), we obtain

$$e^x \geq \sum_{n=0}^{\infty} \frac{x^n}{n!}, \quad x \geq 0.$$

As a consequence (n th term test §1.5),

$$(3.5) \quad \lim_{n \nearrow \infty} \frac{x^n}{n!} = 0$$

for all $x \geq 0$, hence, for all x . Using (3.3), we can derive other limits.

Theorem. For $a > 0$, $b > 0$, and $c > 0$,

- $\lim_{x \rightarrow \infty} x^a e^{-bx} = 0$;
- $\lim_{t \rightarrow 0+} t^b (-\log t)^a = 0$; in particular, $t \log t \rightarrow 0$ as $t \rightarrow 0+$;
- $\lim_{n \nearrow \infty} (\log n)^a / n^b = 0$; in particular, $\log n / n \rightarrow 0$ as $n \nearrow \infty$;
- if $c < 1$, $\lim_{n \nearrow \infty} n^a c^n = 0$. If $c > 1$, $\lim_{n \nearrow \infty} n^{-a} c^n = \infty$;
- $\lim_{n \nearrow \infty} n^{1/n} = 1$.

To obtain the first limit, choose $n > a$, and let $y = bx$. Then, $x \rightarrow \infty$ implies $y \rightarrow \infty$, hence, $x^a e^{-bx} = y^a e^{-y} / b^a < y^n e^{-y} / b^a \rightarrow 0$ by (3.3). Substituting $t = e^{-x}$ in the first yields the second since $e^{-x} \rightarrow 0+$, as $x \rightarrow \infty$. Substituting $t = 1/n$ in the second yields the third. For the fourth, in the first, replace x by n and e^{-b} by c , if $c < 1$. If $c > 1$, $n^{-a} c^n = 1/n^a (1/c)^n \rightarrow \infty$ by what we just derived. For the fifth, take the exponential of both sides of the third with $a = b = 1$. Since e^x is continuous, we obtain the fifth. \square

The moral of the theorem is $\log n \ll n \ll e^n$ as $n \nearrow \infty$, where $A \ll B$ means that any positive power of A is much smaller than B .

Let us use (3.1) to derive the binomial theorem. If $n \geq 1$, then, $(c + x)^n$ is a polynomial of degree n , hence, there are numbers a_0, \dots, a_n with

$$(3.6) \quad f(x) = (c + x)^n = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0.$$

Let us compute the derivatives of f by using either of the expressions in (3.6). The left expression and (3.1) and the chain rule yield $f^{(k)}(0) = n! / (n - k)! c^{n-k}$. The

right expression yields $f^{(k)}(0) = k!a_k$. Hence, $a_k = n!c^{n-k}/(n-k)!k!$. Now, define the *binomial coefficient* (read “ n -choose- k ”)

$$\binom{n}{k} = \frac{n!}{(n-k)!k!}, \quad 0 \leq k \leq n.$$

Then, we obtain the following.

Binomial Theorem. If $n \geq 1$ and $a, b \in \mathbb{R}$, then,

$$\begin{aligned} (a+b)^n &= a^n + \binom{n}{1}a^{n-1}b + \cdots + \binom{n}{n-1}ab^{n-1} + b^n \\ &= \sum_{j=0}^n \binom{n}{j}a^{n-j}b^j. \quad \square \end{aligned}$$

Below is a set of Exercises (7–11) that describe the *conjugate* of a convex function, and derive the result that the conjugate of the conjugate of f is f . Examples of conjugates are given in Exercise 3 of §3.1 and Exercises 2 and 5 of §3.2. The perfect symmetry between f and its conjugate g is exhibited in Exercise 11.

Exercises 3.3

1. Graph $f(x) = (x + 2/x)/2$ for $x > 0$.
2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be convex and, for $b \neq a$, let

$$s[a, b] = \frac{f(b) - f(a)}{b - a}.$$

Show that $a < b < c$ implies $s[a, b] \leq s[a, c] \leq s[b, c]$.

3. Suppose that $f : (a, b) \rightarrow \mathbb{R}$ is convex and differentiable at c with $a < c < b$. Then,

$$\frac{f(x) - f(c)}{x - c} \leq f'(c) \leq \frac{f(y) - f(c)}{y - c}, \quad a < x < c < y < b.$$

Moreover, the graph of f lies above its tangent line at c ,

$$f(x) \geq f'(c)(x - c) + f(c), \quad a < x < b.$$

4. Suppose that $f : (a, b) \rightarrow \mathbb{R}$ is convex. Then, for all $c \in (a, b)$,

$$f'_\pm(c) = \lim_{x \rightarrow c^\pm} \frac{f(x) - f(c)}{x - c}$$

both exist, and $f'_-(c) \leq f'_+(c)$.

5. (Maximum Principle) Suppose that $f : (a, b) \rightarrow \mathbb{R}$ is convex and has a maximum at some c in (a, b) . Then, f is constant.

6. Suppose that $f : (a, b) \rightarrow \mathbf{R}$ is convex, $g : (a, b) \rightarrow \mathbf{R}$ is differentiable, and $f - g$ attains a local maximum at some $a < c < b$. Show that $f'(c)$ exists and equals $g'(c)$.

7. ★ If $f : \mathbf{R} \rightarrow \mathbf{R}$ satisfies

$$\lim_{x \rightarrow \pm\infty} \frac{f(x)}{|x|} = +\infty,$$

we say f is *superlinear*. If f is superlinear and continuous, then (Exercises 20 and 21 of §2.3),

$$g(y) = \max_{-\infty < x < \infty} [xy - f(x)], \quad y \in \mathbf{R},$$

is well defined, superlinear, and continuous. Show that g is convex and, for each y , if x attains the max in the definition of $g(y)$, then, $g'_+(y) \geq x \geq g'_-(y)$. (If y is fixed and x is such that $g(y) = xy - f(x)$, then, $g(z) \geq xz - f(x)$ for all z .)

8. ★ If f is superlinear and strictly convex then, for each y , the max in the definition of $g(y)$ is attained at a unique $x = x^\#(y)$, depending on y . Moreover, in addition, if f is continuous, then, g is differentiable at y with $g'(y) = x^\#(y)$.

9. ★ Suppose that $f : \mathbf{R} \rightarrow \mathbf{R}$ is superlinear, continuous, and strictly convex. If

$$g(y) = \max_{-\infty < x < \infty} (xy - f(x)), \quad y \in \mathbf{R},$$

then, g is superlinear, differentiable, and

$$f(x) = \max_{-\infty < y < \infty} (xy - g(y)).$$

The function g is the *conjugate* of f , and f is the conjugate of g . (Use Exercise 9 of §3.2.)

10. ★ If f is superlinear and differentiable, then, for each y , the max in the definition of $g(y)$ in Exercise 7 is attained at a point x satisfying $f'(x) = y$. Conclude that $f' : \mathbf{R} \rightarrow \mathbf{R}$ is surjective.

11. ★ If $f : \mathbf{R} \rightarrow \mathbf{R}$ is superlinear, differentiable, and strictly convex, then, g is superlinear, differentiable, and strictly convex, with g' equal to the inverse of f' . (Remember, when f is differentiable and strictly convex, f' is strictly increasing.)

12. Graph f , g , f' , and g' where f and g are as in Exercise 2 of §3.2.

13. Show that $f(x) = e^x$ is convex on \mathbf{R} . Deduce the inequality $a^t b^{1-t} \leq ta + (1-t)b$ valid for $a > 0$, $b > 0$, and $0 \leq t \leq 1$.

14. ★ Let f and g be as in Exercise 11. Also assume f is smooth. Then, g is smooth iff $f''(x) > 0$ for all $x \in \mathbf{R}$.

§3.4 Power Series

Let $f : (a, b) \rightarrow \mathbf{R}$ be continuous, and fix c in (a, b) . Suppose that f is differentiable at c , and define $h_1(x)$ by

$$h_1(x) = \begin{cases} \frac{f(x) - f(c)}{x - c}, & x \neq c, \\ f'(c), & x = c. \end{cases}$$

Then, h_1 is continuous, when $x \neq c$, and $\lim_{x \rightarrow c} h_1(x) = f'(c) = h_1(c)$, so, h_1 is continuous at c . Thus, differentiability at c implies there is a continuous function $h_1 : (a, b) \rightarrow \mathbf{R}$ satisfying $h_1(c) = f'(c)$ and

$$(4.1) \quad f(x) = f(c) + h_1(x)(x - c), \quad a < x < b.$$

We wish to derive the analog of this result for higher derivatives.

Let f be differentiable on (a, b) with f' continuous on (a, b) , and suppose that f' is differentiable at $c \in (a, b)$. Define $h_2 : (a, b) \rightarrow \mathbf{R}$ by

$$h_2(x) = \begin{cases} \frac{f(x) - f(c) - f'(c)(x - c)}{(x - c)^2/2}, & x \neq c, \\ f''(c), & x = c. \end{cases}$$

Then, h_2 is continuous. To see this, note that h_2 is continuous when $x \neq c$, whereas applying L'Hopital's rule yields

$$\begin{aligned} \lim_{x \rightarrow c} h_2(x) &= \lim_{x \rightarrow c} \frac{f(x) - f(c) - f'(c)(x - c)}{(x - c)^2/2} \\ &= \lim_{x \rightarrow c} \frac{f'(x) - f'(c)}{x - c} = f''(c). \end{aligned}$$

Thus, $h_2 : (a, b) \rightarrow \mathbf{R}$ is a continuous function satisfying $h_2(c) = f''(c)$ and

$$(4.2) \quad f(x) = f(c) + f'(c)(x - c) + \frac{1}{2}h_2(x)(x - c)^2, \quad a < x < b.$$

Now, we carry out this procedure in the general case. Suppose that f is n times differentiable on (a, b) with $f^{(n)}$ continuous on (a, b) , and assume $f^{(n)}$ is differentiable at c , i.e., assume $f^{(n+1)}(c)$ exists. Define the $(n + 1)$ st remainder

$$\begin{aligned} R_{n+1}(x, c) &= f(x) - \left[f(c) + f'(c)(x - c) \right. \\ &\quad \left. + \frac{f''(c)}{2!}(x - c)^2 + \cdots + \frac{f^{(n)}(c)}{n!}(x - c)^n \right], \quad a < x < b, \end{aligned}$$

and define $h_{n+1} : (a, b) \rightarrow \mathbf{R}$ by

$$h_{n+1}(x) = \begin{cases} \frac{R_{n+1}(x, c)}{(x - c)^{n+1}/(n + 1)!}, & x \neq c, \\ f^{(n+1)}(c), & x = c. \end{cases}$$

Then, h_{n+1} is continuous when $x \neq c$ and $R'_{n+1}(x, c; f) = R_n(x, c; f')$, where $R_n(x, c; f)$ denotes the remainder corresponding to f . Applying l'Hopital's rule n times,

$$\begin{aligned} \lim_{x \rightarrow c} h_{n+1}(x) &= \lim_{x \rightarrow c} \frac{R'_{n+1}(x, c)}{(x - c)^n / n!} \\ &= \lim_{x \rightarrow c} \frac{R_n(x, c; f')}{(x - c)^n / n!} = \lim_{x \rightarrow c} \frac{R_{n-1}(x, c; f'')}{(x - c)^{n-1} / (n-1)!} \\ &= \dots = \frac{R_1(x, c; f^{(n)})}{x - c} = \lim_{x \rightarrow c} \frac{f^{(n)}(x) - f^{(n)}(c)}{x - c} = f^{(n+1)}(c). \end{aligned}$$

Thus, h_{n+1} is continuous on (a, b) .

Taylor's Theorem. Let $n \geq 0$, and suppose that f is n times differentiable on (a, b) with $f^{(n)}$ continuous on (a, b) . If $f^{(n+1)}(c)$ exists at some fixed c in (a, b) , then, there is a continuous function¹⁴ $h_{n+1} : (a, b) \rightarrow \mathbb{R}$ satisfying $h_{n+1}(c) = f^{(n+1)}(c)$ and

$$\begin{aligned} f(x) &= f(c) + f'(c)(x - c) + \frac{f''(c)}{2!}(x - c)^2 + \dots \\ (4.3) \quad &\dots + \frac{f^{(n)}(c)}{n!}(x - c)^n + \frac{h_{n+1}(x)}{(n+1)!}(x - c)^{n+1}. \end{aligned}$$

Moreover, if $f^{(n+1)}$ exists on all of (a, b) , then, for some ξ between c and x , $R_{n+1}(x, c)$ is given by the Cauchy form

$$R_{n+1}(x, c) = \frac{h_{n+1}(x)}{(n+1)!}(x - c)^{n+1} = \frac{f^{(n+1)}(\xi)}{n!}(x - \xi)^n(x - c),$$

and, for some η between c and x , $R_{n+1}(x, c)$ is given by the Lagrange form

$$R_{n+1}(x, c) = \frac{h_{n+1}(x)}{(n+1)!}(x - c)^{n+1} = \frac{f^{(n+1)}(\eta)}{(n+1)!}(x - c)^{n+1}.$$

The derivation of (4.3) is above. To obtain the Cauchy and Lagrange forms, differentiate the expression defining $R_{n+1}(x, c)$ once with respect to c . Then, the sum collapses to (here, ' means derivative with respect to c)

$$R'_{n+1}(x, c) = -\frac{f^{(n+1)}(c)}{n!}(x - c)^n.$$

Now, apply the mean value theorem to $t \mapsto R_{n+1}(x, t)$ on the interval joining c to x . Since $R_{n+1}(x, x) = 0$, this yields the Cauchy form. To obtain the Lagrange form, set $g(t) = (x - t)^{n+1} / (n+1)!$. Then, $g'(t) = -(x - t)^n / n!$, $g(x) = 0$, and $R'_{n+1}(x, t) / g'(t) = f^{(n+1)}(t)$. So, by the generalized mean value theorem, there

¹⁴Taylor's theorem in §4.4 gives a useful formula for h_{n+1} .

is an η between c and x with

$$h_{n+1}(x) = \frac{R_{n+1}(x, x) - R_{n+1}(x, c)}{g(x) - g(c)} = \frac{R'_{n+1}(x, \eta)}{g'(\eta)} = f^{(n+1)}(\eta). \quad \square$$

In particular, the Lagrange form implies that, for f smooth on \mathbb{R} and any $n \geq 1$, for each $x \in \mathbb{R}$, there is an η between 0 and x with

$$(4.4) \quad f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \cdots + \frac{f^{(n)}(0)}{n!}x^n + \frac{f^{(n+1)}(\eta)}{(n+1)!}x^{n+1}.$$

Thus, a smooth function f can be approximated near 0 by an n th degree polynomial with an error $R_n(x, 0)$ given by a certain expression, for every $n \geq 1$.

If the remainder $R_n(x, 0)$ approaches 0 as $n \nearrow \infty$, we can let $n \nearrow \infty$ in (4.4) to obtain a series for f . Thus, if we know enough about $R_n(x, 0)$, we may be able to express f as an “infinite polynomial”

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \cdots + \frac{f^{(n)}(0)}{n!}x^n + \cdots$$

This series is the *Taylor series of f centered at 0*. For example, with $f(x) = e^x$, $f^{(n)}(x) = e^x$, hence, $f^{(n)}(0) = 1$ for all $n \geq 0$. Thus, the absolute value of the n th remainder satisfies

$$|R_n(x, 0)| = \frac{e^\eta}{(n+1)!}|x|^{n+1} \leq \frac{e^{|x|}}{(n+1)!}|x|^{n+1}, \quad n \geq 1.$$

But, now, $R_n(x, 0) \rightarrow 0$ follows from (3.5).

We have shown that

$$(4.5) \quad e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots, \quad x \in \mathbb{R}.$$

In particular, we have arrived at a series for e ,

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots$$

In §1.6 we obtained $2.5 < e \leq 3$. In fact, by the addition of sufficiently many terms, e can be computed to arbitrarily many places.

More generally, we call

$$f(x) = f(c) + f'(c)(x-c) + \frac{f''(c)}{2!}(x-c)^2 + \cdots + \frac{f^{(n)}(c)}{n!}(x-c)^n + \cdots$$

the *Taylor series¹⁵ centered at c* .

Using (4.4) with $f(x) = e^x$, we can show that e is irrational. Indeed, suppose that e were rational. Then, there would be a natural N , such that $n!e \in \mathbb{N}$ for all $n \geq N$. So, choose n greater than 3 and greater than N , and write (4.4) for this n and $x = 1$ to get

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!} + \frac{e^\eta}{(n+1)!}.$$

¹⁵When centered at zero, the Taylor series is also called the *Maclaurin series* of f .

Then

$$n!e = n! \left(1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!} \right) + \frac{e^\eta}{n+1}.$$

So, $e^\eta/(n+1)$ is a natural, which is false since $0 < \eta < 1$ and $n > 3$ imply

$$\frac{e^\eta}{n+1} < \frac{e}{4} \leq \frac{3}{4}.$$

This contradiction allows us to conclude the following.

Theorem. *e is irrational.* \square

The Taylor series of a smooth function may or may not converge for a given x . When it does converge, its sum need not equal the function. In fact, for each x , the Taylor series of f evaluated at x sums to $f(x)$ iff $R_n(x, 0) \rightarrow 0$. However, there are smooth functions f for which $R_n(x, 0) \not\rightarrow 0$. For example (Exercise 3), the function

$$f(x) = \begin{cases} e^{-1/x} & x > 0, \\ 0 & x \leq 0, \end{cases}$$

satisfies $f^{(n)}(0) = 0$ for all $n \geq 0$. It follows that $R_n(x, 0) = f(x)$ for all $n \geq 1$ and $x \in \mathbb{R}$. Thus, for all x , the Taylor series centered at zero converges, since it is identically zero, but $R_n(x, 0) \not\rightarrow 0$ when $x \neq 0$. Hence, the Taylor series does not sum to $f(x)$, except when $x = 0$.

Now, we turn to the general study of series involving powers of x as in (4.5). Below, we denote the series (4.5) by $\exp x$. Thus, $\exp x = e^x$. A *power series* is a series of the form

$$f(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$$

where the coefficients a_n , $n \geq 0$, are reals. We have seen that $f(x) = e^x$ can be expressed as a power series. What other functions can be so expressed? Two examples are the even and the odd parts of \exp .

The even and odd parts (§3.1) of \exp are the *hyperbolic cosine* \cosh and the *hyperbolic sine* \sinh (these are pronounced to rhyme with 'gosh' and 'cinch'). Thus,

$$\sinh x = \frac{e^x - e^{-x}}{2} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots,$$

and

$$\cosh x = \frac{e^x + e^{-x}}{2} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$$

Since $\sum_{n=0}^{\infty} |x^n/n!| = \sum_{n=0}^{\infty} |x|^n/n! = \exp |x|$, the series \exp is absolutely convergent on all of \mathbb{R} . By comparison with \exp , the series for \sinh and \cosh are

also absolutely convergent on all of \mathbb{R} . From their definitions, $\cosh' = \sinh$ and $\sinh' = \cosh$.

Note that the series for $\sinh x$ involves only odd powers of x and the series for $\cosh x$ involves only even powers of x . This holds for all odd and even functions (Exercise 14).

To obtain other examples, we write alternating versions (§1.7) of the last two series obtaining

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots,$$

and

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

These functions are studied in §3.5. Again, by comparison with \exp , \sin and \cos are absolutely convergent series on all of \mathbb{R} . However, unlike \exp , \cosh , and \sinh , we do not as yet know \sin' and \cos' .

In general, since a power series is a series involving a variable x , it may converge at some x 's and diverge at other x 's. For example (§1.6),

$$1 + x + x^2 + \dots \begin{cases} = \frac{1}{1-x}, & |x| < 1, \\ \text{diverges,} & |x| \geq 1. \end{cases}$$

Note that the set of x 's for which this series converges is an interval centered at zero. This is not an accident (Figure 3.10).

Theorem. *Let $\sum a_n x^n$ be a power series. Then, either the series converges absolutely for all x , the series converges only at $x = 0$, or there is an $R > 0$, such that the series converges absolutely if $|x| < R$ and diverges if $|x| > R$.*

To derive the theorem, let $R = \sup\{|x| : \sum a_n x^n \text{ converges}\}$. If $R = 0$, the series converges only for $x = 0$. If $R > 0$ and $|x| < R$, choose c with $|x| < |c| < R$ and $\sum a_n c^n$ convergent. Then, $\{a_n c^n\}$ is a bounded sequence by the n th term test, say $|a_n c^n| \leq C$, $n \geq 1$, and it follows that

$$\sum |a_n x^n| = \sum |a_n c^n| \cdot |x/c|^n \leq C \sum |x/c|^n < \infty,$$

since $|x/c| < 1$ and the last series is geometric. This shows that $\sum a_n x^n$ converges absolutely for all x in $(-R, R)$. On the other hand, if $R < \infty$, the definition of R shows that $\sum a_n x^n$ diverges for $|x| > R$. Finally, if $R = \infty$, this shows that the series converges absolutely for all x . \square




FIGURE 3.10. Region of convergence of a power series.

Note that the theorem says nothing about $x = R$ and $x = -R$. At these two points, anything can happen. For example, the power series

$$f(x) = 1 - \frac{x}{1} + \frac{x^2}{2} - \frac{x^3}{3} + \dots$$

has radius $R = 1$ and converges for $x = 1$ but diverges for $x = -1$. On the other hand, the series $f(-x)$ has $R = 1$ and converges for $x = -1$ but diverges for $x = 1$. The geometric series has $R = 1$ but diverges at $x = 1$ and $x = -1$, whereas

$$1 + \frac{x}{1^2} + \frac{x^2}{2^2} + \frac{x^3}{3^2} + \dots$$

has $R = 1$ and converges at $x = 1$ and $x = -1$. Because the interval of convergence is $(-R, R)$, the number R is called the *radius of convergence*. If the series converges only at $x = 0$, we say $R = 0$, whereas if the series converges absolutely for all x , we say $R = \infty$.

Here are two useful formulas for the radius R .

Theorem (Root Test). *Let $\sum a_n x^n$ be a power series and let L denote the upper limit of the sequence $(|a_n|^{1/n})$. Then, $R = 1/L$ where we take $1/0 = \infty$ and $1/\infty = 0$.*

To derive this, it is enough to show that $LR = 1$. If $|x| < R$, then, $|a_n| \cdot |x|^n = |a_n x^n| \leq C$ for some C , hence, $|a_n|^{1/n} |x| \leq C^{1/n}$. Taking the upper limit of both sides, we conclude that $L|x| \leq 1$. Since $|x|$ may be as close as desired to R , we obtain $LR \leq 1$. On the other hand, if $x > R$, then, $(a_n x^n)$ is unbounded (otherwise, the series converges on $(-x, x)$ contradicting the definition of R). Hence, some subsequence of $(|a_n| |x|^n)$ is bounded below by 1, which implies that some subsequence of $(|a_n|^{1/n} |x|)$ is bounded by 1. We conclude that $L|x| \geq 1$. Since $|x|$ may be as close as desired to R , we obtain $LR \geq 1$. Hence, $LR = 1$. \square

Theorem (Ratio Test). *Let (a_n) be a nonzero sequence, and suppose that*

$$\rho = \lim_{n \rightarrow \infty} |a_n|/|a_{n+1}|$$

exists. Then, ρ equals the radius of convergence R of $\sum a_n x^n$.

To show that $\rho = R$, we show that $\sum a_n x^n$ converges when $|x| < \rho$ and diverges when $|x| > \rho$. If $|x| < \rho$, choose c with $|x| < |c| < \rho$. Then, $|c| \leq |a_n|/|a_{n+1}|$ for $n \geq N$. Hence,

$$|a_{n+1} x^{n+1}| = \left| \frac{a_{n+1}}{a_n} \right| \cdot |c| \cdot |a_n x^n| \cdot \left| \frac{x}{c} \right| \leq |a_n x^n| \cdot \left| \frac{x}{c} \right|, \quad n \geq N.$$

Iterating this, we obtain $|a_{N+2} x^{N+2}| \leq |x/c|^2 |a_N x^N|$. Continuing in this manner, we obtain $|a_n x^n| \leq C |x/c|^{n-N}$, $n \geq N$, for some constant C . Since $\sum |x/c|^n$ converges, this shows that $\sum |a_n x^n|$ converges. On the other hand, if $|x| > \rho$,

then, the same argument shows that $|a_{n+1}x^{n+1}| \geq |a_nx^n|$ for $n \geq N$, hence, $a_nx^n \not\rightarrow 0$. By the n th term test, $\sum a_nx^n$ diverges. Thus, $\rho = R$. \square

A function that can be expressed as a power series with infinite radius of convergence is said to be *entire*. Thus \exp , \sinh , \cosh , \sin , and \cos are entire functions. Clearly the sum of entire functions is entire. Since the Cauchy product (Exercise 7 of §1.7) of absolutely convergent series is absolutely convergent, the product of entire functions is entire. It is not true, however, that the quotient of entire functions is entire. For example, $f(x) = 1$ divided by $g(x) = 1 + x^2$ is smooth but not entire (see below).

Using the formula for the radius of convergence together with the fact that \exp converges everywhere yields

$$(4.6) \quad \lim_{n \rightarrow \infty} (n!)^{1/n} = \infty.$$

This can also be derived directly.

It turns out that functions constructed from power series are smooth in their interval of convergence. They have derivatives of all orders.

Theorem. Let $f(x) = \sum a_nx^n$ be a power series with radius of convergence $R > 0$. Then,

$$(4.7) \quad \sum_{n=1}^{\infty} na_nx^{n-1} = a_1 + 2a_2x + 3a_3x^2 + \dots$$

has radius of convergence R , f is differentiable on $(-R, R)$, and $f'(x)$ equals (4.7) for all x in $(-R, R)$.

In other words, to obtain the derivative of a power series, one need only differentiate the series term by term. To see this, we first show that the radius of the power series $\sum (n+1)a_{n+1}x^n$ is R . Here, the n th coefficient is $b_n = (n+1)a_{n+1}$, so,

$$|b_n|^{1/n} = (n+1)^{1/n} |a_{n+1}|^{1/n} = (n+1)^{1/n} [|a_{n+1}|^{1/(n+1)}]^{(n+1)/n},$$

so, the upper limit of $(|b_n|^{1/n})$ equals the upper limit of $(|a_n|^{1/n})$ since $(n+1)^{1/n} \rightarrow 1$ (§3.3) and $(n+1)/n \rightarrow 1$.

Now we show that $f'(c)$ exists and equals $\sum na_n c^{n-1}$, where $-R < c < R$ is fixed. To do this, let us consider only a single term in the series, i.e., let us consider x^n with n fixed, and pick $|c| < R$. Then, by the binomial theorem (§3.3),

$$\begin{aligned} x^n &= [c + (x - c)]^n = \sum_{j=0}^n \binom{n}{j} c^{n-j} (x - c)^j \\ &= c^n + nc^{n-1}(x - c) + \sum_{j=2}^n \binom{n}{j} c^{n-j} (x - c)^j. \end{aligned}$$

Thus,

$$\frac{x^n - c^n}{x - c} - nc^{n-1} = \sum_{j=2}^n \binom{n}{j} c^{n-j} (x - c)^{j-1}, \quad x \neq c.$$

Now, choose $d > 0$ with $|c| + d < R$. Then, for x satisfying $0 < |x - c| < d$,

$$\begin{aligned} \left| \frac{x^n - c^n}{x - c} - nc^{n-1} \right| &= \left| \sum_{j=2}^n \binom{n}{j} c^{n-j} (x - c)^{j-1} \right| \\ &\leq |x - c| \sum_{j=2}^n \binom{n}{j} |c|^{n-j} |x - c|^{j-2} \\ &\leq |x - c| \sum_{j=2}^n \binom{n}{j} |c|^{n-j} d^{j-2} \\ &= \frac{|x - c|}{d^2} \sum_{j=2}^n \binom{n}{j} |c|^{n-j} d^j \\ &\leq \frac{|x - c|}{d^2} (|c| + d)^n, \end{aligned}$$

where we have used the binomial theorem again. To summarize,

$$(4.8) \quad \left| \frac{x^n - c^n}{x - c} - nc^{n-1} \right| \leq \frac{|x - c|}{d^2} (|c| + d)^n, \quad 0 < |x - c| < d.$$

Assume, temporarily, that the coefficients a_n are nonnegative, $a_n \geq 0$, $n \geq 0$. Multiplying (4.8) by a_n and summing over $n \geq 0$ yields

$$\left| \frac{f(x) - f(c)}{x - c} - \sum_{n=1}^{\infty} na_n c^{n-1} \right| \leq \frac{|x - c|}{d^2} f(|c| + d), \quad 0 < |x - c| < d.$$

Letting $x \rightarrow c$ in the last inequality establishes the result when $a_n \geq 0$, $n \geq 0$. If this is not so, the same argument works, except for a slight modification in the right side of this last inequality, which we leave to the reader. \square

Since this theorem can be applied repeatedly to f , f' , f'' , \dots , every power series with radius of convergence R determines a smooth function f on $(-R, R)$. For example, $f(x) = \sum a_n x^n$ implies

$$f''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2}$$

on the interval of convergence. It follows that every entire function is smooth on all of \mathbb{R} . The converse, however, is not generally true. For example,

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \dots$$

converges only on the interval $(-1, 1)$, but $1/(1+x^2)$ is smooth on \mathbb{R} (Exercise 7).

In particular, \sin and \cos are smooth functions on \mathbb{R} (we already knew this for \exp , \cosh , and \sinh). Moreover, differentiating the series for \sin and \cos , term by term, yields $\sin' = \cos$ and $\cos' = -\sin$.

Our last topic is to describe Newton's generalization of the binomial theorem (§3.3) to nonnatural exponents. The result is the same, except that the sum proceeds to ∞ . For v real and $n \geq 1$, let

$$\binom{v}{n} = \frac{v \cdot (v-1) \cdot \dots \cdot (v-n+1)}{1 \cdot 2 \cdot \dots \cdot n}.$$

Also let $\binom{v}{0} = 1$. If v is a natural, then, $\binom{v}{n}$ is the binomial coefficient defined previously for $0 \leq n \leq v$ and $\binom{v}{n} = 0$ for $n > v$. Then, for any natural N with $n > N \geq v$,

$$\begin{aligned} \left| \binom{v}{n+1} \right| &\leq \frac{N \cdot (N+1) \cdot \dots \cdot (N+n)}{1 \cdot 2 \cdot \dots \cdot (n+1)} \\ &\leq \frac{1 \cdot 2 \cdot \dots \cdot N \cdot \dots \cdot (N+n)}{1 \cdot 2 \cdot \dots \cdot (n+1)} \\ &= (n+2) \cdot \dots \cdot (n+N-1) \cdot (n+N) \leq (n+N)^{N-1}. \end{aligned}$$

Theorem. For v real,

$$(1+x)^v = \sum_{n=0}^{\infty} \binom{v}{n} x^n, \quad -1 < x < 1.$$

To see this, fix a natural $N > v$ and apply the Lagrange form of the remainder in Taylor's theorem to $f(x) = (1+x)^v$ and $0 < x < 1$, $c = 0$. Then, $f'(x) = v(1+x)^{v-1}$, $f''(x) = v(v-1)(1+x)^{v-2}$, and

$$f^{(n)}(x) = v(v-1) \dots (v-n+1)(1+x)^{v-n}.$$

So, $f^{(n)}(x)/n! = \binom{v}{n}(1+x)^{v-n}$. Hence, for some $0 < \eta < x < 1$ (see §3.3 for the limits), we obtain

$$|R_n(x, c)| = \left| \binom{v}{n+1} \right| (1+\eta)^{v-n-1} x^{n+1} \leq (N+n)^{N-1} x^{n+1} \rightarrow 0, n \nearrow \infty.$$


Since $f^{(n)}(0)/n! = \binom{v}{n}$, $n \geq 0$, the Taylor series yields

$$(1+x)^v = \sum_{n=0}^{\infty} \binom{v}{n} x^n, \quad 0 < x < 1,$$

which establishes the theorem on $(0, 1)$.

To establish the result on $(-1, 0)$, apply the Cauchy form of the remainder to $f(x) = (1-x)^v$ and $0 < x < 1$, $c = 0$. Then, $f^{(n+1)}(x)/n! = (-1)^{n+1} \binom{v}{n+1} (1-x)^{v-n-1} (n+1)$, so, for some $0 < \xi < x$, we obtain

$$|R_n(x, c)| = (n+1) \left| \binom{v}{n+1} \right| (1-\xi)^{v-n-1} (x-\xi)^n (x-0)$$

FIGURE 3.11. Graph of the function h in Exercise 3.

$$\begin{aligned}
 &= (n+1)x \left| \binom{v}{n+1} \right| \left(\frac{x-\xi}{1-\xi} \right)^n (1-\xi)^{v-1} \\
 &\leq (n+1)x(N+n)^{N-1} \left(\frac{x-\xi}{1-\xi} \right)^n (1-\xi)^{v-1}.
 \end{aligned}$$

If $v \geq 1$, $(1-\xi)^{v-1} \leq 1$. If $v < 1$, $(1-\xi)^{v-1} \leq (1-x)^{v-1}$. Hence, in both cases, $(1-\xi)^{v-1}$ is bounded by $[1 + (1-x)^{v-1}]$, a fixed quantity independent of n (remember that ξ may depend on n). Moreover, $(x-\xi)/(1-\xi) < (x-0)/(1-0) = x$. Hence,

$$|R_n(x, c)| \leq [1 + (1-x)^{v-1}](n+1)(N+n)^{N-1}x^{n+1},$$

which goes to zero as $n \nearrow \infty$. Since $f^{(n)}(0)/n! = (-1)^n \binom{v}{n}$, $n \geq 0$, the Taylor series yields

$$(1-x)^v = \sum_{n=0}^{\infty} (-1)^n \binom{v}{n} x^n = \sum_{n=0}^{\infty} \binom{v}{n} (-x)^n, \quad 0 < x < 1,$$

which establishes the theorem on $(-1, 0)$. \square

Exercises 3.4

1. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is nonnegative, twice differentiable, and $f''(c) \leq 1/2$ for all $c \in \mathbb{R}$. Use Taylor's theorem to conclude that $|f'(c)| \leq \sqrt{f(c)}$. (Choose $x = c + t$, and note that $f(c+t) \geq 0$.)
2. Use the exponential series to compute e to four decimal places, justifying your reasoning.
3. ★ Let

$$h(x) = \begin{cases} e^{-1/x}, & x > 0, \\ 0, & x \leq 0. \end{cases}$$

By induction on $n \geq 1$, show that

- (1) $h^{(n-1)}(x) = h(x)R_n(x)$, $x > 0$, for some rational function R_n , and
- (2) $h^{(n-1)}(x) = 0$, $x \leq 0$.

Conclude that h (Figure 3.11) is smooth on \mathbb{R} . (Here, do *not* try to compute R_n .)

4. Show directly that $\lim_{n \rightarrow \infty} (n!)^{1/n} = \infty$. (First, show that the lower limit is ≥ 100 .)

5. Show that

$$\frac{1}{\sqrt{1-x^2}} = 1 + \frac{1}{2} \cdot x^2 + \frac{1}{2} \cdot \frac{3}{4} \cdot x^4 + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot x^6 + \dots$$

for $|x| < 1$.

6. Compute the Taylor series of $\log(1+x)$ centered at 0.

7. Suppose that

$$(4.9) \quad f(x) = a_0 + a_1x + a_2x^2 + \dots$$

converges for $|x| < R$. Show that the series in (4.9) is the Taylor series of f centered at 0. Conclude that $\sum a_n x^n = \sum b_n x^n$ for $|x| < R$ implies $a_n = b_n$ for $n \geq 0$. Thus, the coefficients of a power series are uniquely determined.

8. Show that

$$\frac{\log(1+x)}{1+x} = x - \left(1 + \frac{1}{2}\right)x^2 + \left(1 + \frac{1}{2} + \frac{1}{3}\right)x^3 - \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}\right)x^4 + \dots$$

for $|x| < 1$ by considering the product (§1.7) of the series for $1/(1+x)$ and the series for $\log(1+x)$.

9. ★ Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable with f , f' , and f'' continuous. If $f(0) = 1$, $f'(0) = 0$ and $f''(0) = q$, use Taylor's theorem and Exercise 3 of §3.2 to show that

$$\lim_{n \rightarrow \infty} [f(x/\sqrt{n})]^n = \exp(qx^2/2).$$

This result is a key step in the derivation of the *central limit theorem*.

10. ★ Show that the inverse $\operatorname{arcsinh} : \mathbb{R} \rightarrow \mathbb{R}$ of $\sinh : \mathbb{R} \rightarrow \mathbb{R}$ exists and is smooth, and compute $\operatorname{arcsinh}'$. Show that $f(x) = \cosh x$ is superlinear, smooth and strictly convex. Compute the conjugate $g(y)$ (Exercise 11 of §3.3), and check that g is smooth.

11. Compute the radius of convergence of $\sum_{n=0}^{\infty} (-1)^n x^n / 4^n (n!)^2$.

12. ★ What is the radius of convergence of

$$\sum_{n=0}^{\infty} x^{n!} = 1 + x + x^2 + x^6 + x^{24} + x^{120} + \dots?$$

13. Show that

$$\binom{-1/2}{n} = \frac{(-1)^n (2n)!}{4^n (n!)^2}, \quad n \geq 0.$$

14. Let $f(x) = a_0 + a_1x + a_2x^2 + \dots$ converge for $-R < x < R$. Then, f is even iff the coefficients a_{2n-1} , $n \geq 1$, of the odd powers vanish, and f is odd iff the coefficients a_{2n} , $n \geq 1$, of the even powers vanish.

15. Compute

$$\lim_{t \rightarrow 0} \left(\frac{1}{e^t - 1} - \frac{1}{t} \right)$$

by writing $e^t = 1 + t + t^2 h(t)/2$.

§3.5 Trigonometry

In the previous section, we introduced alternating versions of the even and the odd parts of the exponential series, the *sine function*, and the *cosine function*,

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots,$$

and

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

Since these functions are *defined* by these convergent power series, they are smooth everywhere and satisfy (§3.4):

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x,$$

$\sin 0 = 0$, and $\cos 0 = 1$. The sine function is odd and the cosine function is even,

$$\sin(-x) = -\sin x,$$

and

$$\cos(-x) = \cos x.$$

Since

$$(\sin^2 x + \cos^2 x)' = 2 \sin x \cos x + 2 \cos x (-\sin x) = 0,$$

evaluating at $x = 0$ yields

$$(5.1) \quad \sin^2 x + \cos^2 x = 1.$$

If a is a critical point of $\sin x$, then, $\cos a = 0$, hence, $\sin a = \pm 1$. Hence, $\sin a = 1$ at any positive local maximum a of $\sin x$.

Let f, g be differentiable functions satisfying $f' = g$ and $g' = -f$ on \mathbf{R} . Now, the derivatives of $f \sin + g \cos$ and $f \cos - g \sin$ vanish, hence,

$$f(x) \sin x + g(x) \cos x = g(0),$$

and

$$f(x) \cos x - g(x) \sin x = f(0)$$

for all x . Multiplying the first equation by $\sin x$ and the second by $\cos x$ and adding, we obtain

$$f(x) = g(0) \sin x + f(0) \cos x.$$

Multiplying the first by $\cos x$ and the second by $-\sin x$ and adding, we obtain

$$g(x) = g(0) \cos x - f(0) \sin x.$$

Fixing y and taking $f(x) = \sin(x + y)$ and $g(x) = \cos(x + y)$, we obtain the identities

$$(5.2) \quad \begin{aligned} \sin(x + y) &= \sin x \cos y + \cos x \sin y, \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y. \end{aligned}$$

If we replace y by $-y$ and combine the resulting equations with (5.2), we obtain the identities

$$(5.3) \quad \begin{aligned} \sin(x + y) + \sin(x - y) &= 2 \sin x \cos y, \\ \sin(x + y) - \sin(x - y) &= 2 \cos x \sin y, \\ \cos(x - y) - \cos(x + y) &= 2 \sin x \sin y, \\ \cos(x - y) + \cos(x + y) &= 2 \cos x \cos y. \end{aligned}$$

For $0 \leq x \leq 3$, the series

$$x - \sin x = \frac{x^3}{3!} - \frac{x^5}{5!} + \frac{x^7}{7!} - \dots$$

is alternating *with decreasing terms*. Hence, by the Leibnitz test (§1.7),

$$x - \sin x \leq \frac{x^3}{6}, \quad 0 \leq x \leq 3,$$

and

$$x - \sin x \geq \frac{x^3}{6} - \frac{x^5}{120}, \quad 0 \leq x \leq 3.$$

Inserting $x = 1$ in the first inequality and $x = 3$ in the second, we obtain

$$\sin 0 = 0, \quad \sin 1 \geq \frac{5}{6}, \quad \sin 3 \leq \frac{21}{40}.$$

Hence, there is a positive b in $(0, 3)$, where $\sin b$ is a positive maximum, which gives $\cos(b) = 0$ and $\sin(b) = 1$. Let $a = \inf\{b > 0 : \sin b = 1\}$. Then, the continuity of $\sin x$ implies $\sin a = 1$. Since $\sin 0 = 0$, $a > 0$. Since \sin is a specific function, a is a specific real number.

For more than 20 centuries, the real $2a$ has been called “Archimedes’ constant.” More recently, since the eighteenth century, the greek letter π has been used to denote this real. Thus,

$$\sin\left(\frac{\pi}{2}\right) = 1,$$

and

$$\cos\left(\frac{\pi}{2}\right) = 0.$$

As yet, all we know about π is $0 < \pi/2 < 3$. In §5.2, we address the issue of computing π accurately.

Since the slope of the tangent line of $\sin x$ at $x = 0$ is $\cos 0 = 1$, $\sin x > 0$ for $x > 0$ near 0. Since there are no positive local maxima for $\sin x$ in $(0, \pi/2)$ and $\sin 0 = 0$, we must have $\sin x > 0$ on $(0, \pi/2)$. Hence, $\cos x$ is strictly decreasing on $(0, \pi/2)$. Hence, $\cos x$ is positive on $(0, \pi/2)$. Hence, $\sin x$ is strictly increasing on $(0, \pi/2)$. Moreover, since $(\sin x)'' = -\sin x$, and $(\cos x)'' = -\cos x$, $\sin x$ and $\cos x$ are concave on $(0, \pi/2)$. This justifies the graphs of $\sin x$ and $\cos x$ in the interval $[0, \pi/2]$ (Figure 3.12).

Inserting $y = \pi$ and replacing x by $-x$ in (5.2) yields

$$(5.4) \quad \begin{aligned} \sin(\pi - x) &= \sin x, \\ \cos(\pi - x) &= -\cos x. \end{aligned}$$

These identities justify the graphs of $\sin x$ and $\cos x$ in the interval $[\pi/2, \pi]$. Hence, the graphs are now justified on $[0, \pi]$. Replacing x by $-x$ in (5.4), we obtain

$$\sin(x + \pi) = -\sin x,$$

and

$$\cos(x + \pi) = -\cos x.$$

These identities justify the graphs of $\sin x$ and $\cos x$ on $[0, 2\pi]$. Repeating this reasoning once more,

$$\sin(x + 2\pi) = \sin(x + \pi + \pi) = -\sin(x + \pi) = \sin x,$$

and

$$\cos(x + 2\pi) = \cos(x + \pi + \pi) = -\cos(x + \pi) = \cos x,$$

showing that 2π is a *period* of $\sin x$ and $\cos x$. In fact, repeating this reasoning,

$$\sin(x + 2\pi n) = \sin x, \quad n \in \mathbf{Z},$$

and

$$\cos(x + 2\pi n) = \cos x, \quad n \in \mathbf{Z},$$

showing that every integer multiple of 2π is a period of $\sin x$ and $\cos x$. If $\sin x$ or $\cos x$ had any other period p , then, by subtracting from p an appropriate integral multiple of 2π , we would obtain a period in $(0, 2\pi)$, contradicting the graphs. Hence, $2\pi\mathbf{Z}$ is the set of periods of $\sin x$ and of $\cos x$.

If we set $x = y$ in (5.2), we obtain

$$\sin(2x) = 2 \sin x \cos x,$$

and

$$\cos(2x) = \cos^2 x - \sin^2 x.$$

By (5.1), the second identity implies

$$\cos^2 x = \frac{1 + \cos(2x)}{2},$$

and

$$\sin^2 x = \frac{1 - \cos(2x)}{2}.$$




FIGURE 3.12. The graphs of sine and cosine.

By the inverse function theorem, $\sin x$ has an inverse on $[-\pi/2, \pi/2]$ and $\cos x$ has an inverse on $[0, \pi]$. These inverses are $\arcsin : [-1, 1] \rightarrow [-\pi/2, \pi/2]$ and $\arccos : [-1, 1] \rightarrow [0, \pi]$. Since $\cos x > 0$ on $(-\pi/2, \pi/2)$, by (5.1), $\cos(\arcsin x) = \sqrt{1 - x^2}$ on $[-1, 1]$. Similarly, since $\sin x > 0$ on $(0, \pi)$, $\sin(\arccos x) = \sqrt{1 - x^2}$ on $[-1, 1]$. Thus, the derivatives of the inverse functions are given by

$$(\arcsin x)' = \frac{1}{\cos(\arcsin x)} = \frac{1}{\sqrt{1 - x^2}}, \quad -1 < x < 1,$$

and


$$(\arccos x)' = \frac{1}{-\sin(\arccos x)} = \frac{-1}{\sqrt{1 - x^2}}, \quad -1 < x < 1.$$

As an application,

$$(2 \arcsin \sqrt{x})' = 2 \cdot \frac{1}{\sqrt{1 - \sqrt{x}^2}} \cdot \frac{1}{2\sqrt{x}} = \frac{1}{\sqrt{x(1 - x)}}, \quad 0 < x < 1.$$

Now, we make the connection with the unit circle. The *unit circle* is the subset $\{(x, y) : x^2 + y^2 = 1\}$ of \mathbb{R}^2 (§2.1). The interior $\{(x, y) : x^2 + y^2 < 1\}$ of the unit circle is the (*open*) *unit disk*.

Theorem. *If (x, y) is a point on the unit circle, there is a real θ with $(x, y) = (\cos \theta, \sin \theta)$. If ϕ is any other such real, then, $\theta - \phi$ is an integral multiple of*

FIGURE 3.13. The graphs of $\tan x$ and $\arctan x$.

2π . If (x, y) is any point in \mathbb{R}^2 , then, $(x, y) = (r \cos \theta, r \sin \theta)$ for some uniquely determined $r \geq 0$ and real θ , with θ determined up to an additive multiple of 2π , when $r > 0$.

Since $x^2 + y^2 = 1$, $|x| \leq 1$. Let $\theta = \arccos x$. Then, $\sin^2 \theta + \cos^2 \theta = 1$ implies $y = \pm \sin \theta$. If $y = \sin \theta$, we have found a real θ , as required. Otherwise, replace θ by $-\theta$. This does not change $\cos \theta = \cos(-\theta)$, but changes $\sin \theta = -\sin(-\theta)$. For the second statement, suppose that $(\cos \theta, \sin \theta) = (\cos \phi, \sin \phi)$. Then,

$$\cos(\theta - \phi) = \cos \theta \cos \phi + \sin \theta \sin \phi = 1.$$

Hence, $\theta - \phi$ is a multiple of 2π . For general (x, y) , set $r = \sqrt{x^2 + y^2}$. If $r = 0$, any θ yields $(x, y) = (r \cos \theta, r \sin \theta)$, whereas, if $r > 0$, $(x/r, y/r)$ is on the unit circle, so, we can choose θ by the first part. \square

Of course (r, θ) are the usual *polar coordinates* of the point (x, y) .

The *tangent function*, $\tan x = \sin x / \cos x$, is smooth everywhere except at odd multiples of $\pi/2$ where the denominator vanishes. Moreover, $\tan x$ is an odd function, and

$$\tan(x + \pi) = \frac{\sin(x + \pi)}{\cos(x + \pi)} = \frac{-\sin x}{-\cos x} = \tan x.$$

So, π is the period for $\tan x$. By the quotient rule

$$(\tan x)' = \frac{(\sin x)' \cos x - \sin x (\cos x)'}{(\cos x)^2} = \frac{1}{\cos^2 x}, \quad -\pi/2 < x < \pi/2.$$

Thus, $\tan x$ is strictly increasing on $(-\pi/2, \pi/2)$. Moreover,

$$\begin{aligned} \tan(\pi/2-) &= \infty, \\ \tan(-\pi/2+) &= -\infty, \end{aligned}$$

and

$$(\tan x)'' = \left(\frac{1}{\cos^2 x} \right)' = 2 \frac{\tan x}{\cos^2 x}.$$

Thus, $\tan x$ is convex on $(0, \pi/2)$ and concave on $(-\pi/2, 0)$. The graph is as shown in Figure 3.13.

By the inverse function theorem, $\tan x$ has an inverse on $(-\pi/2, \pi/2)$. This inverse, $\arctan : (-\infty, \infty) \rightarrow (-\pi/2, \pi/2)$, is smooth and

$$(\arctan x)' = 1/\cos(\arctan x)^{-2} = \cos^2(\arctan x).$$

Since $\cos x$ is positive on $(-\pi/2, \pi/2)$, dividing (5.1) by $\cos^2 x$, we have $\tan^2 x + 1 = 1/\cos^2 x$. Hence, $\cos^2 x = 1/(1 + \tan^2 x)$. Thus,

$$(\arctan x)' = \frac{1}{1 + x^2}.$$

It follows that $\arctan x$ is strictly increasing on \mathbb{R} . Since $(\arctan x)'' = -2x/(1 + x^2)^2$, $\arctan x$ is convex for $x < 0$ and concave for $x > 0$. Moreover, $\arctan \infty = \pi/2$ and $\arctan(-\infty) = -\pi/2$. The graph is as shown in Figure 3.13.

Often, we will use the convenient abbreviations $\sec x = 1/\cos x$, $\csc x = 1/\sin x$, and $\cot x = 1/\tan x$. These are the *secant*, *cosecant*, and *cotangent* functions. For example, $(\tan x)' = \sec^2 x$, and $(\cot x)' = -\csc^2 x$.

If $t = \tan(\theta/2)$, we have the *half-angle formulas*

$$\begin{aligned}\sin \theta &= \frac{2t}{1 + t^2}, \\ \cos \theta &= \frac{1 - t^2}{1 + t^2},\end{aligned}$$

and

$$\tan \theta = \frac{2t}{1 - t^2}.$$

The rest of the section is a review of euclidean geometry in the plane. The key concept is that of a euclidean motion.

Let $(a, b) \in \mathbb{R}^2$. A *translation* or a *translation by* (a, b) is the mapping $T = T_{(a,b)} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$(x, y) \mapsto (x + a, y + b).$$

Let θ be a real. A *rotation* or a *rotation by* θ is the mapping $R = R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$(x, y) \mapsto (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta).$$

A *euclidean motion* is a translation or a rotation or a composition of the two.

The basic group properties of euclidean motions are as follows. If T, T' are translations by $(a, b), (a', b')$, then, $T \circ T'$ is a translation by $(a + a', b + b')$. If R, R' are rotations by θ, θ' , then, $R \circ R'$ is a rotation by $\theta + \theta'$ (this follows from (5.2)). The translation by $(0, 0)$ and the rotation by 0 are both the identity mapping and the inverses of $T_{(a,b)}$ and R_θ are $T_{(-a,-b)}$ and $R_{-\theta}$, respectively.

If $A = (x, y)$ and $A' = (x', y')$ are points in \mathbb{R}^2 , their *sum* is $A + A' = (x + x', y + y')$. If k is a real and $A = (x, y)$ is a point in \mathbb{R}^2 , its *multiple* kA is the point (kx, ky) . If $k > 0$, kA is also called a *dilate* of A . The origin is $O = (0, 0)$, and a point is *nonzero* if $A \neq O$. The set of all nonnegative multiples $\{kA : k \geq 0\}$ of a fixed nonzero point A , by definition, is the *ray through* A .

If $A = (x, y)$ and $A' = (x', y')$ are any two points in \mathbb{R}^2 , the *distance between A and A'* is

$$|AA'| = \sqrt{(x - x')^2 + (y - y')^2}.$$

Note that the distance between A and A' equals the distance between $A - A'$ and O . If a translation sends A and B to A' and B' , then, the distance formula shows that $|AB| = |A'B'|$. Thus, *distances are unchanged by translation*.

If a rotation sends A, B to A' and B' , then, it sends $A + B$ and kA to $A' + B'$ and kA' . This *linearity* property follows from the definition of R_θ . It follows that rotations send rays to rays. If $A = (x, y)$ is rotated into $A' = (x', y')$, then $|OA| = |OA'|$ since

$$x'^2 + y'^2 = (x \cos \theta - y \sin \theta)^2 + (x \sin \theta + y \cos \theta)^2 = x^2 + y^2.$$

In particular, this is so when $|OA| = 1$, i.e., when A is on the unit circle. Hence, rotations send the unit circle into itself. If a rotation sends A and B to A' and B' , then,

$$|AB| = |O(A - B)| = |O(A - B)'| = |O(A' - B')| = |A'B'|.$$

Thus, *distances are unchanged by rotation*.

Given any two nonzero points A, B we define the *angle $\angle AOB$* (with vertex at O). We say $\angle AOB = \theta$ if R_θ maps the ray through A to the ray through B (Figure 3.14). Note that this notion of angle is *oriented*, i.e., $\angle AOB = -\angle BOA$. Moreover, $\angle AOB$ is defined up to an additive multiple of 2π and $\angle AOB = \angle A'OB'$ for any dilates A', B' of A, B , since the rays through A and B equal the rays through A' and B' , respectively. Since $R_{\alpha+\beta} = R_\alpha \circ R_\beta$, angles add as they should, i.e., for any three nonzero points A, B , and C , $\angle AOB + \angle BOC = \angle AOC$.




FIGURE 3.14. Definition of the angle $\angle AOB = \theta$.

More generally, given any three points A, B , and C with $A \neq B, C \neq B$, we define $\angle ABC$ (with vertex at B) to be $\angle A'OC'$ where $A' = A - B$ and $C' = C - B$.

If a translation sends A, B , and C to A', B' , and C' , then, $A - B = A' - B'$ and $C - B = C' - B'$. Hence,

$$\angle ABC = \angle(A - B)O(C - B) = \angle(A' - B')O(C' - B') = \angle A'B'C'.$$

Thus, *angles are unchanged by translation.*

If a rotation sends nonzero points A, B, C to A', B', C' , then, $\angle AOA' = \angle BOB'$. Hence,

$$\begin{aligned}\angle AOB &= \angle AOA' + \angle A'OB' + \angle B'OB \\ &= \angle AOA' + \angle A'OB' - \angle BOB' = \angle A'OB'.\end{aligned}$$

Now, the rotation sends $A - B$ to $A' - B'$ and $C - B$ to $C' - B'$. Hence,

$$\angle ABC = \angle(A - B)O(C - B) = \angle(A' - B')O(C' - B') = \angle A'B'C'.$$

Thus, *angles are unchanged by rotation.*

Using the euclidean invariance of distance and angle, one can derive any euclidean identity by reducing it to a simple situation. For example, given two points $A = (x, y)$ and $A' = (x', y')$, their *inner product* is

$$\langle OA, OA' \rangle = xx' + yy'.$$

Then, $\langle OA, OA' \rangle$ is unchanged by rotation (Exercise 8). It follows that

$$(5.5) \quad \langle OA, OA' \rangle = |OA| |OA'| \cos(\angle AOA').$$

To see this, note that both sides are invariant under rotation, hence, we may assume $A = (r \cos \theta, r \sin \theta)$, $A' = (r', 0)$. The result now falls out since both sides equal $rr' \cos \theta$.

Exercises 3.5

1. Derive the Cauchy–Schwarz inequality

$$\langle OA, OA' \rangle^2 \leq |OA|^2 |OA'|^2$$

directly, i.e., without using (5.5).

2. Derive the half-angle formulas.

3. Let $f(x) = x \sin(1/x)$, $x \neq 0$, $f(0) = 0$. Show that f is continuous at all points, but is not of bounded variation on any interval (a, b) containing 0.

4. Let $f(x) = x^2 \sin(1/x)$, $x \neq 0$, $f(0) = 0$. Show that f is differentiable at all points with $|f'(x)| \leq 1 + 2|x|$. Show that f is of bounded variation on any bounded interval.

5. Show that the composition of a rotation by ϕ with a rotation by θ is a rotation by $\phi + \theta$.

6. For any three distinct points A, B , and C , $\angle ABC + \angle BCA + \angle CAB = \pi$.

7. Show that, for $0 \leq x \leq 3$, the series

$$\frac{x^3}{3!} - \frac{x^5}{5!} + \frac{x^7}{7!} - \dots$$

has decreasing terms.

8. Given two points $A = (x, y)$ and $A' = (x', y')$, let $\langle OA, OA' \rangle = xx' + yy'$. Show that $\langle OA, OA' \rangle$ is unchanged by rotation.

9. For any three distinct points A, B , and C ,

$$|AB|^2 + |BC|^2 - 2|AB||BC|\cos(\angle ABC) = |AC|^2.$$

(First, assume that $B = O$ and expand the squares in $|AC|^2$.)

10. Show that $\cos(\pi/9)\cos(2\pi/9)\cos(4\pi/9) = 1/8$.

11. Compute the sine, cosine and tangent of $\pi/6, \pi/4$, and $\pi/3$.

12. ★ Use (5.2) and induction to show that

$$1 + 2\cos x + 2\cos(2x) + \cdots + 2\cos(nx) = \frac{\sin\left[\left(n + \frac{1}{2}\right)x\right]}{\sin\left(\frac{1}{2}x\right)}$$

for $n \geq 1$ and $x \notin 2\pi\mathbb{Z}$.

13. Show that $2\cot(2x) = \cot x - \tan x$.

14. Show that

$$(x^2 - 2x\cos\theta + 1) \cdot (x^2 - 2x\cos(\pi - \theta) + 1) = (x^4 - 2x^2\cos(2\theta) + 1).$$

Use this to derive the identity

$$(n) \quad x^{2n} - 1 = (x^2 - 1) \cdot \prod_{k=1}^{n-1} (x^2 - 2x\cos(k\pi/n) + 1)$$

for $n = 2, 4, 8, 16, \dots$. Here, $\prod_{k=1}^{n-1} a_k$ means $a_1 a_2 \dots a_{n-1}$. (Assuming (n) is true, establish (2n).)

15. ★ Use the Dirichlet test (§1.7) to show that $\sum_{n=1}^{\infty} \cos(nx)/n$ is convergent for $x \notin 2\pi\mathbb{Z}$.

§3.6 Primitives

Let f be defined on (a, b) . A differentiable function F is a *primitive* of f if

$$F'(x) = f(x), \quad a < x < b.$$

For example, $f(x) = x^3$ has the primitive $F(x) = x^4/4$ on \mathbb{R} since $(x^4/4)' = (4x^3)/4 = x^3$.

Not every function has a primitive on a given open interval (a, b) . Indeed, if $f : (a, b) \rightarrow \mathbb{R}$ has a primitive F on (a, b) , then, by Exercise 8 of §3.2, $f = F'$ satisfies the intermediate value property. Hence, $f((a, b))$ must be an interval.

Moreover, Exercise 6 of §3.1 shows that the presence of a jump discontinuity in f at a single point in (a, b) is enough to prevent the existence of a primitive F on (a, b) . In other words, if f is defined on (a, b) and $f(c+)$, $f(c-)$ exist but are not both equal to $f(c)$, for some $c \in (a, b)$, then, f has no primitive on (a, b) .

Later (§4.4), we see that *every continuous function has a primitive* on any open interval of definition.

Now, we investigate the converse of the last statement: To what extent does the existence of a primitive F of f determine the continuity of f ? To begin, it is possible (Exercise 7) for a function f to have a primitive and to be discontinuous at some points, so, the converse is, in general, false. However, the previous paragraph shows that such discontinuities cannot be jump discontinuities but must be wild, in the terminology of §2.3. In fact, it turns out that, wherever f is of bounded variation, the existence of a primitive forces the continuity of f (Exercise 8). Thus, a function f that has a primitive on (a, b) and is discontinuous at a particular point $c \in (a, b)$ must have unbounded variation near c , i.e., must be similar to the example in Exercise 7.

From the mean value theorem, we have the following simple but fundamental fact.

Theorem. *Any two primitives of f differ by a constant.*

Indeed, if F and G are primitives of f , then, $H = F - G$ is a primitive of zero, i.e., $H'(x) = (F(x) - G(x))' = 0$ for all $a < x < b$. Hence, $H(x) - H(y) = H'(c)(x - y) = 0$ for $a < x < y < b$, i.e., H is a constant. \square

For example, all the primitives of $f(x) = x^3$ are $F(x) = x^4/4 + C$ with C a real constant. Sometimes, F is called the antiderivative or the indefinite integral of f . We shall use only the term *primitive* and, symbolically, we write

$$(6.1) \quad F(x) = \int f(x) dx$$

to mean — no more, no less — $F'(x) = f(x)$ on the interval under consideration. The reason for the unusual notation (6.1) is due to the connection between the primitive and the integral. This is explained in §4.4. With this notation,

$$\int f'(x) dx = f(x)$$

is a tautology. As an additional mnemonic device, we sometimes write $d[f(x)] = f'(x)dx$. With this notation, \int and d “cancel”: $\int d[f(x)] = f(x)$.

Based on the derivative formulas in Chapter 3, we can list the primitives known to us at this point. These identities are valid on any open interval of definition. These formulas, like any formula involving primitives, can be checked by differentiation.

$$\int x^a dx = \frac{x^{a+1}}{a+1}, \quad a \neq -1,$$

$$\begin{aligned}
\int \frac{1}{x} dx &= \log |x|, \\
\int a^x dx &= \frac{1}{\log a} a^x, \quad a > 0, \\
\int \cos x dx &= \sin x, \\
\int \sin x dx &= -\cos x, \\
\int \sec^2 x dx &= \tan x, \\
\int \csc^2 x dx &= -\cot x, \\
\int \frac{1}{\sqrt{1-x^2}} dx &= \arcsin x,
\end{aligned}$$

and

$$\int \frac{1}{1+x^2} dx = \arctan x.$$

From the linearity of derivatives, we obtain the following.

Theorem. *If f and g have primitives on (a, b) and k is a real, so do $f + g$ and kf , and*

$$\int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx,$$

and

$$\int kf(x) dx = k \int f(x) dx. \quad \square$$

From the product rule, we obtain the following analog for primitives of summation by parts (§1.7).

Theorem (Integration By Parts). *If f and g are differentiable on (a, b) and $f'g$ has a primitive on (a, b) , then, so does fg' , and*

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx.$$

To see this, let F be a primitive for $f'g$. Then, $(fg - F)' = f'g + fg' - F' = fg'$, so, $fg - F$ is a primitive for fg' . \square We caution the reader that integration is taken up in §4.3. Here, the term *integration by parts* is just a name.

From the chain rule, we obtain the following.

Theorem (Substitution). If g is differentiable on (a, b) with $g[(a, b)] \subset (c, d)$ and $\int f(x) dx = F(x)$ on (c, d) , then,

$$\int f[g(x)]g'(x) dx = F[g(x)], \quad a < x < b. \quad \square$$

We work out some examples. It is convenient to allow the undefined symbol dx to enter into the expression for f and to write, e.g., $\int dx$ instead of $\int 1 dx$ and $\int dx/x$ instead of $\int (1/x)dx$.

Substitution is often written $\int f[g(x)]g'(x) dx = \int f(u)du$, $u = g(x)$. For example,

$$\begin{aligned} \int \cot x dx &= \int \frac{\cos x}{\sin x} dx \\ &= \int \frac{d \sin x}{\sin x} = \int \frac{du}{u} = \log |u| = \log |\sin x|. \end{aligned}$$

Similarly,

$$\begin{aligned} \int \frac{2x+1}{1+x^2} dx &= \int \frac{2x}{1+x^2} dx + \int \frac{1}{1+x^2} dx \\ &= \int \frac{d(1+x^2)}{1+x^2} + \arctan x \\ &= \log(1+x^2) + \arctan x. \end{aligned}$$

Particularly useful special cases of substitution are

$$\int f(x) dx = F(x) \quad \text{implying} \quad \int f(x+a) dx = F(x+a),$$

and

$$\int f(x) dx = F(x) \quad \text{implying} \quad \int f(ax) dx = \frac{1}{a} F(ax), \quad a \neq 0.$$

We call these the *translation* and *dilation* properties. Thus, for example,

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0.$$

Integration by parts is often written $\int u dv = uv - \int v du$. If we take $u = \log x$, $dv = dx$, then, $du = u' dx = dx/x$, $v = x$, so,

$$\int \log x dx = x \log x - \int x(dx/x) = x \log x - \int dx = x \log x - x.$$

Similarly,

$$\begin{aligned} \int x \log x dx &= \frac{x^2}{2} \log x - \int \frac{x^2}{2} (dx/x) \\ &= \frac{x^2}{2} \log x - \int \frac{x}{2} dx \end{aligned}$$

$$= \frac{x^2}{2} \log x - \frac{x^2}{4}.$$

By a trigonometric formula (§3.5),

$$\begin{aligned} \int \cos^2 x \, dx &= \int \frac{1 + \cos(2x)}{2} \, dx = \frac{x}{2} + \frac{1}{2} \int \cos(2x) \, dx \\ &= \frac{x}{2} + \frac{\sin(2x)}{4} = \frac{2x + \sin(2x)}{4}. \end{aligned}$$

Since $4x(1-x) = 4x - 4x^2 = 1 - (2x-1)^2$,

$$\int \frac{dx}{\sqrt{x(1-x)}} = \int \frac{2dx}{\sqrt{4x-4x^2}} = \int \frac{d(2x-1)}{\sqrt{1-(2x-1)^2}} = \arcsin(2x-1)$$

by translation and dilation. Of course, we already know (§3.5) that another primitive is $2 \arcsin \sqrt{x}$, so, the two primitives must differ by a constant (Exercise 9). The reduction $4x - 4x^2 = 1 - (2x-1)^2$ is the usual technique of *completing the square*.

To compute $\int \sqrt{1-x^2} \, dx$, let $x = \sin \theta$, $dx = \cos \theta \, d\theta$. Then,

$$\begin{aligned} \int \sqrt{1-x^2} \, dx &= \int \sqrt{1-\sin^2 \theta} \cos \theta \, d\theta \\ &= \int \cos^2 \theta \, d\theta \\ &= \frac{1}{4}[2\theta + \sin(2\theta)] \\ &= \frac{1}{2}(\theta + \sin \theta \cos \theta) \\ &= \frac{1}{2} \left(\arcsin x + x\sqrt{1-x^2} \right). \end{aligned}$$

Alternatively, let $u = \sqrt{1-x^2}$, $dv = dx$. Then, $du = -x dx / \sqrt{1-x^2}$, $v = x$. So,

$$\begin{aligned} \int \sqrt{1-x^2} \, dx &= x\sqrt{1-x^2} - \int x(-x dx / \sqrt{1-x^2}) \\ &= x\sqrt{1-x^2} + \int \frac{x^2 dx}{\sqrt{1-x^2}} \\ &= x\sqrt{1-x^2} + \int \frac{(x^2-1)dx}{\sqrt{1-x^2}} + \int \frac{dx}{\sqrt{1-x^2}} \\ &= x\sqrt{1-x^2} - \int \sqrt{1-x^2} \, dx + \arcsin x. \end{aligned}$$

Moving the second term on the right to the left side, we obtain the same result.

To compute $\int \frac{dx}{1-x^2}$, write

$$\frac{1}{1-x^2} = \frac{1}{2} \left(\frac{1}{1+x} + \frac{1}{1-x} \right)$$

to get

$$\int \frac{dx}{1-x^2} = \frac{1}{2} [\log(1+x) - \log(1-x)] = \frac{1}{2} \log \left(\frac{1+x}{1-x} \right).$$

If f is given as a power series, the primitive is easily found as another power series.

Theorem. *If $R > 0$ is the radius of convergence of*

$$f(x) = a_0 + a_1x + a_2x^2 + \dots,$$

then, R is the radius of convergence of $\sum a_n x^{n+1}/(n+1)$, and

$$(6.2) \quad \int f(x) dx = a_0x + \frac{a_1x^2}{2} + \frac{a_2x^3}{3} + \dots$$

on $(-R, R)$.

To see this, one first checks that the radius of convergence of the series in (6.2) is also R using $n^{1/n} \rightarrow 1$, as in the previous section. Now, differentiate the series in (6.2) obtaining $f(x)$. Hence, the series in (6.2) is a primitive. \square

For example, using the geometric series with $-x$ replacing x ,

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots, \quad |x| < 1.$$

Hence, by the theorem

$$(6.3) \quad \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots, \quad |x| < 1.$$

Indeed, both sides are primitives of $1/(1+x)$, and both sides equal zero at $x = 0$. Similarly, using the geometric series with $-x^2$ replacing x ,

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \dots, \quad |x| < 1.$$

Hence, by the theorem,

$$(6.4) \quad \arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots, \quad |x| < 1.$$

This follows since both sides are primitives of $1/(1+x^2)$, and both sides vanish at $x = 0$. To obtain the sum of the series (7.5) in §1.7, we seek to insert $x = 1$ in (6.4). We cannot do this directly since (6.4) is valid only for $|x| < 1$. Instead,

we let $s_n(x)$ denote the n th partial sum of the series in (6.4). Since this series is alternating with decreasing terms (§1.7) when $0 < x < 1$,

$$s_{2n}(x) \leq \arctan x \leq s_{2n-1}(x), \quad n \geq 1.$$

In this last inequality, the number of terms in the partial sums is finite. Letting $x \nearrow 1$, we obtain

$$s_{2n}(1) \leq \arctan 1 \leq s_{2n-1}(1), \quad n \geq 1.$$

Now, letting $n \nearrow \infty$ and recalling $\arctan 1 = \pi/4$, we arrive at the sum of the *Leibnitz series*

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots,$$

first discussed in §1.7. In particular, we conclude that

$$\frac{8}{3} < \pi < 4.$$

In §5.2, we obtain the sum of the Leibnitz series by a procedure that will be useful in many other situations.

The Leibnitz series is “barely convergent” and is not useful for computing π . To compute π , the traditional route is to insert $x = 1/5$ and $x = 1/239$ in (6.4) and to use Machin’s 1706 formula

$$\frac{\pi}{4} = 4 \arctan \frac{1}{5} - \arctan \frac{1}{239}.$$

Exercises 3.6

1. Compute $\int e^x \cos x \, dx$.
2. Compute $\int e^{\arcsin x} \, dx$.
3. Compute $\int \frac{x+1}{\sqrt{1-x^2}} \, dx$.
4. Compute $\int \frac{\arctan x}{1+x^2} \, dx$.
5. Compute $\int x^2 (\log x)^2 \, dx$.
6. Compute $\int \sqrt{1-e^{-2x}} \, dx$.
7. Let $F(x) = x^2 \sin(1/x)$, $x \neq 0$, and let $F(0) = 0$. Show that the derivative $F'(x) = f(x)$ exists for all x , but f is not continuous at $x = 0$.
8. If f is of bounded variation and $F' = f$ on (a, b) , then, f is continuous (Exercise 18 of §2.3).

9. Show directly (i.e., without derivatives) that

$$(6.5) \quad 2 \arcsin \sqrt{x} = \arcsin(2x - 1) + \pi/2.$$

10. Show that

$$\arcsin x = x + \frac{1}{2} \cdot \frac{x^3}{3} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{x^5}{5} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{x^7}{7} + \dots$$

for $|x| < 1$.

11. If f is a polynomial, then,

$$\int e^x f(-x) dx = e^x (f(-x) + f'(-x) + f''(-x) + f'''(-x) + \dots).$$

12. Show that

$$\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}.$$

Use this to derive Machin's formula.

13. Use Machin's formula and (6.4) to obtain $\pi = 3.14\dots$ to within an error of 10^{-2} .

14. Simplify $\arcsin(\sin 100)$. (The answer is not 100.)

15. Compute $\int \frac{-4x}{1-x^2} dx$.

16. Compute $\int \frac{4\sqrt{2} - 4x}{x^2 - \sqrt{2}x + 1} dx$.

17. Show that

$$\log 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

following the procedure discussed above for the Leibnitz series.

Integration

§4.1 The Cantor Set

The subject of this chapter is the measurement of the areas of subsets of the plane \mathbb{R}^2 . The areas of elementary geometric figures, such as squares, rectangles, and triangles, are already known to us. By *known to us* we mean that, e.g., by defining the area of a rectangle to be the product of the lengths of its sides, we obtain quantities that agree with our intuition. Since every right-angle triangle is half a rectangle, the areas of right-angle triangles are also known to us. Similarly, we can obtain the area of a general triangle.

How does one approach the problem of measuring the area of an unfamiliar figure or subset of \mathbb{R}^2 , say a subset that cannot be broken up into triangles? For example, how does one measure the area of the unit disk

$$D = \{(x, y) : x^2 + y^2 < 1\}?$$

One solution is to arbitrarily define the area of D to equal whatever one feels is right. The Egyptian book of Ahmes (~ 1900 B.C.) states that the area of D is $(16/9)^2$. In the Indian Śulbastras (written down ~ 500 B.C.), the area of D is taken to equal $(26/15)^2$. Albrecht Dürer (1471–1528) of Nuremburg solved a related problem which amounted to taking the area of D to equal $25/8$.

Which of these answers should we accept as the area of D ? If we treat these answers as *estimates* of the area of D , then, in our minds, we must have the presumption that such a quantity — the area of D — has a meaningful existence. In that case, we have no way of judging the merit of an estimate except by the quality of the reasoning leading to it.

Realizing this, by reasoning that remains perfectly valid today, Archimedes (~ 250 B.C.) carefully established,

$$\frac{223}{71} < \text{area}(D) < \frac{22}{7}.$$

In §4.4, we show that $\text{area}(D) = \pi$, where π , “Archimedes’ constant”, is the real number defined in §3.5.

At the basis of the Greek mathematicians’ computations of area was the *method of exhaustion*. This asserted that the area of a set $A \subset \mathbb{R}^2$ could be computed as

the limit of areas of a sequence of inscribed sets (A_n) that filled out more and more of A as $n \nearrow \infty$ (Figure 4.1). Nevertheless the Greeks were apparently uncomfortable with the concept of infinity and never used this method as stated. Instead, for example, in dealing with D , Archimedes used inscribed and circumscribed polygons with 96 sides to obtain the above result. He never explicitly passed to the limit. It turns out, however, that the method of exhaustion is so important to integration that in §4.5 we give a careful derivation of it.




FIGURE 4.1. The method of exhaustion.

Now, the unit disk is not a totally unfamiliar set to the reader. But, if we are presented with some genuinely unfamiliar subset C , the situation changes, and we may no longer have any clear conception of the area of C . If we are unable to come up with a procedure leading us to the area, then, we may be forced to reexamine our intuitive notion of area. In particular, we may be led to the conclusion that the “true area” of C may have no meaning. Let us describe such a subset.

Recall (§2.1) that an *open rectangle* is a subset of the form $(a, b) \times (c, d)$ where a or c may equal $-\infty$ and b or d may equal $+\infty$. A *compact rectangle* is a subset of the form $[a, b] \times [c, d]$.

Let C_0 denote the compact unit square $[0, 1] \times [0, 1]$. Divide C_0 into nine equal subsquares and take out from C_0 all but the four compact corner subsquares. Let C_1 be the remainder, i.e., the union of the four remaining compact subsquares. Repeat this process with each of the four subsquares. Divide each subsquare into nine equal compact sub-subsquares and take out, in each subsquare, all but the four compact corner sub-subsquares. Call the union of the remaining sixteen sub-subsquares C_2 . Continuing in this manner yields a sequence $C_0 \supset C_1 \supset C_2 \supset \dots$. The *Cantor set* is the common part (Figure 4.2) of all these sets, i.e., their intersection

$$C = \bigcap_{n=1}^{\infty} C_n.$$

At first glance it is not clear that C is not empty. But $(0, 0) \in C$! Moreover, the sixteen corners of the set C_1 are in C . Similarly, any corner of any subsquare, at any level, lies in C . But the set of such points is countable (§1.7), and it turns out that there is much more: There are as many points in C as there are in the unit square C_0 . In particular C is uncountable.




FIGURE 4.2. The Cantor set.

To see this, recall the concept of ternary expansions (§1.6). Let $a \in [0, 1]$. We say that

$$a = .a_1a_2 \dots$$

is the ternary expansion of a if the naturals a_n are ternary digits 0, 1, 2, and

$$a = \sum_{n=1}^{\infty} a_n 3^{-n}.$$

Now, let $(a, b) \in C_0$, and let

$$a = .a_1a_2 \dots$$

and

$$b = .b_1b_2 \dots$$

be ternary expansions of a and b . If $a_1 \neq 1$ and $b_1 \neq 1$, then, (a, b) is in C_1 . Similarly, in addition, if $a_2 \neq 1$ and $b_2 \neq 1$, $(a, b) \in C_2$. Continuing in this manner, we see that, if $a_n \neq 1$ and $b_n \neq 1$ for all $n \geq 1$, $(a, b) \in C$. Conversely, $(a, b) \in C$ implies that there are ternary expansions of a and b as stated. Thus, $(a, b) \in C$ iff a and b have ternary expansions in which the digits are equal to 0 or 2.

Now, although some reals may have more than one ternary expansion, a real a cannot have more than one ternary expansion $.a_1a_2 \dots$ where $a_n \neq 1$ for all $n \geq 1$ because any two ternary expansions yielding the same real must have their n th digits differing by 1 for some $n \geq 1$ (Exercise 2 of §1.6). Thus, the mapping

$$(a, b) = (.a_1a_2a_3 \dots, .b_1b_2b_3 \dots) \mapsto \left(\sum_{n=1}^{\infty} a'_n 2^{-n}, \sum_{n=1}^{\infty} b'_n 2^{-n} \right),$$

where $a'_n = a_n/2$, $b'_n = b_n/2$, $n \geq 1$, is well defined. Since any real in $[0, 1]$ has a binary expansion, this mapping is a surjection of the Cantor set C onto the unit square C_0 . Since C_0 is uncountable (Exercise 4 of §1.7), we conclude that C is uncountable (§1.7).

The difficulty of measuring the size of the Cantor set underscores the difficulty in arriving at a consistent notion of area. Above, we saw that the Cantor set is uncountable. In this sense, the Cantor set is “big”. On the other hand, note that

the areas of the subsquares removed from C_0 to obtain C_1 sum to $5/3^2$. Similarly, the areas of the sub-subsquares removed from C_1 to obtain C_2 sum to $20/9^2$. Similarly, at the next stage, we remove squares with areas summing to $80/27^2$. Thus, the sum of the areas of all the removed squares is

$$\frac{5}{9} + \frac{20}{9^2} + \frac{80}{27^2} + \cdots = \frac{5}{9} \left(1 + \frac{4}{9} + \left(\frac{4}{9}\right)^2 + \cdots \right) = 1.$$

Since C is the complement of all these squares in C_0 and C_0 has area 1, the area of C is $1 - 1 = 0$. Thus, in the sense of area, the Cantor set is “small”.

This argument is perfectly reasonable, except for one aspect. We are assuming that areas can be added and subtracted in the usual manner, even when there are *infinitely* many sets involved. In §4.2, we show that, with an appropriate definition of area, this argument can be modified to become correct, and the area of C is in fact zero.


Another indication of the smallness of C is the fact that C has no interior. To explain this, given any set $A \subset \mathbb{R}^2$, let us say that A *has interior* if we can fit some open rectangle Q within A , i.e., $Q \subset A$. If we cannot fit an open rectangle, no matter how small, within A , then, we say that A *has no interior*. For example, the unit disk has interior but a line segment has no interior. The Cantor set C has no interior, because there is a point in every open rectangle whose coordinate ternary expansions contain at least one digit 1. Alternatively, if C contained a rectangle Q , then, the area of C would be at least as much as the area of Q , which is positive. But we saw above that the area of C equals zero.

Since this reasoning applies to any set, we see that if $A \subset \mathbb{R}^2$ has interior, then, the area of A is positive. The surprising fact is that the converse of this statement is false. There are sets $A \subset \mathbb{R}^2$ that have positive area but have no interior. Such a set is described in Exercise 2.

These issues are discussed to point out the existence of unavoidable phenomena involving area where things do not behave as simply as triangles. In the first three decades of this century, these issues were finally settled. The solution to *the problem of area*, analyzed extensively by Archimedes more than two thousand years ago, can now be explained in a few pages. Why did it take so long for the solution to be discovered? It should not be too surprising that one missing ingredient was the completeness property of the set of real numbers, the importance of which was not fully realized until the nineteenth century.

Exercises 4.1

1. Let $C_0 = [0, 1] \times [0, 1]$ denote the unit square, and let C'_1 be obtained by throwing out from C_0 the middle subrectangle $(1/3, 2/3) \times [0, 1]$ of width $1/3$ and height 1. Then, C'_1 consists of two compact subrectangles. Let C'_2 be obtained from C'_1 by throwing out, in each of the subrectangles, the middle sub-subrectangles $(1/9, 2/9) \times [0, 1]$ and $(7/9, 8/9) \times [0, 1]$, each of width $1/3^2$ and height 1. Then, C'_2 consists of four compact sub-subrectangles. Similarly C'_3 consists of eight

FIGURE 4.3. A and B are rectangles, but C is not.

compact sub-sub-subrectangles, obtained by throwing out from C'_2 the middle sub-sub-subrectangles of width $1/3^3$ and height 1. Continuing in this manner, we have $C'_1 \supset C'_2 \supset C'_3 \supset \dots$. Let $C' = \bigcap_{n=1}^{\infty} C'_n$. Show that $\text{area}(C') = 0$ and C' has no interior.

2. Fix a real $0 < \alpha < 1$ (e.g., $\alpha = .7$) and let $C_0 = [0, 1] \times [0, 1]$ be the unit square. Let C_1^α be obtained from C_0 by throwing out the middle subrectangle of width $\alpha/3$ and height 1. Then, C_1^α consists of two subrectangles. Let C_2^α be obtained from C_1^α by throwing out, in each of the subrectangles, the *middle* sub-subrectangles of width $\alpha/3^2$ and height 1. Then, C_2^α consists of four sub-subrectangles. Similarly C_3^α consists of eight sub-sub-subrectangles, obtained by throwing out from C_2^α the middle sub-sub-subrectangles of width $\alpha/3^3$ and height 1. Continuing in this manner, we have $C_1^\alpha \supset C_2^\alpha \supset C_3^\alpha \supset \dots$. Let $C^\alpha = \bigcap_{n=1}^{\infty} C_n^\alpha$. Show that $\text{area}(C^\alpha) > 0$, but C^α has no interior.

3. For $A \subset \mathbb{R}^2$ let

$$A + A = \{(x + x', y + y') : (x, y) \in A, (x', y') \in A\}$$

be the *set of sums*. Show that $C + C = [0, 2] \times [0, 2]$ (Exercise 1.6.6).

§4.2 Area

Recall that $(a, b) = \{x : a < x < b\}$ and $[a, b] = \{x : a \leq x \leq b\}$. An open rectangle is a subset (Figure 4.3) of the form $(a, b) \times (c, d)$, where a or c may equal $-\infty$ and b or d may equal ∞ . In particular, \mathbb{R}^2 is an open rectangle. We say that an open rectangle is *bounded* if a, b, c , and d are reals. A compact rectangle is a subset of the form $[a, b] \times [c, d]$ where a, b, c , and d are reals. In particular, the vertical line segment $\{a\} \times [c, d]$ is a compact rectangle. Similarly, so is a horizontal line segment. A single point is a compact rectangle. If $Q = (a, b) \times (c, d)$ is a bounded, nonempty, open rectangle, the compact rectangle $\overline{Q} = [a, b] \times [c, d]$ is its *compactification*. Note that an open rectangle may be empty, for example $(a, a) \times (c, d)$ is empty.

Let A be a subset of \mathbb{R}^2 . A *cover* of A is a sequence of sets (A_n) , such that A is contained in their union,

$$A \subset \bigcup_{n=1}^{\infty} A_n.$$

In a given cover, the sets (A_n) may overlap, i.e., intersect. If for some N , $A_n = \emptyset$ for $n > N$, we say that (A_1, \dots, A_N) is a *finite cover* (Figure 4.4).




FIGURE 4.4. A finite cover.

A *paving* of A is a cover (Q_n) , where the sets Q_n , $n \geq 1$, are *open rectangles*. A *finite paving* is a finite cover that is also a paving (Figure 4.5). Every subset $A \subset \mathbb{R}^2$ has at least one (not very interesting) paving $Q_1 = \mathbb{R}^2$, $Q_2 = \emptyset$, $Q_3 = \emptyset$,




FIGURE 4.5. A finite paving.

For any nonempty, open rectangle $Q = (a, b) \times (c, d)$ or compact rectangle $Q = [a, b] \times [c, d]$, let

$$\|Q\| = (b - a) \cdot (d - c).$$

Then, $\|Q\|$ is a positive real or 0 or ∞ . We also take $\|\emptyset\| = 0$.

Let A be a subset of \mathbb{R}^2 . The *area* of A is defined by

$$\text{area}(A) = \inf \left\{ \sum_{n=1}^{\infty} \|Q_n\| : \text{all pavings } (Q_n) \text{ of } A \right\}.$$

This definition of area is at the basis of all that follows. It is necessarily complicated because it applies to all subsets A of \mathbb{R}^2 . As an immediate consequence of the

definition, $\text{area}(\emptyset) = 0$. Similarly, the area of a finite vertical line segment A is zero since A can be covered by a thin open rectangle of arbitrarily small area.

In words, the definition says that to find the area of a set A , we cover A by a sequence Q_1, Q_2, \dots of open rectangles, measure the sum of their areas, and take this sum as an estimate for the area of A . Of course, we expect that this sum will be an overestimate of the area of A for two reasons. The paving may cover a superset of A , and we are not taking into account any overlaps when computing the sum. Then, we define the area of A to be the inf of these sums.

Of course, carrying out this procedure explicitly, even for simple sets A , is completely impractical. Because of this, we almost never use the definition directly to compute areas. Instead, as is typical in mathematics, we derive the elementary properties of area from the definition, and we use them to compute areas.

We emphasize that according to the above definitions, a rotated rectangle A is not a rectangle. Hence, $\|A\|$ is not defined unless the sides of the rectangle A are parallel to the axes. Nevertheless, we will see, below, that area is rotation-invariant. Hence, $\text{area}(A)$ turns out as expected.

Whether or not we can compute the area of a given set, the above definition applies consistently to every subset A . In particular, this is so whether A is a rectangle, a triangle, a smooth graph, or the Cantor set C . Now, let us derive the properties of area that follow immediately from the definition.

Since every open rectangle Q is a paving of itself, $\text{area}(Q) \leq \|Q\|$. Below, we obtain $\text{area}(Q) = \|Q\|$ for a rectangle Q . Until we establish this, we refer to $\|Q\|$ as the *naive area* of Q . Note that, although area is defined for every subset, naive area is defined only for (nonrotated) rectangles.

If (a, b) is a point in \mathbb{R}^2 and $A \subset \mathbb{R}^2$, the set

$$A + (a, b) = \{(x + a, y + b) : (x, y) \in A\}$$

is the *translate* of A by (a, b) . Then, $[A + (a, b)] + (c, d) = A + (a + c, b + d)$ and, for any rectangle Q , $\|Q + (a, b)\| = \|Q\|$. From this follows the *translation invariance of area*,

$$\text{area}[A + (a, b)] = \text{area}(A), \quad A \subset \mathbb{R}^2.$$


To see this, let (Q_n) be a paving of A . Then, $(Q_n + (a, b))$ is a paving of $A + (a, b)$, so

$$\text{area}[A + (a, b)] \leq \sum_{n=1}^{\infty} \|Q_n + (a, b)\| = \sum_{n=1}^{\infty} \|Q_n\|.$$

Since $\text{area}(A)$ is the inf of the sums on the right, $\text{area}[A + (a, b)] \leq \text{area}(A)$. Now, in this last inequality, replace, in order, (a, b) by $(-a, -b)$ and A by $A + (a, b)$. We obtain $\text{area}(A) \leq \text{area}[A + (a, b)]$. Hence, $\text{area}(A) = \text{area}([A + (a, b)] + (-a, -b))$, establishing translation invariance (Figure 4.6).

If $k > 0$ is real and $A \subset \mathbb{R}^2$, the set

$$kA = \{(kx, ky) : (x, y) \in A\}$$

FIGURE 4.6. $\text{area}(A) = \text{area}[A + (a, b)]$.


is the *dilate* of A by k . Then, $k(cA) = (kc)A$ for k and c positive, and $\|kQ\| = k^2\|Q\|$ for every rectangle Q . From this follows the *dilation invariance of area*,

$$\text{area}(kA) = k^2 \cdot \text{area}(A), \quad A \subset \mathbb{R}^2.$$

To see this, let (Q_n) be a paving of A . Then, (kQ_n) is a paving of kA so

$$\text{area}(kA) \leq \sum_{n=1}^{\infty} \|kQ_n\| = k^2 \left(\sum_{n=1}^{\infty} \|Q_n\| \right).$$

Since $\text{area}(A)$ is the inf of the sums on the right, we obtain $\text{area}(kA) \leq k^2 \cdot \text{area}(A)$. Now, in this last inequality, replace, in order, k by $1/k$ and A by kA . We obtain $k^2 \cdot \text{area}(A) \leq \text{area}(kA)$. Hence, $\text{area}(kA) = k^2 \cdot \text{area}(A)$, establishing dilation invariance.

FIGURE 4.7. $\text{area}(kA) = k^2 \cdot \text{area}(A)$.

Instead of dilation from the origin, we can dilate from any point in \mathbb{R}^2 . In particular, by elementary geometry, certain subsets such as rectangles, triangles, and parallelograms have well defined centers. Given a subset A with a center (a, b) , its *centered dilation* kA is the set (Figure 4.8) obtained by translating (a, b) to $(0, 0)$, dilating as above by the factor $k > 0$, then, translating $(0, 0)$ back




FIGURE 4.8. Centered dilation.

to (a, b) . Then, $\text{area}(kA) = k^2 \cdot \text{area}(A)$ for centered dilations as well. For example, if $k < 1$, kA is A shrunk towards its center.

The diagonal line segment $A = \{(x, y) : 0 \leq x \leq 1, y = x\}$ has zero area (Figure 4.9). To see this, choose $n \geq 1$ and let $Q_k = \{(x, y) : (k-1)/n < x < k/n, (k-1)/n < y < k/n\}$, $k = 1, \dots, n$. Then, (Q_1, \dots, Q_n) is almost a paving of A , except for the corner points of Q_k in A , $k = 1, \dots, n$, which are not covered. To remedy this, for $c > 1$, let cQ_k denote the centered dilations of Q_k , $k = 1, \dots, n$. Then, (cQ_1, \dots, cQ_n) is a paving of A . Hence, $\text{area}(A) \leq \|cQ_1\| + \dots + \|cQ_n\| = n \cdot c^2 \cdot \frac{1}{n^2} = c^2/n$. Since $n \geq 1$ may be arbitrarily large, we conclude that $\text{area}(A) = 0$. Similarly, the area of any finite line segment is zero.




FIGURE 4.9. Area of a diagonal line segment.

As with dilation, setting $-A = \{(-x, -y) : (x, y) \in A\}$, we have *reflection invariance of area*,

$$\text{area}(-A) = \text{area}(A), \quad A \subset \mathbb{R}^2,$$

and *monotonicity*,

$$\text{area}(A) \leq \text{area}(B), \quad A \subset B \subset \mathbb{R}^2.$$

Another property is *subadditivity*. For any cover (A_n) of a given set A ,

$$(2.1) \quad \text{area}(A) \leq \sum_{n=1}^{\infty} \text{area}(A_n).$$

Here, the sets A_n , $n \geq 1$, need not be rectangles. For future reference, we call the sum on the right side of (2.1) the *area of the cover* (A_n) .

In particular, since (A, B) is a cover of $A \cup B$,

$$\text{area}(A \cup B) \leq \text{area}(A) + \text{area}(B).$$

Similarly, (A_1, A_2, \dots, A_n) is a cover of $A_1 \cup A_2 \cup \dots \cup A_n$, so,

$$\text{area}(A_1 \cup A_2 \cup \dots \cup A_n) \leq \text{area}(A_1) + \text{area}(A_2) + \dots + \text{area}(A_n).$$

To obtain subadditivity, note that, if the right side of (2.1) is ∞ , there is nothing to show, since in that case (2.1) is true. Hence, we may safely assume $\text{area}(A_n) < \infty$ for all $n \geq 1$. Let $\epsilon > 0$, and, for each $k \geq 1$, choose a paving $(Q_{k,n})$ of A_k satisfying

$$(2.2) \quad \sum_{n=1}^{\infty} \|Q_{k,n}\| < \text{area}(A_k) + \epsilon 2^{-k}.$$

This is possible since $\text{area}(A_k)$ is the inf of sums of the form $\sum_{n=1}^{\infty} \|Q_n\|$. Then, the double sequence $(Q_{k,n})$ is a cover by open rectangles, hence, a paving of A . Summing (2.2) over $k \geq 1$, we obtain

$$\begin{aligned} \text{area}(A) &\leq \sum_{k=1}^{\infty} \left(\sum_{n=1}^{\infty} \|Q_{k,n}\| \right) \\ &\leq \sum_{k=1}^{\infty} (\text{area}(A_k) + \epsilon 2^{-k}) = \left(\sum_{k=1}^{\infty} \text{area}(A_k) \right) + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, subadditivity follows.

From subadditivity and monotonicity, we see that the area of any (infinite) line is zero and $\text{area}(Q) = \text{area}(\overline{Q})$ for any bounded open rectangle Q . Moreover, $\text{area}(Q) \leq \|Q\|$ for any compact rectangle Q .

Theorem. *The area of a (compact or open) rectangle equals the product of the lengths of its sides.*

We see, below, that this holds for rotated rectangles as well. Before we derive this Theorem, we make some elementary observations concerning the naive area (Figure 4.10).

Let A be a compact rectangle. If $A = Q_1 \cup \dots \cup Q_N$ with $A = [a, b] \times [c, d]$, $Q_n = [a_{n-1}, a_n] \times [c, d]$, $n = 1, \dots, N$, and $a = a_0 < a_1 < \dots < a_N = b$, then, by definition of $\|\cdot\|$,

$$(2.3) \quad \|A\| = \|Q_1\| + \dots + \|Q_N\|.$$

Let us call such a cover (Q_1, \dots, Q_N) of A horizontal type I. Similarly, we can define vertical type I covers. (Note that, since the rectangles are compact, type I covers are not pavings.) If $A = Q_1 \cup \dots \cup Q_N$ is a vertical type I cover and each Q_i can be broken into a horizontal type I cover, $Q_i = Q_{i1} \cup \dots \cup Q_{iN_i}$,

$i = 1, \dots, N$, we call $(Q_{ij} : 1 \leq i \leq N, 1 \leq j \leq N_i)$ a type II cover of A . Clearly, for a type II cover, $\|A\|$ equals the sum of $\|Q_{ij}\|$ over all $1 \leq j \leq N_i$, $1 \leq i \leq N$. If we rewrite the rectangles in a type II cover as (Q_1, \dots, Q_N) , we see that (2.3) holds also for any type II cover of A . If $A = Q_1 \cup \dots \cup Q_N$, where $Q_n, n = 1, \dots, N$, are possibly overlapping compact rectangles, then, we say that (Q_1, \dots, Q_N) is a type III cover. Given a type III cover of A , by extending all the horizontal edges of all the rectangles in the cover till they intersect the edges of A , we obtain a type II cover. Moreover, because the overlaps are counted more than once and for type II covers we have (2.3), we conclude that

$$(2.4) \quad \|A\| \leq \|Q_1\| + \dots + \|Q_N\|$$

for type III covers, where $A = Q_1 \cup \dots \cup Q_N$.




FIGURE 4.10. Type I, type II, type III covers.

Now, we can derive the theorem. Let A be a compact rectangle. We already know (Exercise 2) that $\text{area}(A) \leq \|A\|$, so, we need only derive $\|A\| \leq \text{area}(A)$. To establish this we need to show that

$$(2.5) \quad \|A\| \leq \sum_{n=1}^{\infty} \|Q_n\|$$

for every paving (Q_n) of A . Let (Q_n) be a paving of A . We want to apply (2.4) to (Q_n) . To this end, we need to make four reductions that turn the paving (Q_n) into a type III cover.

If any one of the rectangles Q_n is unbounded, then, the right side of (2.5) is infinite, so, (2.5) is true. Therefore, we may assume that all the rectangles Q_n , $n \geq 1$, are bounded. This is the first reduction. Now, since the rectangles Q_n are open, there is (§2.1) a natural N , such that (Q_1, \dots, Q_N) is a finite paving of A . This is the second reduction. Then, $(\overline{Q}_1, \dots, \overline{Q}_N)$ is a finite cover of A by compact rectangles. This is the third reduction. Now, cutting off the excess leads us to the type III cover $(\overline{Q}_1 \cap A, \dots, \overline{Q}_N \cap A)$. This is the fourth reduction. Hence, by (2.4), we conclude that

$$\|A\| \leq \sum_{n=1}^N \|\overline{Q}_n \cap A\| \leq \sum_{n=1}^N \|\overline{Q}_n\| = \sum_{n=1}^N \|Q_n\| \leq \sum_{n=1}^{\infty} \|Q_n\|,$$

establishing (2.5). Taking the inf over all pavings (Q_n) of A in (2.5), we obtain $\|A\| \leq \text{area}(A)$, hence, $\text{area}(A) = \|A\|$ for A compact. Since $\text{area}(A) =$

area (\overline{A}) and $\|A\| = \|\overline{A}\|$ for A bounded, open, we also obtain the result for bounded, open rectangles A . When A is an unbounded open rectangle, we know that $\|A\| = \infty$. On the other hand, since there are bounded subrectangles Q in A for which $\text{area}(Q) = \|Q\|$ is arbitrarily large, we obtain $\text{area}(A) = \infty$ by monotonicity. Hence, $\text{area}(A) = \|A\|$ for this case also. \square

Now, we can compute the area of a triangle A with a horizontal base of length b and height h by constructing a cover of A consisting of thin horizontal strips (Figure 4.11). Let S^- and S^+ denote the nonhorizontal sides, where we take S^- to the left of S^+ and let $\|A\|$ denote the naive area of A , i.e., $\|A\| = hb/2$. By reflection invariance, we may assume A lies above its base.

For each $n \geq 1$ on S^- , mark $n + 1$ equally spaced points (x_i^-, y_i) , $i = 0, 1, \dots, n$. Similarly, on S^+ , mark (x_i^+, y_i) , $i = 0, 1, \dots, n$. Here, we take $y_0 = 0$, $y_n = h$, $y_0 < y_1 < \dots < y_n$ and $y_i - y_{i-1} = h/n$, $i = 1, \dots, n$. Then, (x_0^-, y_0) and (x_0^+, y_0) are the endpoints of the base, and $x_0^+ - x_0^-$ is the length of the base. Two cases can occur. Either S^- has positive slope, in which case $x_0^- < x_1^- < \dots < x_n^-$, or S^- has negative slope, in which case $x_0^- > x_1^- > \dots > x_n^-$. To take care of both cases simultaneously, let $x_i^{-*} = \min(x_{i-1}^-, x_i^-)$ for each $i = 1, \dots, n$. Similarly, let $x_i^{+*} = \max(x_{i-1}^+, x_i^+)$, $i = 1, \dots, n$.




FIGURE 4.11. Cover of a triangle.

Now, let Q_i denote the compact rectangle $[x_i^{-*}, x_i^{+*}] \times [y_{i-1}, y_i]$, $i = 1, \dots, n$. Then, we obtain a cover (Q_1, \dots, Q_n) of A (the Q_i 's cover by definition of $x_i^{\pm*}$). Since the height of each Q_i is h/n and the width is $x_i^{+*} - x_i^{-*}$, by subadditivity, we obtain

$$\text{area}(A) \leq h \cdot \frac{1}{n} \sum_{i=1}^n (x_i^{+*} - x_i^{-*})$$

which says $\text{area}(A)$ is less or equal to h times the average of the widths of the rectangles Q_i , $i = 1, \dots, n$. But the average of the widths of the rectangles Q_i , $i = 1, \dots, n$, differs from $b/2$, at most, by b/n . Now, let $n \nearrow \infty$. Then, the sequence of averages converges to $b/2$, and we obtain $\text{area}(A) \leq \|A\|$.

To obtain the reverse inequality, draw two other triangles B , C with horizontal bases, such that $A \cup B \cup C$ is a rectangle and A , B , and C intersect only along their edges. Then, by simple arithmetic, the sum of the naive areas of A , B , and C equals the naive area of $A \cup B \cup C$, so, by subadditivity of area,

$$\begin{aligned} \|A\| + \|B\| + \|C\| &= \|A \cup B \cup C\| \\ &= \text{area}(A \cup B \cup C) \end{aligned}$$

$$\begin{aligned} &\leq \text{area}(A) + \text{area}(B) + \text{area}(C) \\ &\leq \text{area}(A) + \|B\| + \|C\|. \end{aligned}$$

Cancelling $\|B\|$, $\|C\|$, we obtain the reverse inequality $\|A\| \leq \text{area}(A)$.

Theorem. *The area of a triangle equals half the product of the length of its base and its height.* \square

We have derived this theorem assuming that the base of the triangle is horizontal. The general case follows from rotation invariance, which we do below. Now, let P be a parallelogram with horizontal base and let $\|P\|$ denote its naive area, i.e., the product of the length of its base and its height. Then, we leave it as an exercise to show that $\text{area}(P) = \|P\|$.

Theorem. *The area of a parallelogram equals the product of the length of its base and height.* \square

By the next theorem, the areas of rectangles, triangles and parallelograms are given by the last three theorems, even when rotated. Recall that rotations were defined in §3.5.

Theorem (Rotation Invariance). *Let $A \subset \mathbb{R}^2$. If A is rotated into A' , then, $\text{area}(A) = \text{area}(A')$.*

To see this, first, assume that Q is a bounded rectangle. Hence, Q' is a rotated rectangle. Now, decompose Q' into the union of a parallelogram P and two triangles S, T , all with horizontal bases and intersecting along their edges (Figure 4.12). Then, by simple arithmetic, the sum of the naive areas of S, T , and P equals the naive area of Q , so by subadditivity

$$\begin{aligned} \text{area}(Q') &\leq \text{area}(S) + \text{area}(P) + \text{area}(T) \\ &= \|S\| + \|P\| + \|T\| \\ &= \|Q\| = \text{area}(Q). \end{aligned}$$

We have just shown that $\text{area}(Q') \leq \text{area}(Q)$ for any bounded rectangle Q . If Q is an unbounded rectangle, this last inequality is clearly true. Hence, $\text{area}(Q') \leq \text{area}(Q)$ for every rectangle Q .

Now, let A be any subset, and let (Q_n) be a paving of A . Suppose that the rotation sending A to A' sends Q_n to Q'_n , $n \geq 1$. Then, $A' \subset \bigcup_{n=1}^{\infty} Q'_n$. So, by subadditivity

$$\text{area}(A') \leq \sum_{n=1}^{\infty} \text{area}(Q'_n) \leq \sum_{n=1}^{\infty} \text{area}(Q_n).$$

Here, we used the inequality derived in the previous paragraph. Taking the inf over all pavings of A , we obtain $\text{area}(A') \leq \text{area}(A)$. If we apply this last inequality




FIGURE 4.12. Rotation invariance of area.

to A' instead of A and to the inverse rotation, we obtain the reverse inequality $\text{area}(A) \leq \text{area}(A')$. We conclude that $\text{area}(A') = \text{area}(A)$. \square

Area also has good invariance properties under other types of dilation. For example, let $k > 0$ and set $H(x, y) = (kx, y)$, $V(x, y) = (x, ky)$. Then, the mappings $H : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $V : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ dilate area in the sense $\text{area}[H(A)] = k \cdot \text{area}(A) = \text{area}[V(A)]$. To see this, first, check that this is so on rectangles. Then, use the definition of area to arrive at the general case.

Our last item is additivity. In general, we do not expect that $\text{area}(A \cup B) = \text{area}(A) + \text{area}(B)$ because A and B may overlap, i.e., intersect. If A and B are disjoint, one expects to have additivity. Here, we establish additivity only for the case when A and B are well separated. Exercises 12 and 13 of §4.5 discuss a broader case.

If $A \subset \mathbb{R}^2$ and $B \subset \mathbb{R}^2$, set

$$d(A, B) = \inf \sqrt{(a - c)^2 + (b - d)^2},$$

where the \inf is over all points $(a, b) \in A$ and points $(c, d) \in B$. We say A and B are *well separated* if $d(A, B)$ is positive (Figure 4.13). For example, although $\{(2, 0)\}$ and the unit disk are well separated, $\mathbb{Q} \times \mathbb{Q}$ and $\{(\sqrt{2}, 0)\}$ are disjoint but not well separated. Note that, since $\inf \emptyset = \infty$, A empty implies $d(A, B) = \infty$. Hence, the empty set is well separated from any subset of \mathbb{R}^2 .




FIGURE 4.13. Well Separated sets.

If the lengths of the sides of a rectangle Q are a and b , by the *diameter* of Q , we mean the length of the diagonal $\sqrt{a^2 + b^2}$.

Theorem (Additivity). *If A and B are well separated, then,*

$$\text{area}(A \cup B) = \text{area}(A) + \text{area}(B).$$

By subadditivity, $\text{area}(A \cup B) \leq \text{area}(A) + \text{area}(B)$, so, we need show only that

$$(2.6) \quad \text{area}(A \cup B) \geq \text{area}(A) + \text{area}(B).$$

If $\text{area}(A \cup B) = \infty$ (2.6) is true, so, assume $\text{area}(A \cup B) < \infty$. In this case, to compute the area of $A \cup B$, we need consider only pavings involving bounded open rectangles, since the sum of the areas of open rectangles with at least one unbounded, open rectangle, is ∞ . Let $\epsilon = d(A, B) > 0$. If (Q_n) is a paving of $A \cup B$ with bounded, open rectangles $Q_n, n \geq 1$, decompose the compactification \overline{Q}_n of each bounded open rectangle Q_n into a type II cover (see above) where the compact subrectangles are all of diameter less than ϵ . Since $\|Q_n\|$, then, equals the sum of the areas of its subrectangles, by replacing each Q_n by its subrectangles, we obtain a cover (\overline{Q}'_n) of $A \cup B$ by compact rectangles, where each \overline{Q}'_n has diameter less than ϵ and

$$\sum_{n=1}^{\infty} \|\overline{Q}'_n\| = \sum_{n=1}^{\infty} \|Q_n\|.$$


Thus, for each n , \overline{Q}'_n intersects A or B or neither but not both. For each n , if \overline{Q}'_n intersects A , denote it \overline{Q}^A_n . If it intersects B , denote it \overline{Q}^B_n . Because no \overline{Q}'_n intersects both A and B , (\overline{Q}^A_n) must be a cover of A and (\overline{Q}^B_n) must be a cover of B . Hence, by subadditivity,

$$\begin{aligned} \text{area}(A) + \text{area}(B) &\leq \sum_{n=1}^{\infty} \text{area}(\overline{Q}^A_n) + \sum_{n=1}^{\infty} \text{area}(\overline{Q}^B_n) \\ &\leq \sum_{n=1}^{\infty} \text{area}(\overline{Q}'_n) \\ &= \sum_{n=1}^{\infty} \|Q_n\|. \end{aligned}$$

Taking the inf over all pavings (Q_n) of $A \cup B$, we obtain the result. \square

As an application, let A denote the unit square $[0, 1] \times [0, 1]$ and B the triangle obtained by joining the three points $(1, 0)$, $(1, 1)$ and $(2, 1)$. We already know that $\text{area}(A) = 1$ and $\text{area}(B) = 1/2$, and we want to conclude that $\text{area}(A \cup B) = 1 + 1/2 = 3/2$ (Figure 4.14). But A and B are not well separated, so, we do not have additivity directly. Instead, we dilate A by a factor $0 < \alpha < 1$ towards its center. Then, the shrunk set αA and B are well separated. Moreover, $\alpha A \subset A$, so,

$$\begin{aligned} \text{area}(A \cup B) &\geq \text{area}((\alpha A) \cup B) \\ &= \text{area}(\alpha A) + \text{area}(B) \\ &= \alpha^2 \cdot \text{area}(A) + \text{area}(B) \\ &= \alpha^2 + 1/2. \end{aligned}$$

FIGURE 4.14. Area of $A \cup B$.

Since α can be arbitrarily close to 1, we conclude that $\text{area}(A \cup B) \geq 3/2$. Since subadditivity yields $\text{area}(A \cup B) \leq \text{area}(A) + \text{area}(B) = 1 + 1/2 = 3/2$, we obtain the result we seek, $\text{area}(A \cup B) = 3/2$.

Additivity holds by induction for several sets. If A_1, A_2, \dots, A_n are pairwise well separated subsets of \mathbb{R}^2 , then,

$$(2.7) \quad \text{area}(A_1 \cup \dots \cup A_n) = \text{area}(A_1) + \dots + \text{area}(A_n).$$

To see this, (2.7) is trivially true for $n = 1$, so, assume (2.7) is true for a particular $n \geq 1$, and let A_1, \dots, A_{n+1} be pairwise well separated. Let $\epsilon_j = d(A_j, A_{n+1}) > 0$, $j = 1, \dots, n$. Since

$$d(A_1 \cup \dots \cup A_n, A_{n+1}) = \min(\epsilon_1, \dots, \epsilon_n) > 0,$$

A_{n+1} and $A_1 \cup \dots \cup A_n$ are well separated. Hence, by the inductive hypothesis,

$$\begin{aligned} \text{area}(A_1 \cup \dots \cup A_{n+1}) &= \text{area}(A_1 \cup \dots \cup A_n) + \text{area}(A_{n+1}) \\ &= \text{area}(A_1) + \dots + \text{area}(A_n) + \text{area}(A_{n+1}). \end{aligned}$$

By induction, this establishes (2.7) for all $n \geq 1$.

More generally, if (A_n) is a sequence of pairwise well separated sets, then,

$$(2.8) \quad \text{area}\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \text{area}(A_n).$$

To see this, subadditivity yields

$$\text{area}\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \text{area}(A_n).$$

For the reverse inequality, apply (2.7) and monotonicity to the first N sets, yielding

$$\text{area}\left(\bigcup_{n=1}^{\infty} A_n\right) \geq \text{area}\left(\bigcup_{n=1}^N A_n\right) = \sum_{n=1}^N \text{area}(A_n).$$

Now let $N \nearrow \infty$, obtaining

$$\text{area}\left(\bigcup_{n=1}^{\infty} A_n\right) \geq \sum_{n=1}^{\infty} \text{area}(A_n).$$

This establishes (2.8).

As an application of (2.8), we can now compute the area of the Cantor set C . The Cantor set is constructed by removing, at successive stages, smaller and smaller

open subsquares of $C_0 = [0, 1] \times [0, 1]$. Denote these subsquares Q_1, Q_2, \dots (at what stage they are removed is not important). Then, for each n , C and Q_n are disjoint, so, for $0 < \alpha < 1$, C and the centered dilations αQ_n are well separated. Moreover for each m, n the centered dilations αQ_n and αQ_m are well separated. But the union of C with all the squares αQ_n , $n \geq 1$, lies in the unit square C_0 . Hence, by (2.8),

$$\text{area}(C) + \sum_{n=1}^{\infty} \text{area}(\alpha Q_n) = \text{area}\left(C \cup \left(\bigcup_{n=1}^{\infty} \alpha Q_n\right)\right) \leq \text{area}(C_0) = 1.$$

In the previous section we obtained, $\sum_{n=1}^{\infty} \text{area}(Q_n) = 1$. By dilation invariance, this implies $\text{area}(C) + \alpha^2 \leq 1$. Letting $\alpha \nearrow 1$, we obtain $\text{area}(C) + 1 \leq 1$ or $\text{area}(C) = 0$.

Theorem. *The area of the Cantor set is zero.* \square

Exercises 4.2

1. Establish reflection invariance and monotonicity of area.
2. Show that the area of a bounded line segment is zero, the area of any line is zero, and $\text{area}(Q) = \text{area}(\overline{Q})$ for any bounded open rectangle Q . Show also that $\text{area}(Q) \leq \|Q\|$ for any compact rectangle Q .
3. Let P be a parallelogram with a horizontal base, and let $\|P\|$ denote the product of the length of its base and its height. Then, $\text{area}(P) = \|P\|$.
4. Compute the area of a trapezoid.
5. If A and B are rectangles, then, $\text{area}(A \cup B) = \text{area}(A) + \text{area}(B) - \text{area}(A \cap B)$.
6. For $k \in \mathbb{R}$, define $H : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $V : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $H(x, y) = (kx, y)$ and $V(x, y) = (x, ky)$. Then, $\text{area}[V(A)] = |k| \cdot \text{area}(A) = \text{area}[H(A)]$ for every $A \subset \mathbb{R}^2$.
7. ★ A mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is *linear* if it is of the form $L(x, y) = (ax + by, cx + dy)$ with $a, b, c, d \in \mathbb{R}$. Show that a linear mapping sends lines to (possibly collapsed) lines and parallelograms to (possibly collapsed) parallelograms. Show that a linear mapping L is invertible (i.e., a bijection §1.1) iff the real $\det(L) = ad - bc$ is not zero. In this case, show that the inverse K of L is linear and compute $\det(K)$.
8. ★ Let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be an invertible linear mapping. Show that

$$(2.9) \quad \text{area}[L(A)] = |\det(L)| \cdot \text{area}(A), \quad A \subset \mathbb{R}^2.$$

Thus, L is area-preserving iff $\det(L) = \pm 1$. Such an L is called *affine*, and this result is *affine-invariance* of area. (Do this for rectangles first.)




FIGURE 4.15. Affine invariance of area.

9. ★ Let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a noninvertible linear mapping. Show that $L(\mathbb{R}^2)$ is contained in a line and (2.9) holds.
10. Show that $\{(\sqrt{2}, 0)\}$ and the unit disk are well separated, but $\{(\sqrt{2}, 0)\}$ and $\mathbb{Q} \times \mathbb{Q}$ are not.
11. Let D be the unit disk, and let $D^+ = \{(x, y) : x^2 + y^2 < 1 \text{ and } y > 0\}$. Show that $\text{area}(D) = 2 \cdot \text{area}(D^+)$.
12. Compute the area of the sets C' and C^α described in Exercises 1 and 2 of §4.1, using the properties of area.
13. Let $P_k = (\cos(2\pi k/n), \sin(2\pi k/n))$, $k = 0, 1, \dots, n$. Then, P_0, P_1, \dots, P_n are evenly spaced points on the unit circle with $P_n = P_0$. Let D_n denote the n -sided polygon obtained by joining the points P_k . Compute $\text{area}(D_n)$.
14. Let $A \subset \mathbb{R}^2$. A *triangular paving* of A is a cover (T_n) of A where each T_n , $n \geq 1$, is a triangle (oriented arbitrarily). With $\text{area}(A)$ as defined previously, show that

$$\text{area}(A) = \inf \left\{ \sum_{n=1}^{\infty} \|T_n\| : \text{all triangular pavings } (T_n) \text{ of } A \right\}.$$

Here, $\|T\|$ denotes the naive area of the triangle T , i.e., half the product of the length of the base times the height.

15. ★ Let $A \subset \mathbb{R}^2$. If $\text{area}(A) > 0$ and $0 < \alpha < 1$, there is *some* open rectangle Q , such that $\text{area}(Q \cap A) > \alpha \cdot \text{area}(Q)$. (Argue by contradiction, and use the definition of area.)

§4.3 The Integral

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function defined on an open interval (a, b) , where, as usual, a may equal $-\infty$ or b may equal ∞ . We say f is *bounded* if $|f(x)| \leq M$, $a < x < b$, for some real M . If f is *nonnegative*, i.e., if $f(x) \geq 0$, $a < x < b$, the *subgraph* of f over (a, b) is the set (Figure 4.16)

$$G = \{(x, y) : a < x < b, 0 < y < f(x)\} \subset \mathbb{R}^2.$$




FIGURE 4.16. Subgraphs of nonnegative functions.

Note that the inequalities in this definition are strict.

For nonnegative f , we define the *integral* $\int_a^b f(x) dx$ of f from a to b to be the area of its subgraph G ,

$$\int_a^b f(x) dx = \text{area}(G).$$

Then, the integral is either 0, a positive real, or ∞ . The reason for the unusual notation is explained below.

Thus, according to our definition, *every nonnegative function has an integral* and integrals of nonnegative functions are areas — nothing more, nothing less — of certain subsets of \mathbb{R}^2 .

Since the empty set has zero area, we always have $\int_a^a f(x) dx = 0$. For each $k \geq 0$, the subgraph of $f(x) = k$, $a < x < b$, over (a, b) is an open rectangle, so,

$$\int_a^b k dx = k(b - a).$$

Since the area is monotone, so is the integral: If $0 \leq f \leq g$ on (a, b) ,

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

In particular, $0 \leq f \leq M$ on (a, b) implies $0 \leq \int_a^b f(x) dx \leq M(b - a)$.

A nonnegative function f is *integrable* over (a, b) if $\int_a^b f(x) dx < \infty$. For example, we have just seen that every bounded nonnegative f is integrable over a bounded interval (a, b) . Now, we discuss the integral of a *signed* function, i.e., a function that takes on positive and negative values.


Given a function $f : (a, b) \rightarrow \mathbb{R}$, we set

$$f^+(x) = \max[f(x), 0],$$

and

$$f^-(x) = \max[-f(x), 0].$$

These are (Figure 4.17) the *positive part* and the *negative part* of f , respectively. Note that $f^+ - f^- = f$ and $f^+ + f^- = |f|$.

FIGURE 4.17. Positive and negative parts of $\sin x$.

We say a signed function f is *integrable over* (a, b) if

$$\int_a^b |f(x)| dx < \infty.$$

In this case, $\int_a^b f^\pm(x) dx \leq \int_a^b |f(x)| dx$ are both finite. For integrable f , we define the *integral* $\int_a^b f(x) dx$ of f from a to b by

$$\int_a^b f(x) dx = \int_a^b f^+(x) dx - \int_a^b f^-(x) dx.$$

From this follows

$$\int_a^b [-f(x)] dx = - \int_a^b f(x) dx$$

for every integrable function f , since $g = -f$ implies $g^+ = f^-$ and $g^- = f^+$.

We warn the reader that, although $\int_a^b f(x) dx = \int_a^b f^+(x) dx - \int_a^b f^-(x) dx$ is a definition, the identity $\int_a^b |f(x)| dx = \int_a^b f^+(x) dx + \int_a^b f^-(x) dx$ is not true for general integrable f . However, it is true when f is continuous. This property (linearity) is discussed in the next section.

Also, from the above discussion, we see that *every bounded (signed) function is integrable over a bounded interval*. For example, $\sin x$ and $\sin x/x$ are integrable over $(0, \pi)$. In fact, both functions are integrable over $(0, b)$ for any finite b and, hence, $\int_0^b \sin x dx$ and $\int_0^b (\sin x/x) dx$ are defined.

It is reasonable to expect that $\sin x$ is not integrable over $(0, \infty)$. Indeed the subgraph of $|\sin x|$ consists of a union of sets $G_n, n \geq 1$, where each G_n denotes the subgraph over $((n-1)\pi, n\pi)$. By translation invariance, the sets $G_n, n \geq 1$, have the same positive area and the sets G_1, G_3, G_5, \dots , are well separated. Hence, we obtain

$$\int_0^\infty (\sin x)^+ dx = \text{area} \left(\bigcup_{n=1}^\infty G_{2n-1} \right) = \sum_{n=1}^\infty \text{area}(G_{2n-1}) = \infty.$$

By considering, instead, G_2, G_4, G_6, \dots , we obtain $\int_0^\infty (\sin x)^- dx = \infty$. Thus,

$$\int_0^\infty (\sin x)^+ dx - \int_0^\infty (\sin x)^- dx = \infty - \infty.$$

Hence, $\int_0^\infty \sin x dx$ cannot be defined as a difference of two areas.

It turns out that $\sin x/x$ is also not integrable over $(0, \infty)$. To see this, let G_n denote the subgraph of $|\sin x/x|$ over $((n-1)\pi, n\pi)$, $n \geq 1$ (Figure 4.18). In each G_n , we can insert a rectangle Q_n of area $\sqrt{2}/(4n-1)$ so that the rectangles are well separated. By additivity, then, we obtain

$$\int_0^\infty \frac{|\sin x|}{x} dx \geq \text{area} \left(\bigcup_{n=1}^\infty Q_n \right) = \sum_{n=1}^\infty \text{area}(Q_n) = \sum_{n=1}^\infty \frac{\sqrt{2}}{4n-1} = \infty,$$

by comparison with the harmonic series. Thus, $\sin x/x$ is not integrable over $(0, \infty)$. More explicitly, this reasoning also shows that

$$\int_0^\infty \left(\frac{\sin x}{x} \right)^+ dx \geq \text{area}(Q_1) + \text{area}(Q_3) + \text{area}(Q_5) + \dots = \infty$$

and

$$\int_0^\infty \left(\frac{\sin x}{x} \right)^- dx \geq \text{area}(Q_2) + \text{area}(Q_4) + \text{area}(Q_6) + \dots = \infty.$$

Thus,

$$\int_0^\infty \left(\frac{\sin x}{x} \right)^+ dx - \int_0^\infty \left(\frac{\sin x}{x} \right)^- dx = \infty - \infty,$$

hence, $\int_0^\infty \sin x/x dx$ also cannot be defined as a difference of two areas.




FIGURE 4.18. The graphs of $\sin x/x$ and $|\sin x|/x$.

To summarize, *the integral of an integrable function is the area of the subgraph of its positive part minus the area of the subgraph of its negative part.* Every property of $\int_a^b f(x) dx$ ultimately depends on a corresponding property of area.

Frequently, one checks integrability of a given f by first applying one or more of the properties below to the nonnegative function $|f|$. For example, consider the function $g(x) = 1/x^2$ for $x > 1$, and, for each $n \geq 1$, let G_n denote the compact rectangle $[n, n+1] \times [0, 1/n^2]$. Then, (G_n) is a cover of the subgraph of g over $(1, \infty)$ (Figure 4.19). Hence,

$$\int_1^\infty \frac{1}{x^2} dx \leq \sum_{n=1}^\infty \frac{1}{n^2},$$

which is finite (§1.6). Thus, g is integrable over $(1, \infty)$. Since the signed function $f(x) = \cos x/x^2$ satisfies $|f(x)| \leq g(x)$ for $x > 1$, by monotonicity, we conclude that

$$(3.1) \quad \int_1^\infty \left| \frac{\cos x}{x^2} \right| dx < \infty.$$

Hence, $\cos x/x^2$ is integrable over $(1, \infty)$.




FIGURE 4.19. A cover of the subgraph of $1/x^2$ over $(1, \infty)$.


Of course, functions may be unbounded and integrable. For example, the function $f(x) = 1/\sqrt{x}$ is integrable over $(0, 1)$. To see this, let G_n denote the compact rectangle $[1/(n+1)^2, 1/n^2] \times [0, n+1]$. Then, (G_n) is a cover of the subgraph of f over $(0, 1)$ (Figure 4.20). Hence,

$$\int_0^1 \frac{1}{\sqrt{x}} dx \leq \sum_{n=1}^\infty (n+1) \left(\frac{1}{n^2} - \frac{1}{(n+1)^2} \right) = \sum_{n=1}^\infty \frac{2n+1}{n^2(n+1)} \leq 2 \sum_{n=1}^\infty \frac{1}{n^2},$$

which is finite. Thus, f is integrable over $(0, 1)$.

Theorem (Monotonicity). *Suppose that f and g are both nonnegative or both integrable on (a, b) . If $f \leq g$ on (a, b) , then,*

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

FIGURE 4.20. A cover of the subgraph of $1/\sqrt{x}$ over $(0, 1)$.

If $0 \leq f \leq g$, we already know this. For the integrable case, note that $f \leq g$ implies $f^+ = \max(f, 0) \leq \max(g, 0) = g^+$ and $g^- = \max(-g, 0) \leq \max(-f, 0) = f^-$ on (a, b) . Hence,

$$\int_a^b f^+(x) dx \leq \int_a^b g^+(x) dx,$$

and

$$\int_a^b f^-(x) dx \geq \int_a^b g^-(x) dx.$$

Subtracting the second inequality from the first, the result follows. \square

Since $\pm f \leq |f|$, the theorem implies

$$\pm \int_a^b f(x) dx = \int_a^b \pm f(x) dx \leq \int_a^b |f(x)| dx$$

which yields

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

for every integrable f .

Theorem (Translation and Dilation Invariance). *Let f be nonnegative or integrable on (a, b) . Choose $c \in \mathbb{R}$ and $k > 0$. Then,*

$$\int_a^b f(x+c) dx = \int_{a+c}^{b+c} f(x) dx,$$

$$\int_a^b kf(x) dx = k \int_a^b f(x) dx,$$

and

$$\int_a^b f(kx) dx = \frac{1}{k} \int_{ka}^{kb} f(x) dx.$$

If f is nonnegative, let G denote the subgraph of $f(x + c)$ over (a, b) (Figure 4.21). Then, the translate $G + (c, 0)$ equals

$$\{(x, y) : a + c < x < b + c, 0 < y < f(x)\},$$

which is the subgraph of $f(x)$ over the interval $(a + c, b + c)$. By translation invariance of area, we obtain translation invariance of the integral in the nonnegative case. If f is integrable, by the nonnegative case,

$$\int_a^b f^+(x + c) dx = \int_{a+c}^{b+c} f^+(x) dx,$$

and

$$\int_a^b f^-(x + c) dx = \int_{a+c}^{b+c} f^-(x) dx.$$

Now, if $g(x) = f(x + c)$, then, $g^+(x) = f^+(x + c)$ and $g^-(x) = f^-(x + c)$. So, subtracting the last equation from the previous one, we obtain translation invariance in the integrable case.

For the second equation and f nonnegative, recall that from the previous section the dilation mapping $V(x, y) = (x, ky)$, and let G denote the subgraph of f over (a, b) . Then, $V(G) = \{(x, y) : a < x < b, 0 < y < kf(x)\}$. Hence, $\text{area}(V(G)) = \int_a^b kf(x) dx$. Now, dilation invariance of the area (Exercise 6 of §4.2) yields $\int_a^b kf(x) dx = k \int_a^b f(x) dx$ for f nonnegative. For integrable f , the result follows by applying, as above, the nonnegative case to f^+ and f^- .

For the third equation, let $H(x, y) = (kx, y)$, and let G denote the subgraph of $f(kx)$ over (a, b) . Then, $H(G) = \{(x, y) : ka < x < kb, 0 < y < f(x)\}$. The third equation now follows, as before, by dilation invariance. For integrable f , the result follows by applying the nonnegative case to f^\pm . \square




FIGURE 4.21. Translation and dilation invariance of integrals.

By similar reasoning, one can also derive (Figure 4.22)

$$\int_{-b}^{-a} f(-x) dx = \int_a^b f(x) dx,$$

valid for f nonnegative or integrable over (a, b) .

The next property is additivity.




FIGURE 4.22. Reflection invariance of integrals.

Theorem (Additivity). Suppose that f is nonnegative or integrable over (a, b) , and choose $a < c < b$. Then,

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

To see this, first, assume that f is nonnegative. Since the vertical line $x = c$ has zero area, subadditivity yields

$$\int_a^b f(x) dx \leq \int_a^c f(x) dx + \int_c^b f(x) dx.$$

So, we need only show that

$$(3.2) \quad \int_a^b f(x) dx \geq \int_a^c f(x) dx + \int_c^b f(x) dx.$$

If f is not integrable, (3.2) is immediate since the left side is infinite, so, assume f is nonnegative and integrable. Now, choose any strictly increasing sequence $a < c_1 < c_2 < \dots$ converging to c . Then, for $n \geq 1$, the subgraph of f over (a, c_n) and the subgraph of f over (c, b) are well separated (Figure 4.23). So, by monotonicity and well separated additivity,

$$(3.3) \quad \int_a^b f(x) dx \geq \int_a^{c_n} f(x) dx + \int_c^b f(x) dx.$$

We wish to send $n \nearrow \infty$ in (3.3). To this end, for each $n \geq 1$, let G_n denote the subgraph of f over (c_n, c_{n+1}) , and let L_n denote the vertical line segment $\{(x, y) : x = c_n, 0 < y < f(x)\}$. Since G_2, G_4, G_6, \dots , are pairwise well separated,

$$\text{area}(G_2) + \text{area}(G_4) + \text{area}(G_6) + \dots \leq \int_a^c f(x) dx < \infty.$$

Since G_1, G_3, G_5, \dots , are pairwise well separated,

$$\text{area}(G_1) + \text{area}(G_3) + \text{area}(G_5) + \dots \leq \int_a^c f(x) dx < \infty.$$

Adding the last two inequalities yields the convergence of $\sum_{n=1}^{\infty} \text{area}(G_n)$. Hence, the tail (§1.6) goes to zero:

$$\lim_{n \nearrow \infty} \sum_{k=n}^{\infty} \text{area}(G_k) = 0.$$

Since the subgraph of f over (c_n, c) equals $G_n \cup L_{n+1} \cup G_{n+1} \cup L_{n+2} \cup G_{n+2} \cup L_{n+3} \cup \dots$, subadditivity implies

$$0 \leq \int_{c_n}^c f(x) dx \leq \sum_{k=n}^{\infty} \text{area}(G_k), \quad n \geq 1.$$

Hence, we obtain

$$(3.4) \quad \lim_{n \nearrow \infty} \int_{c_n}^c f(x) dx = 0.$$

Since by monotonicity and subadditivity, again,

$$\int_a^{c_n} f(x) dx \leq \int_a^c f(x) dx \leq \int_a^{c_n} f(x) dx + \int_{c_n}^c f(x) dx, \quad n \geq 1,$$

we conclude that

$$\lim_{n \nearrow \infty} \int_a^{c_n} f(x) dx = \int_a^c f(x) dx.$$

Now, sending $n \nearrow \infty$ in (3.3) yields (3.2). Hence, the result for f nonnegative. If f is integrable, apply the nonnegative case to f^+ and f^- . Then,

$$\int_a^b f^+(x) dx = \int_a^c f^+(x) dx + \int_c^b f^+(x) dx,$$

and

$$\int_a^b f^-(x) dx = \int_a^c f^-(x) dx + \int_c^b f^-(x) dx.$$

Subtracting the second equation from the first, we obtain the result in the integrable case. \square




FIGURE 4.23. Additivity of integrals.

The bulk of the derivation above involves establishing (3.4). If f is bounded, say by M , then, the integral in (3.4) is no more than $M(c - c_n)$, hence, trivially, goes to zero. The delicacy is necessary to handle unbounded situations.

The main point in the derivation is that, although the subgraph G of f over (a, c) and the subgraph G' of f over (c, b) are not well separated, we still have additivity, because we know something — the existence of the vertical edges — about the geometry of G and G' .

In the previous section, when we wanted to apply additivity to several sets (for example, when we computed the area of the Cantor set) that were not well separated, we dilated them by a factor $0 < \alpha < 1$ and, then, applied additivity to the shrunken sets.

Why don't we use the same trick here for G or G' ? The reason is that if the graph of f is sufficiently "jagged" (Figure 4.24), we do not have $\alpha G \subset G$, a necessary step in applying the shrinking trick of the previous section.




FIGURE 4.24. A "jagged" function.

By induction, additivity holds for a partition (§2.2) of (a, b) : If $a = x_0 < x_1 < \dots < x_{n+1} = b$ and f is nonnegative or integrable over (a, b) , then,

$$\int_a^b f(x) dx = \sum_{k=1}^{n+1} \int_{x_{k-1}}^{x_k} f(x) dx.$$

Since the right side does not involve the values of f at the points defining the partition, we conclude that *the integrals of two functions $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ are equal, whenever they differ only on finitely many points $a < x_1 < \dots < x_n < b$.*

Another application of additivity is to piecewise constant functions. A function $f : (a, b) \rightarrow \mathbb{R}$ is *piecewise constant* if there is a partition $a = x_0 < x_1 < \dots < x_{n+1} = b$, such that f , restricted to each open subinterval (x_{i-1}, x_i) , $i = 1, \dots, n + 1$, is constant. (Note that the values of a piecewise constant function at the partition points x_i , $1 \leq i \leq n$, are not restricted in any way.) In this case, additivity implies

$$\int_a^b f(x) dx = \sum_{i=1}^{n+1} c_i \Delta x_i,$$

where $\Delta x_i = x_i - x_{i-1}$, $i = 1, \dots, n + 1$. Since a continuous function can be closely approximated by a piecewise constant function (§2.3), the integral should

be thought of as a sort of sum with Δx_i “infinitely small,” hence, the notation dx replacing Δx_i and \int replacing \sum .

This view is supported by Exercise 3. Indeed, by defining integrals as areas of subgraphs, we capture the intuition that integrals are approximately sums of areas of rectangles in *any* paving, and not just finite vertical pavings as given by the “Riemann sums” of Exercise 3.

Also, since the integral is, by definition, a combination of certain areas and the notation $\int_a^b f(x) dx$ is just a mnemonic device, the variable inside the integral sign is a “dummy” variable, i.e., $\int_a^b f(x) dx = \int_a^b f(t) dt$. Nevertheless, the interpretation of the integral as a “continuous sum” is basic, useful, and important.

Let us go back to the integrals of $\sin x$ and $\sin x/x$ over $(0, \infty)$. Above, we saw that these functions were not integrable over $(0, \infty)$, and, so, $\int_0^\infty \sin x dx$ and $\int_0^\infty \sin x/x dx$ could not be defined as the difference of the areas of the positive and the negative parts. An alternate approach is to consider $F(b) = \int_0^b \sin x dx$ and to take the limit $F(\infty) = \lim_{b \rightarrow \infty} F(b)$. However, since the areas of the sets G_n , $n \geq 1$, are equal, by additivity, $F(n\pi) = \text{area}(G_1) - \text{area}(G_2) + \cdots \pm \text{area}(G_n)$ equals $\text{area}(G_1)$ or zero according to whether n is odd or even. Thus, the limit $F(\infty)$ does not exist and this approach fails for $\sin x$.

For $\sin x/x$, however, it is a different story. Let $F(b) = \int_0^b \sin x/x dx$, and let G_n denote the subgraph of $|\sin x|/x$ over $((n-1)\pi, n\pi)$ for each $n \geq 1$. Then, by additivity $F(n\pi) = \text{area}(G_1) - \text{area}(G_2) + \cdots \pm \text{area}(G_n)$. Hence,

$$\lim_{n \nearrow \infty} \int_0^{n\pi} \frac{\sin x}{x} dx = \text{area}(G_1) - \text{area}(G_2) + \text{area}(G_3) - \dots$$

But this last series has a finite sum since it is alternating with decreasing terms! Thus,

$$(3.5) \quad \int_0^\infty \frac{\sin x}{x} dx \neq \lim_{n \nearrow \infty} \int_0^{n\pi} \frac{\sin x}{x} dx$$

since the left side is not defined and the right side is a well defined, finite real. The limit $\lim_{n \nearrow \infty} F(n\pi)$ is computed in Exercise 12 of §5.4.

On the other hand, when f is nonnegative or integrable, its integral over an interval (a, b) can be obtained as a limit of integrals over subintervals (a_n, b_n) (Figure 4.25), and the behavior (3.5) does not occur.

Theorem (Continuity At The Endpoints). *If f is nonnegative or integrable on (a, b) and $a_n \rightarrow a+$, $b_n \rightarrow b-$, then,*

$$(3.6) \quad \int_a^b f(x) dx = \lim_{n \nearrow \infty} \int_{a_n}^{b_n} f(x) dx.$$

If f is integrable on (a, b) and $a_n \rightarrow a+$, $b_n \rightarrow b-$, then, in addition,

$$(3.7) \quad \lim_{n \nearrow \infty} \int_a^{a_n} f(x) dx = 0,$$

and

$$(3.8) \quad \lim_{n \nearrow \infty} \int_{b_n}^b f(x) dx = 0.$$




FIGURE 4.25. Continuity at the endpoints.

To see this, first assume that f is nonnegative and $b_n \nearrow b$, and fix $a < c < b$. Since area is monotone, the sequence $\int_c^{b_n} f(x) dx$, $n \geq 1$, is increasing and

$$\lim_{n \nearrow \infty} \int_c^{b_n} f(x) dx \leq \int_c^b f(x) dx.$$

For the reverse inequality, let G_n denote the subgraph of f over (b_n, b_{n+1}) , $n \geq 1$. By additivity,

$$\int_c^{b_n} f(x) dx = \int_c^{b_1} f(x) dx + \sum_{k=1}^{n-1} \text{area}(G_k),$$

so, taking the limit and using subadditivity,

$$\begin{aligned} \lim_{n \nearrow \infty} \int_c^{b_n} f(x) dx &= \int_c^{b_1} f(x) dx + \sum_{k=1}^{\infty} \text{area}(G_k) \\ &\geq \int_c^{b_1} f(x) dx + \int_{b_1}^b f(x) dx \geq \int_c^b f(x) dx. \end{aligned}$$

Hence,

$$(3.9) \quad \lim_{n \nearrow \infty} \int_c^{b_n} f(x) dx = \int_c^b f(x) dx.$$

In general, if $b_n \rightarrow b-$, then, $b_{n*} \nearrow b$ (§1.5), and $b_{n*} \leq b_n < b$. Hence,

$$\int_c^{b_{n*}} f(x) dx \leq \int_c^{b_n} f(x) dx \leq \int_c^b f(x) dx, \quad n \geq 1,$$

which implies (3.9), for general $b_n \rightarrow b-$. Since

$$\int_{a_n}^c f(x) dx = \int_{-c}^{-a_n} f(-x) dx,$$

applying what we just learned to $f(-x)$ yields

$$\lim_{n \nearrow \infty} \int_{a_n}^c f(x) dx = \int_a^c f(x) dx.$$

Hence,

$$\begin{aligned} \int_a^b f(x) dx &= \int_a^c f(x) dx + \int_c^b f(x) dx \\ &= \lim_{n \nearrow \infty} \int_{a_n}^c f(x) dx + \lim_{n \nearrow \infty} \int_c^{b_n} f(x) dx \\ &= \lim_{n \nearrow \infty} \int_{a_n}^{b_n} f(x) dx. \end{aligned}$$

For the integrable case, apply (3.6) to f^\pm to get (3.6) for f . Since

$$\int_a^{a_n} f(x) dx = \int_a^b f(x) dx - \int_{a_n}^b f(x) dx,$$

we get (3.7). Similarly, we get (3.8). \square

For example,

$$\int_0^1 x^r dx = \lim_{a \rightarrow 0+} \int_a^1 x^r dx,$$

and

$$\int_1^\infty x^r dx = \lim_{b \rightarrow \infty} \int_1^b x^r dx,$$

both for r real.

When f is integrable, the last theorem can be improved: We have continuity of the integral at every point in (a, b) .

Theorem (Continuity). *Suppose that f is integrable over (a, b) , and set*

$$F(t) = \int_a^t f(x) dx, \quad a < t < b.$$

Then, F is continuous on (a, b) .

To see this, fix $a < c < b$, and let $c_n \rightarrow c-$. Applying the previous theorem on (a, c) , we obtain $F(c_n) \rightarrow F(c)$. Hence, we obtain continuity of F from the left at every real in (a, b) .

Now, let $g(x) = f(-x)$, $-b < x < -a$, and

$$G(t) = \int_t^{-a} g(x) dx, \quad -b < t < -a.$$

Since, by additivity,

$$G(t) = \int_{-b}^{-a} g(x) dx - \int_{-b}^t g(x) dx, \quad -b < t < -a,$$

by the previous paragraph applied to g , the function G is continuous from the left at every point in $(-b, -a)$. Thus, the function

$$G(-t) = \int_{-t}^{-a} f(-x) dx = \int_a^t f(x) dx = F(t), \quad a < t < b,$$

is continuous from the right at every point in (a, b) . This establishes continuity of F on (a, b) . \square

Our last item is the integral test for positive series.

Integral Test. *Let $f : (0, \infty) \rightarrow (0, \infty)$ be decreasing. Then,*

$$(3.10) \quad \gamma = \lim_{n \nearrow \infty} \left[\sum_{k=1}^n f(k) - \int_1^{n+1} f(x) dx \right]$$

exists and $0 \leq \gamma \leq f(1)$. In particular, the integral $\int_1^\infty f(x) dx$ is finite iff the sum $\sum_{n=1}^\infty f(n)$ converges.




FIGURE 4.26. Integral test.

For each $n \geq 1$, let $B_n = (n, n+1) \times (0, f(n))$, $B'_n = (n, n+1) \times [f(n+1), f(n)]$, and let G_n denote the subgraph of f over $(n, n+1)$ (Figure 4.26). Since f is decreasing, $G_n \subset B_n \subset G_n \cup B'_n$, for all $n \geq 1$. Then, the quantity, whose limit is the right side of (3.10), equals

$$\sum_{k=1}^n [\text{area}(B_k) - \text{area}(G_k)],$$

which is clearly increasing with n (here, we used additivity). Hence, the limit $\gamma \geq 0$ exists. On the other hand, by subadditivity, we get

$$\text{area}(B_k) - \text{area}(G_k) \leq \text{area}(B'_k) = f(k) - f(k+1).$$

So,

$$\gamma = \sum_{n=1}^{\infty} [\text{area}(B_k) - \text{area}(G_k)] \leq \sum_{n=1}^{\infty} [f(k) - f(k+1)] = f(1).$$

Thus, $\gamma \leq f(1)$. If either $\int_1^\infty f(x) dx$ or $\sum_{n=1}^\infty f(n)$ is finite, (3.10) simplifies to

$$\gamma = \sum_{n=1}^\infty f(n) - \int_1^\infty f(x) dx,$$

which shows that the sum is finite iff the integral is finite. \square

Exercises 4.3

1. Show that $\int_0^\infty f(kx)x^{-1} dx = \int_0^\infty f(x)x^{-1} dx$ for $k > 0$ and $f(x)/x$ nonnegative or integrable over $(0, \infty)$.
2. Show that $\int_{-b}^{-a} f(-x) dx = \int_a^b f(x) dx$ for f nonnegative or integrable over (a, b) .
3. ★ Let $f : [a, b] \rightarrow \mathbf{R}$ be continuous. If $a = x_0 < x_1 < \cdots < x_{n+1} = b$ is a partition of $[a, b]$, a *Riemann sum* corresponding to this partition is the real (Figure 4.27)

$$\sum_{i=1}^{n+1} f(x_i^\#)(x_i - x_{i-1}),$$

where $x_i^\#$ is arbitrarily chosen in (x_{i-1}, x_i) , $i = 1, \dots, n+1$. Let $I = \int_a^b f(x) dx$. Show that, for every $\epsilon > 0$, there is a $\delta > 0$, such that

$$(3.11) \quad \left| I - \sum_{i=1}^{n+1} f(x_i^\#)(x_i - x_{i-1}) \right| \leq \epsilon$$

for any partition $a = x_0 < x_1 < \cdots < x_{n+1} = b$ of mesh less than δ and choice of points $x_1^\#, \dots, x_{n+1}^\#$. (Approximate f by a piecewise constant f_ϵ as in §2.3.)

4. Let $f : (0, 1) \rightarrow \mathbf{R}$ be given by

$$f(x) = \begin{cases} x & \text{if } x \text{ irrational,} \\ 0 & \text{if } x \text{ rational.} \end{cases}$$

Compute $\int_0^1 f(x) dx$.

5. ★ Let $f : (a, b) \rightarrow \mathbf{R}$ be nonnegative, and suppose that $g : (a, b) \rightarrow \mathbf{R}$ is nonnegative and piecewise constant. Use additivity to show that

$$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$$

(First, do this for g constant.)

6. Let $f : (0, \infty) \rightarrow \mathbf{R}$ be nonnegative and equal to a constant c_n on each subinterval $(n-1, n)$ for $n = 1, 2, \dots$. Then,

$$\int_0^\infty f(x) dx = \sum_{n=1}^\infty c_n.$$




FIGURE 4.27. Riemann sums.

Instead, if f is integrable, then, $\sum_{n=1}^{\infty} c_n$ is absolutely convergent and the equality holds.

7. ★ A function $f : (a, b) \rightarrow \mathbb{R}$ is *Riemann integrable over* (a, b) if there is a real I satisfying the following property: For all $\epsilon > 0$, there is a $\delta > 0$, such that (3.11) holds for any partition $a = x_0 < x_1 < \dots < x_{n+1} = b$ of mesh less than δ and choice of intermediate points $x_1^\#, \dots, x_n^\#$. Thus, Exercise 3 says every function continuous on a compact interval $[a, b]$ is Riemann integrable over (a, b) . Let $f(x) = 0$ for $x \in \mathbb{Q}$ and $f(x) = 1$ for $x \notin \mathbb{Q}$. Show that this f is *not* Riemann integrable over $(0, 1)$.

8. Let $g : (0, \infty) \rightarrow (0, \infty)$ be decreasing and bounded. Show that

$$\lim_{\delta \rightarrow 0+} \delta \sum_{n=1}^{\infty} g(n\delta) = \int_0^{\infty} g(x) dx.$$

(Apply the integral test to $f(x) = g(x\delta)$.)

9. Let $f : (-b, b) \rightarrow \mathbb{R}$ be nonnegative or integrable. If f is even, then, $\int_{-b}^b f(x) dx = 2 \int_0^b f(x) dx$. Now, let f be integrable. If f is odd, then, $\int_{-b}^b f(x) dx = 0$.

10. Show that $\int_{-\infty}^{\infty} e^{-a|x|} dx < \infty$ for $a > 0$.

11. ★ If $f : \mathbb{R} \rightarrow \mathbb{R}$ is superlinear (Exercise 20 of §2.3) and continuous, the *Laplace transform*

$$L(s) = \int_{-\infty}^{\infty} e^{sx} e^{-f(x)} dx$$

is finite for all $s \in \mathbb{R}$. (Write $\int_{-\infty}^{\infty} = \int_{-\infty}^a + \int_a^b + \int_b^{\infty}$ for appropriately chosen a and b .)

12. ★ A function $\delta : \mathbb{R} \rightarrow \mathbb{R}$ is a *Dirac delta function* if it is nonnegative and satisfies

$$(3.12) \quad \int_{-\infty}^{\infty} \delta(x) f(x) dx = f(0)$$

for *all* continuous nonnegative $f : \mathbb{R} \rightarrow \mathbb{R}$. Show that there is no such function. (Construct continuous f 's which take on the two values 0 or 1 on most or all of \mathbb{R} , and insert them into (3.12).)

§4.4 The Fundamental Theorem of Calculus

By constructing appropriate covers, Archimedes was able to compute areas and integrals in certain situations. For example, he knew that $\int_0^1 x^2 dx = 1/3$. On the other hand, Archimedes was also able to compute tangent lines to certain curves and surfaces. However, he apparently had no idea that these two processes were intimately related, through the fundamental theorem of calculus. It was the discovery of the fundamental theorem, in the seventeenth century, that turned the computation of areas from a mystery to a simple and straightforward reality.

In this section, all functions will be continuous. Since we will use f^+ and f^- repeatedly, it is important to note that (§2.3) *a function is continuous iff both its positive and negative parts are continuous*.

Let f be continuous on (a, b) , and let $[c, d]$ be a compact subinterval. Since (§2.3) continuous functions map compact intervals to compact intervals, f is bounded on $[c, d]$, hence, integrable over (c, d) .

Let f be continuous on (a, b) , fix $a < c < b$, and set

$$F_c(x) = \begin{cases} \int_c^x f(t) dt, & c \leq x < b, \\ -\int_x^c f(t) dt, & a < x \leq c. \end{cases}$$

By the previous paragraph, $F_c(x)$ is finite for all $a < x < b$. From the previous section, we know that F_c is continuous. Here, we show that F_c is differentiable and $F'_c(x) = f(x)$ on (a, b) (Figure 4.28). We will need the modulus of continuity μ_x (§2.3) of f at x . To begin, by additivity, $F_c(y) - F_c(x) = F_x(y) - F_x(x)$ for any two points x, y in (a, b) , whether they are to the right or the left of c .

Then, for $a < x < t < y < b$, $|f(t) - f(x)| \leq \mu_x(y - x)$. Thus, $f(t) \leq f(x) + \mu_x(y - x)$. Hence,

$$\begin{aligned} \frac{F_c(y) - F_c(x)}{y - x} &= \frac{F_x(y) - F_x(x)}{y - x} \\ &= \frac{1}{y - x} \int_x^y f(t) dt \\ &\leq \frac{1}{y - x} \int_x^y [f(x) + \mu_x(y - x)] dt = f(x) + \mu_x(y - x). \end{aligned}$$

Similarly, since $a < x < t < y < b$ implies $f(x) - \mu_x(y - x) \leq f(t)$,

$$\frac{F_c(y) - F_c(x)}{y - x} \geq f(x) - \mu_x(y - x).$$

Combining the last two inequalities, we obtain

$$\left| \frac{F_c(y) - F_c(x)}{y - x} - f(x) \right| \leq \mu_x(y - x)$$

for $a < x < y < b$. If $a < y < x < b$, repeating the same steps yields

$$\left| \frac{F_c(y) - F_c(x)}{y - x} - f(x) \right| \leq \mu_x(x - y).$$

Hence, if $a < x \neq y < b$,

$$\left| \frac{F_c(y) - F_c(x)}{y - x} - f(x) \right| \leq \mu_x(|y - x|),$$

which implies, by continuity of f at x ,

$$\lim_{y \rightarrow x} \frac{F_c(y) - F_c(x)}{y - x} = f(x).$$

Hence, $F'_c(x) = f(x)$. We have established the following result, first mentioned in §3.6.




FIGURE 4.28. The derivative at x of the integral of f is $f(x)$.

Theorem. Every continuous $f : (a, b) \rightarrow \mathbb{R}$ has a primitive on (a, b) . \square

When f is continuous and integrable on (a, b) , we can do better.

Theorem. Let $f : (a, b) \rightarrow \mathbb{R}$ be continuous and integrable. Then,

$$F(x) = \int_a^x f(t) dt, \quad a < x < b,$$

implies

$$F'(x) = f(x), \quad a < x < b,$$

and

$$F(x) = \int_x^b f(t) dt, \quad a < x < b,$$

implies

$$F'(x) = -f(x), \quad a < x < b.$$

To see this, for the first implication, write $\int_a^x f(t) dt = \int_a^c f(t) dt + F_c(x)$, and use $F'_c(x) = f(x)$. Since, by additivity, $\int_a^x f(t) dt + \int_x^b f(t) dt$ equals the constant $\int_a^b f(x) dx$, the second implication follows. \square

For example, if

$$F(x) = \int_0^{\tan x} e^{-t^2} dt, \quad 0 < x < \frac{\pi}{2},$$

then, $F'(x) = e^{-\tan^2 x} \sec^2 x$ by the above theorem combined with the chain rule. We will need this in §5.4.

The last two results show that integrals yield primitives. This is one version of the *fundamental theorem of calculus*. The other version of the fundamental theorem states that primitives yield integrals. When one is seeking areas or integrals, it is this version that is all-important.

Fundamental Theorem of Calculus. *Let f be nonnegative or integrable over (a, b) . Suppose that f is continuous on (a, b) , and let F be any primitive of f on (a, b) . Then, $F(b-)$ and $F(a+)$ exist, and*

$$\int_a^b f(x) dx = F(b-) - F(a+).$$

To see this, first, assume that f is nonnegative. Then, F is increasing ($F' = f \geq 0$). Hence, $F(b-)$ and $F(a+)$ exist for any primitive F . In particular, with F_c as above, $F_c(b-)$ and $F_c(a+)$ exist. Since $F_c - F = k$ is a constant, by continuity at the endpoints,

$$\begin{aligned} \int_a^b f(x) dx &= \int_a^c f(x) dx + \int_c^b f(x) dx \\ &= \lim_{n \nearrow \infty} \int_{a+1/n}^c f(x) dx + \lim_{n \nearrow \infty} \int_c^{b-1/n} f(x) dx \\ &= -\lim_{n \nearrow \infty} F_c(a + 1/n) + \lim_{n \nearrow \infty} F_c(b - 1/n) \\ &= F_c(b-) - F_c(a+) \\ &= (F(b-) + k) - (F(a+) + k) = F(b-) - F(a+). \end{aligned}$$

For the integrable case, let F^\pm denote primitives of f^\pm (here, F^\pm are *not* the positive and negative parts of F). Then, $F^+ - F^-$ differs from any primitive F of

f by a constant k . Since $F^\pm(b-)$ and $F^\pm(a+)$ exist, so do $F(b-)$ and $F(a+)$. Hence,

$$\begin{aligned}\int_a^b f(x) dx &= \int_a^b f^+(x) dx - \int_a^b f^-(x) dx \\ &= F^+(b-) - F^+(a+) - F^-(b-) + F^-(a+) \\ &= F(b-) + k - F(a+) - k = F(b-) - F(a+). \quad \square\end{aligned}$$

Note that, in the fundamental theorem, as stated above, a or b or $F(a+)$ or $F(b-)$ may be infinite.

When $a, b, F(b-)$ and $F(a+)$ are all finite, the fundamental theorem simplifies slightly. Indeed, in this case, by defining $F(b) = F(b-)$, $F(a) = F(a+)$, the primitive F extends to a continuous function on the compact interval $[a, b]$ and the fundamental theorem becomes

$$\int_a^b f(x) dx = F(b) - F(a).$$

In particular, this simpler form of the fundamental theorem applies when f and (a, b) are both bounded. All primitives displayed below were obtained in §3.6.

For example, $\sin x$ is bounded and has the primitive $-\cos x$ on $(0, \pi)$. So,

$$\int_0^\pi \sin x dx = (-\cos \pi) - (-\cos 0) = 2.$$

Similarly, x^n , $n \geq 0$, is bounded and has the primitive $x^{n+1}/(n+1)$ over any bounded interval (a, b) , so,

$$(4.1) \quad \int_a^b x^n dx = \frac{b^{n+1}}{n+1} - \frac{a^{n+1}}{n+1}, \quad n \geq 0.$$

Below, it is convenient to denote $F(b) - F(a) = F(x)|_a^b$. Since, in §3.6, a primitive of f was written $\int f(x) dx$, the fundamental theorem becomes

$$\int_a^b f(x) dx = \int f(x) dx \Big|_a^b.$$

This explains the notation $\int f(x) dx$ for primitives. (The notation $\int_a^b f(x) dx$ for integrals was explained in §4.3.)

Also $f(x) = 1/\sqrt{x(1-x)} > 0$ has the primitive $F(x) = 2 \arcsin \sqrt{x}$ continuous over $[0, 1]$, so,

$$\int_0^1 \frac{dx}{\sqrt{x(1-x)}} = 2 \arcsin \sqrt{x} \Big|_0^1 = 2 \arcsin 1 = \pi.$$

Similarly, since $f(x) = 1/(1+x^2)$ is nonnegative and has the primitive $F(x) = \arctan x$ over \mathbb{R} ,

$$\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \arctan x \Big|_{-\infty}^{\infty} = \frac{\pi}{2} - \left(-\frac{\pi}{2}\right) = \pi.$$

The unit disk

$$D = \{(x, y) : x^2 + y^2 < 1\}$$

is the disjoint union of a horizontal line segment and the two half-disks

$$D^\pm = \{(x, y) : x^2 + y^2 < 1, \pm y > 0\}.$$

Then, $\text{area}(D) = 2 \cdot \text{area}(D^+)$ (Exercise 11 of §4.2). But D^+ is the subgraph of $f(x) = \sqrt{1 - x^2}$ over $(-1, 1)$, which has a primitive continuous on $[-1, 1]$. Hence,

$$\int_{-1}^1 \sqrt{1 - x^2} dx = \frac{1}{2} \left(\arcsin x + x\sqrt{1 - x^2} \right) \Big|_{-1}^1 = \frac{\pi}{2}.$$

This yields the following.

Theorem. *The area of the unit disk is π .* \square

Of course, by translation and dilation invariance, the area of any disk of radius $r > 0$ is πr^2 . Another integral is

$$\int_0^1 (-\log x) dx = (x - x \log x) \Big|_0^1 = 1 + \lim_{x \rightarrow 0^+} x \log x = 1 + 0 = 1.$$

Our next item is the linearity of the integral.

Theorem (Linearity). *Suppose that f, g are continuous on (a, b) . If f and g are both nonnegative or both integrable over (a, b) , then,*

$$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$$

To see this, let F and G be primitives corresponding to f and g . Then, $f + g = F' + G' = (F + G)'$. So, $F + G$ is a primitive of $f + g$. By the fundamental theorem,

$$\begin{aligned} \int_a^b [f(x) + g(x)] dx &= F(b-) + G(b-) - F(a+) - G(a+) \\ &= \int_a^b f(x) dx + \int_a^b g(x) dx. \quad \square \end{aligned}$$

We say $f : (a, b) \rightarrow \mathbb{R}$ is *piecewise continuous* if there is a partition $a = x_0 < x_1 < \dots < x_{n+1} = b$, such that f is continuous on each subinterval (x_{i-1}, x_i) , $i = 1, \dots, n + 1$. Now, by additivity, the integral \int_a^b can be broken up into $\int_{x_{i-1}}^{x_i}$, $i = 1, \dots, n + 1$. We conclude that *linearity also holds for piecewise continuous functions*.

By induction, linearity holds for finitely many (piecewise) continuous functions. If f_1, \dots, f_n are (piecewise) continuous and all nonnegative or all integrable over

(a, b) , then,

$$\int_a^b \sum_{k=1}^n f_k(x) dx = \sum_{k=1}^n \int_a^b f_k(x) dx.$$

Since primitives are connected to integrals by the fundamental theorem, there is an integration by parts (§3.6) result for integrals.

Theorem (Integration By Parts). *Let f and g be differentiable on (a, b) with $f'g$ and fg' continuous. If $f'g$ and fg' are both nonnegative or both integrable, then,*

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_{a+}^{b-} - \int_a^b f'(x)g(x) dx.$$

This follows by applying the fundamental theorem to $f'g + fg' = (fg)'$ and using linearity. \square

Since primitives are connected to integrals by the fundamental theorem, there is a substitution (§3.6) result for integrals. Recall (§2.3) that continuous strictly monotone functions map open intervals to open intervals.

Theorem (Substitution). *Let g be differentiable and strictly monotone on an interval (a, b) with g' continuous, and let $(m, M) = g[(a, b)]$. Let $f : (m, M) \rightarrow \mathbb{R}$ be continuous. If f is nonnegative or integrable over (m, M) , then, $f[g(t)]|g'(t)|$ is nonnegative or integrable over (a, b) , and*

$$(4.2) \quad \int_m^M f(x) dx = \int_a^b f[g(t)]|g'(t)| dt.$$

To see this, first, assume that g is strictly increasing and f is nonnegative, let F be a primitive of f , let $H(t) = F[g(t)]$, and let $h(t) = f[g(t)]g'(t)$. Then, $(m, M) = (g(a+), g(b-))$ and $H'(t) = F'[g(t)]g'(t) = f[g(t)]g'(t) = h(t)$ by the chain rule. Hence, H is a primitive for h . Moreover, h is continuous and nonnegative, $F(M-) = H(b-)$, and $F(m+) = H(a+)$. By the fundamental theorem,

$$\begin{aligned} \int_m^M f(x) dx &= F(M-) - F(m+) \\ &= H(b-) - H(a+) \\ &= \int_a^b h(t) dt \\ &= \int_a^b f[g(t)]g'(t) dt. \end{aligned}$$

Since $|g'(t)| = g'(t)$, this establishes the case with g strictly increasing and f nonnegative. If f is integrable, apply the nonnegative case to f^\pm . Since the

positive and negative parts of $f[g(t)]g'(t)$ are $f^\pm[g(t)]g'(t)$, the integrable case follows.

If g is strictly decreasing, then, $(m, M) = (g(b-), g(a+))$. Now, $h(t) = g(-t)$ is strictly increasing, $h((-b, -a)) = (m, M)$, and $h'(-t) = -g'(t) = |g'(t)|$ is nonnegative on (a, b) . Applying what we just learned to f and h over $(-b, -a)$ yields

$$\begin{aligned}\int_m^M f(x) dx &= \int_{-b}^{-a} f[h(t)]h'(t) dt \\ &= \int_a^b f[h(-t)]h'(-t) dt \\ &= \int_a^b f[g(t)]|g'(t)| dt. \quad \square\end{aligned}$$

If g is not monotone, then, (4.2) has to be reformulated (Exercise 21). To see what happens, let us consider a simple example with $f(x) = 1$. Let $g : (a, b) \rightarrow (m, M)$ be *piecewise linear* with line segments inclined at $\pm\pi/4$. By this, we mean g is continuous on (a, b) and the graph of g is a line segment with slope ± 1 on each subinterval (t_{i-1}, t_i) , $i = 1, \dots, n+1$, for some partition $a = t_0 < t_1 < \dots < t_{n+1} = b$ of (a, b) (Figure 4.29). Then, $|g'(t)| = 1$ for all but finitely many t , so, $\int_a^b |g'(t)| dt = b - a$. On the other hand, substituting $f(x) = 1$ in (4.2) gives $\int_a^b |g'(t)| dt = M - m$. Thus, in such a situation, (4.2) cannot be correct unless the domain and the range have the same length, i.e., $M - m = b - a$.

To fix this, we have to take into account the extent to which g is not a bijection. To this end, for each x in (m, M) , let $\#(x)$ denote the number of points in the inverse image $g^{-1}(\{x\})$. Since (m, M) is the range of g , $\#(x) \geq 1$ for all x . The correct replacement for (4.2) with $f(x) = 1$ is

$$(4.3) \quad \int_m^M \#(x) dx = \int_a^b |g'(t)| dt.$$


This holds as long as g is continuous on (a, b) and there is a partition $a = t_0 < t_1 < \dots < t_{n+1} = b$ of (a, b) with g differentiable, g' continuous, and g strictly monotone on each subinterval (t_{i-1}, t_i) , for each $i = 1, \dots, n+1$ (Exercise 21).

For example, supposing $g : (a, b) \rightarrow (m, M)$ piecewise linear with slopes ± 1 , reduces (4.3) to $\int_m^M \#(x) dx = b - a$. Dividing by $M - m$ yields

$$(4.4) \quad \frac{1}{M - m} \int_m^M \#(x) dx = \frac{b - a}{M - m}.$$

Now, the left side of (4.4) may be thought of as the average value of $\#(x)$ over (m, M) . We conclude that, for a piecewise linear g with slopes ± 1 , the average value of the number of inverse images equals the ratio of the lengths of the domain over the range.

Now, we derive the integral version of

FIGURE 4.29. Piecewise linear: $\#(x) = 4$, $\#(x') = 3$.

Taylor's Theorem. Let $n \geq 0$ and suppose that f is $(n+1)$ times differentiable on (a, b) , with $f^{(n+1)}$ continuous on (a, b) . Suppose that $f^{(n+1)}$ is nonnegative or integrable over (a, b) , and fix $a < c < b$. Then,

$$f(x) = f(c) + f'(c)(x - c) + \frac{f''(c)}{2!}(x - c)^2 + \dots \\ \dots + \frac{f^{(n)}(c)}{n!}(x - c)^n + \frac{h_{n+1}(x)}{(n+1)!}(x - c)^{n+1},$$

where

$$h_{n+1}(x) = (n+1) \int_0^1 (1-s)^n f^{(n+1)}[c + s(x-c)] ds.$$

To see this, recall, in §3.4, that we obtained $R_{n+1}(x, x) = 0$ and (here, ' denotes derivative with respect to t)

$$R'_{n+1}(x, t) = -\frac{f^{(n+1)}(t)}{n!}(x - t)^n.$$

Now, apply the fundamental theorem to $-R'_{n+1}(x, t)$ and substitute $t = c + s(x - c)$, $dt = (x - c)ds$, obtaining

$$R_{n+1}(x, c) = \frac{1}{n!} \int_c^x f^{(n+1)}(t)(x - t)^n dt \\ = \frac{(x - c)^{n+1}}{n!} \int_0^1 f^{(n+1)}(c + s(x - c))(1 - s)^n ds \\ = \frac{(x - c)^{n+1}}{(n+1)!} h_{n+1}(x). \quad \square$$

In contrast with the Lagrange and Cauchy forms (§3.4) of the remainder, here, we need continuity and nonnegativity or integrability of $f^{(n+1)}$.

Our last item is the integration of power series. Since we already know (§3.6) how to find primitives of power series, the fundamental theorem and (4.1) yield the following.

Theorem. Suppose that $R > 0$ is the radius of convergence of

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

If $[a, b] \subset (-R, R)$, then,

$$(4.5) \quad \int_a^b f(x) dx = \sum_{n=0}^{\infty} \int_a^b a_n x^n dx. \quad \square$$

For example, substituting $-x^2$ for x in the exponential series,

$$e^{-x^2} = 1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \frac{x^8}{4!} - \dots$$

Integrating this over $(0, 1)$, we obtain

$$\int_0^1 e^{-x^2} dx = 1 - \frac{1}{1!3} + \frac{1}{2!5} - \frac{1}{3!7} + \frac{1}{4!9} - \dots$$

This last result is, in general, false if $a = -R$ or $b = R$. For example, with $f(x) = e^{-x} = \sum_{n=0}^{\infty} (-1)^n x^n / n!$ and $(a, b) = (0, \infty)$, (4.5) reads $1 = \infty - \infty + \infty - \infty + \dots$. Under additional assumptions, however, (4.5) is true, even in these cases (see §5.2).

Exercises 4.4

1. Compute $\int_0^{\infty} e^{-sx} dx$ for $s > 0$.
2. Compute $\int_0^1 x^{r-1} dx$ and $\int_1^{\infty} x^{r-1} dx$ and $\int_0^{\infty} x^{r-1} dx$ for r real. (There are three cases, $r < 0$, $r = 0$, and $r > 0$.)
3. Suppose that f is continuous over (a, b) , and let F be any primitive. If f and (a, b) are both bounded, then, f is integrable, and $F(a+)$ and $F(b-)$ are finite.
4. Let $f(x) = \sin x / x$, $x > 0$, and let $F(b) = \int_0^b f(x) dx$, $b > 0$. Show that $F(\infty) = \lim_{b \rightarrow \infty} F(b)$ exists and is finite. (Write $F(b) = \int_0^1 f(x) dx + \int_1^b f(x) dx$, integrate the second integral by parts, and use (3.1). This limit is computed in Exercise 12 of §5.4.)
5. For f continuous and nonnegative or integrable over $(0, 1)$,

$$\int_0^1 f(x) dx = \int_0^{\infty} e^{-t} f(e^{-t}) dt.$$

6. Compute $\int_0^{\infty} e^{-sx} x^n dx$ for $s > 0$ and $n \geq 0$. (Integration by parts.)
7. Compute $\int_0^{\infty} e^{-nx} \sin(sx) dx$ and $\int_0^{\infty} e^{-nx} \cos(sx) dx$ for $n \geq 1$. (Integration by parts.)

8. Show that $\int_0^\infty e^{-t^2/2} t^x dt = (x-1) \int_0^\infty e^{-t^2/2} t^{x-2} dt$ for $x > 1$. Use this to derive

$$\int_0^\infty e^{-t^2/2} t^{2n+1} dt = 2^n n!, \quad n \geq 0.$$

(Integration by parts.)

9. Compute $\int_0^1 (1-t)^n t^{x-1} dt$ for $x > 0$ and $n \geq 1$. (Integration by parts.)

10. Compute $\int_0^1 (-\log x)^n dx$.

11. ★ Show that

$$\int_{-1}^1 (x^2 - 1)^n dx = (-1)^n \frac{2n \cdot (2n-2) \cdot \dots \cdot 2}{(2n+1) \cdot (2n-1) \cdot \dots \cdot 3} \cdot 2.$$

(Integrate by parts.)

12. ★ For $n \geq 0$, the *Legendre polynomial* P_n (of degree n) is given by $P_n(x) = f^{(n)}(x)/2^n n!$, where $f(x) = (x^2 - 1)^n$. Show that

$$\int_{-1}^1 P_n(x)^2 dx = \frac{2}{2n+1}.$$

13. Use the integral test (§4.3) to show that

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad s > 1,$$

converges.

14. Use the integral test (§4.3) to show that

$$\gamma = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \log n \right)$$

exists and $0 < \gamma < 1$. This particular real γ is *Euler's constant*.

15. Compute $\int_{-\pi}^{\pi} x \cos(nx) dx$ and $\int_{-\pi}^{\pi} x \sin(nx) dx$ for $n \geq 0$. (Integration by parts.)

16. ★ Compute $\int_{-\pi}^{\pi} f(nx)g(mx) dx$, $n, m \geq 0$, with $f(x)$ and $g(x)$ equal to $\sin x$ or $\cos x$ (three possibilities — use (5.3) in §3.5).

17. ★ If $f, g : (a, b) \rightarrow \mathbb{R}$ are nonnegative and continuous, derive the *Cauchy-Schwarz inequality*

$$\left[\int_a^b f(x)g(x) dx \right]^2 \leq \left[\int_a^b f(x)^2 dx \right] \cdot \left[\int_a^b g(x)^2 dx \right].$$

(Use the fact that $q(t) = \int_a^b [f(x) + tg(x)]^2 dx$ is a nonnegative quadratic polynomial and Exercise 5 of §1.4.)

18. For $n \geq 1$, show that

$$\int_0^n \frac{1 - (1-t/n)^n}{t} dt = 1 + \frac{1}{2} + \dots + \frac{1}{n}.$$

19. We say $f : (a, b) \rightarrow \mathbf{R}$ is *piecewise differentiable* if there is a partition $a = x_0 < x_1 < \cdots < x_{n+1} = b$, such that f restricted to (x_{i-1}, x_i) is differentiable for $i = 1, \dots, n+1$. Let $f : (a, b) \rightarrow \mathbf{R}$ be piecewise continuous and integrable. Show that $F(x) = \int_a^x f(t) dt$, $a < x < b$, is continuous on (a, b) and piecewise differentiable on (a, b) .

20. ★ If $f : (a, b) \rightarrow \mathbf{R}$ is nonnegative and $g : [a, b] \rightarrow \mathbf{R}$ is nonnegative and continuous, then,

$$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$$

(Use Exercise 5 of §4.3 and approximate g by a piecewise constant g_ϵ as in §2.3. Since f is arbitrary, linearity may not be used directly.)

21. ★ Suppose that $g : (a, b) \rightarrow (m, M)$ is continuous, and suppose that there is a partition $a = t_0 < t_1 < \cdots < t_{n+1} = b$ of (a, b) , such that g is differentiable, g' is continuous, and g is strictly monotone on each subinterval (t_{i-1}, t_i) , for each $i = 1, \dots, n+1$. For each x in (m, M) , let $\#(x)$ denote the number of points in the inverse image $g^{-1}(\{x\})$. Also let $f : (m, M) \rightarrow \mathbf{R}$ be continuous and nonnegative. Then,

$$\int_m^M f(x)\#(x) dx = \int_a^b f[g(t)]|g'(t)| dt.$$

(Use additivity on the integral \int_a^b .)

22. ★ Let f be differentiable with f' continuous on an interval containing $[a, b]$. Show that the variation of f corresponding to any partition in (a, b) (Exercise 4 of §2.2) is bounded by $\int_a^b |f'(x)| dx$. Use Exercise 3 of §4.3 to show that the total variation of f over (a, b) equals $\int_a^b |f'(x)| dx$. (Rewrite the variation of f over a given partition as a Riemann sum for $|f'|$.)

§4.5 The Method of Exhaustion

In this section, we compute the area of the unit disk D via the Method of Exhaustion.

For $n \geq 3$, let $P_k = (\cos(2\pi k/n), \sin(2\pi k/n))$, $0 \leq k \leq n$. Then, the points P_k are evenly spaced about the unit circle $\{(x, y) : x^2 + y^2 = 1\}$, and $P_n = P_0$. Let $D_n \subset D$ be the interior of the inscribed regular n -sided polygon obtained by joining the points P_0, P_1, \dots, P_n (we do not include the edges of D_n in the definition of D_n). Then (Exercise 13 of §4.2),

$$\text{area}(D_n) = \frac{n}{2} \sin(2\pi/n) = \pi \cdot \frac{\sin(2\pi/n)}{2\pi/n}.$$

Since $\lim_{x \rightarrow 0} \frac{\sin x}{x} = \sin' 0 = \cos 0 = 1$, we obtain

$$(5.1) \quad \lim_{n \nearrow \infty} \text{area}(D_n) = \pi.$$

Since

$$(5.2) \quad D_4 \subset D_8 \subset D_{16} \subset \dots, \quad \text{and} \quad D = \bigcup_{n=2}^{\infty} D_{2^n},$$

it is reasonable to make the guess that

$$(5.3) \quad \text{area}(D) = \lim_{n \nearrow \infty} \text{area}(D_{2^n}),$$

and, hence, conclude that $\text{area}(D) = \pi$. The reasoning that leads from (5.2) to (5.3) is generally correct. The result is called the *Method of Exhaustion*.

Although $\text{area}(D)$ was computed in the previous section using the fundamental theorem, in Chapter 5 we will need the Method to compute other areas.

We say that a sequence of sets (A_n) is *increasing* (Figure 4.30) if $A_1 \subset A_2 \subset A_3 \subset \dots$.




FIGURE 4.30. An increasing sequence of sets.

Theorem (Method of Exhaustion). *If $A_1 \subset A_2 \subset \dots$ is an increasing sequence of subsets of \mathbb{R}^2 , then,*

$$\text{area} \left(\bigcup_{n=1}^{\infty} A_n \right) = \lim_{n \nearrow \infty} \text{area}(A_n).$$

We warn the reader that the result is false, in general, for decreasing sequences. For example, take $A_n = (n, \infty) \times (-\infty, \infty)$, $n \geq 1$. Then, $\text{area}(A_n) = \infty$ for all $n \geq 1$, but $\bigcap_{n=1}^{\infty} A_n = \emptyset$ so $\text{area}(\bigcap_{n=1}^{\infty} A_n) = 0$. This lack of symmetry between increasing and decreasing sequences is a reflection of the lack of symmetry in the definition of area: $\text{area}(A)$ is defined as an inf of over-estimates, not as a sup of underestimates.

The Method is established in three stages: first, when (definitions below) the sets A_n , $n \geq 1$, are open; then, when the sets A_n , $n \geq 1$, are interopen; finally, for

arbitrary sets A_n , $n \geq 1$. *Open* and *interopen* are structural properties of sets that we describe below.

We call a set $G \subset \mathbb{R}^2$ *open* if every point $(a, b) \in G$ can be surrounded by a nonempty, open rectangle wholly contained in G . For example, an open rectangle is an open set, but a compact rectangle Q is not, since no point on the edges of Q can be surrounded by a rectangle wholly contained in Q . The n -sided polygon D_n , considered above, is an open set as is the unit disk D . Since there are no points in \emptyset for which the open criterion fails, \emptyset is open.

For our purposes, the most important example of an open set is given by the following.

Theorem. *If $f \geq 0$ is continuous on (a, b) , its subgraph is an open subset of \mathbb{R}^2 .*

To see this, pick x and y with $a < x < b$ and $0 < y < f(x)$. We have to find a rectangle Q containing (x, y) and contained in the subgraph. Pick $y < y_1 < f(x)$. We claim there is a $c > 0$, such that $|t - x| < c$ implies $a < t < b$ and $f(t) > y_1$. If not, then, for all $n \geq 1$, we can find a real t_n in the interval $(x - 1/n, x + 1/n)$ contradicting the stated property, i.e., satisfying $f(t_n) \leq y_1$. Then, $t_n \rightarrow x$, so, by continuity $f(t_n) \rightarrow f(x)$. Hence, $f(x) \leq y_1$, contradicting our initial choice of y_1 . Thus, there is a $c > 0$, such that the rectangle $Q = (x - c, x + c) \times (0, y_1)$ contains (x, y) and lies in the subgraph. \square

Thus, *the integral of a continuous nonnegative function is the area of an open set.*

An alternative description of open sets is in terms of distance. If (a, b) is a point and A is a set, then, the distance $d((a, b), A)$ between the point (a, b) and A , by definition, is the distance between the set $\{(a, b)\}$ and the set A (§4.2). For example, if Q is an open rectangle and $(a, b) \in Q$, then, $d((a, b), Q^c)$ is positive. Here and below, $A^c = \mathbb{R}^2 \setminus A$.

Theorem. *A set G is open iff $d((a, b), G^c) > 0$ for all points $(a, b) \in G$.*

Indeed, if $(a, b) \in G$ and $Q \subset G$ contains (a, b) , then, $Q^c \supset G^c$, so, $d((a, b), G^c) \geq d((a, b), Q^c) > 0$. Conversely, if $d = d((a, b), G^c) > 0$, then, the disk R of radius $d/2$ and center (a, b) lies wholly in G . Now, choose any rectangle Q in R containing (a, b) . \square

If G, G' are open subsets, so, are $G \cup G'$ and $G \cap G'$. In fact, if (G_n) is a sequence of open sets, then, $G = \bigcup_{n=1}^{\infty} G_n$ is open. To see this, if $(a, b) \in G$, then, $(a, b) \in G_n$ for some specific n . Since the specific G_n is open, there is a rectangle Q with $(a, b) \in Q \subset G_n \subset G$. Hence, G is open. Thus, an infinite union of open sets is open. If G_1, \dots, G_n are finitely many open sets, then, $G = G_1 \cap G_2 \cap \dots \cap G_n$ is open. To see this, if $(a, b) \in G$, then, $(a, b) \in G_k$ for all $1 \leq k \leq n$, so, there are open rectangles Q_k with $(a, b) \in Q_k \subset G_k$, $1 \leq k \leq n$. Hence, $Q = \bigcap_{k=1}^n Q_k$ is an open rectangle containing (a, b) and contained in G (a finite intersection of open rectangles is an open rectangle). Thus,

a finite intersection of open sets is open. However, an infinite intersection of open sets need not be open.

If $A \subset \mathbb{R}^2$ is any set and $\epsilon > 0$, by definition of area, we can find an open set G containing A and satisfying $\text{area}(G) \leq \text{area}(A) + \epsilon$ (Exercise 6). If we had additivity and $\text{area}(A) < \infty$, writing $\text{area}(G) = \text{area}(A) + \text{area}(G \setminus A)$, we would conclude that $\text{area}(G \setminus A) \leq \epsilon$. Conversely, if we are seeking properties of sets that guarantee additivity, we may, instead, focus on those sets M in \mathbb{R}^2 satisfying the above approximability condition: *For all $\epsilon > 0$, there is an open superset G of M , such that $\text{area}(G \setminus M) \leq \epsilon$.* Instead of doing this, however, it will be quicker for us to start with an alternate equivalent (Exercise 16) formulation.

We say a set $M \subset \mathbb{R}^2$ is *measurable* if

$$(5.4) \quad \text{area}(A) = \text{area}(A \cap M) + \text{area}(A \cap M^c), \quad \text{for all } A \subset \mathbb{R}^2.$$

For example, the empty set is measurable and M is measurable iff M^c is measurable. Below, we show that every open set is measurable. Measurability may be looked upon as a strengthened form of additivity, since the equality in (5.4) is required to hold for *every* $A \subset \mathbb{R}^2$. Note that the trick, below, of summing alternate areas C_1, C_3, C_5, \dots was already used in deriving additivity in §4.3. Compare the next derivation with that derivation!

In §4.2, we established additivity when the sets were well separated. Now, we establish a similar result involving open sets.

Theorem. *If G is open, then, G is measurable.*

To see this, we need show only that

$$(5.5) \quad \text{area}(A) \geq \text{area}(A \cap G) + \text{area}(A \cap G^c)$$

for every $A \subset \mathbb{R}^2$, since the reverse inequality follows by subadditivity. Let $A \subset \mathbb{R}^2$ be arbitrary. If $\text{area}(A) = \infty$, (5.5) is immediate, so, let us assume that $\text{area}(A) < \infty$. Let G_n be the set of points in G whose distance from G^c is at least $1/n$. Since $A \cap G_n$ and $A \cap G^c$ are well separated (Figure 4.31),

$$\text{area}(A) \geq \text{area}(A \cap G_n) + \text{area}(A \cap G^c).$$

By subadditivity,

$$\text{area}(A \cap G) \leq \text{area}(A \cap G_n) + \text{area}(A \cap G \cap G_n^c).$$

Combining the last two inequalities, we obtain

$$(5.6) \quad \text{area}(A) \geq \text{area}(A \cap G) + \text{area}(A \cap G^c) - \text{area}(A \cap G \cap G_n^c).$$

Thus, if we show that

$$(5.7) \quad \lim_{n \nearrow \infty} \text{area}(A \cap G \cap G_n^c) = 0,$$

letting $n \nearrow \infty$ in (5.6), we obtain (5.5), hence, the result.

To obtain (5.7), let C_n be the set of points (a, b) in G satisfying $1/(n+1) \leq d((a, b), G^c) < 1/n$. Since G is open, $d((a, b), G^c) > 0$ for every point in G . Thus,

$$G \cap G_n^c = C_n \cup C_{n+1} \cup C_{n+2} \cup \dots$$

But the sets $C_n, C_{n+2}, C_{n+4}, \dots$, are well separated. Hence,

$$\text{area}(A \cap C_n) + \text{area}(A \cap C_{n+2}) + \text{area}(A \cap C_{n+4}) + \dots \leq \text{area}(A \cap G).$$

Since $C_{n+1}, C_{n+3}, C_{n+5}, \dots$, are well separated,

$$\text{area}(A \cap C_{n+1}) + \text{area}(A \cap C_{n+3}) + \text{area}(A \cap C_{n+5}) + \dots \leq \text{area}(A \cap G).$$

Adding the last two inequalities, by subadditivity, we obtain

$$(5.8) \quad \text{area}(A \cap G \cap G_n^c) \leq \sum_{k=n}^{\infty} \text{area}(A \cap C_k) \leq 2 \text{area}(A \cap G) < \infty.$$

Now, (5.8) with $n = 1$ shows that the series $\sum_{k=1}^{\infty} \text{area}(A \cap C_k)$ converges. Thus, the tail series, starting from $k = n$ in (5.8), approaches zero, as $n \nearrow \infty$. This establishes (5.7). \square




FIGURE 4.31. An open set is measurable.

Now we establish the Method for measurable, hence, for open sets. In fact, we need to establish a strengthened form of the Method for measurable sets.

Theorem (Measurable Method of Exhaustion). *If $M_1 \subset M_2 \subset \dots$ is an increasing sequence of measurable subsets of \mathbb{R}^2 and $A \subset \mathbb{R}^2$ is arbitrary, then,*

$$\text{area} \left[A \cap \left(\bigcup_{n=1}^{\infty} M_n \right) \right] = \lim_{n \nearrow \infty} \text{area}(A \cap M_n).$$

As stated previously, the Method is obtained by choosing $A = \mathbb{R}^2$. To derive this, let $M_{\infty} = \bigcup_{n=1}^{\infty} M_n$. Since $A \cap M_n \subset A \cap M_{\infty}$, by monotonicity, the sequence $(\text{area}(A \cap M_n))$ is increasing and bounded above by $\text{area}(A \cap M_{\infty})$. Thus,

$$\lim_{n \nearrow \infty} \text{area}(A \cap M_n) \leq \text{area}(A \cap M_{\infty}).$$

To obtain the reverse inequality, apply (5.4) with M and A , there, replaced by M_1 and $A \cap M_2$ respectively, obtaining

$$\text{area}(A \cap M_2) = \text{area}(A \cap M_2 \cap M_1) + \text{area}(A \cap M_2 \cap M_1^c).$$

Since $A \cap M_2 \cap M_1 = A \cap M_1$, this implies that

$$\text{area}(A \cap M_2) = \text{area}(A \cap M_1) + \text{area}(A \cap M_2 \cap M_1^c).$$

Now, apply (5.4) with M and A , there, replaced by M_2 and $A \cap M_3$ respectively, obtaining

$$\begin{aligned} \text{area}(A \cap M_3) &= \text{area}(A \cap M_2) + \text{area}(A \cap M_3 \cap M_2^c) \\ &= \text{area}(A \cap M_1) + \text{area}(A \cap M_2 \cap M_1^c) \\ &\quad + \text{area}(A \cap M_3 \cap M_2^c). \end{aligned}$$

Proceeding in this manner, we obtain

$$\text{area}(A \cap M_n) = \text{area}(A \cap M_1) + \sum_{k=2}^n \text{area}(A \cap M_k \cap M_{k-1}^c).$$

Sending $n \nearrow \infty$, we obtain

$$\lim_{n \nearrow \infty} \text{area}(A \cap M_n) = \text{area}(A \cap M_1) + \sum_{k=2}^{\infty} \text{area}(A \cap M_k \cap M_{k-1}^c).$$

Since

$$M_1 \cup (M_2 \cap M_1^c) \cup (M_3 \cap M_2^c) \cup \dots = M_{\infty},$$

subadditivity implies that

$$\text{area}(A \cap M_{\infty}) \leq \text{area}(A \cap M_1) + \sum_{k=2}^{\infty} \text{area}(A \cap M_k \cap M_{k-1}^c).$$

Hence, we obtain the reverse inequality

$$\lim_{n \nearrow \infty} \text{area}(A \cap M_n) \geq \text{area}(A \cap M_{\infty}). \quad \square$$

By choosing $A = \mathbb{R}^2$, we conclude that the Method is valid for measurable, hence, open sets. *This completes stage one of the derivation of the Method.*

Next we establish the Method for interopen sets. A set $I \subset \mathbb{R}^2$ is *interopen* if I is the infinite intersection of a sequence of open sets (G_n) , $I = \bigcap_{n=1}^{\infty} G_n$. Of course, every open set is interopen. Also, every compact rectangle is interopen (Exercise 5). The key feature of interopen sets is that any set A can be covered by some interopen set I , $A \subset I$, having the same area, $\text{area}(A) = \text{area}(I)$ (Exercise 7).

Theorem. *If (M_n) is a sequence of measurable sets, then, $\bigcap_{n=1}^{\infty} M_n$ is measurable.*

To derive this theorem, we start with two measurable sets M, N , and we show that $M \cap N$ is measurable. First, note that

$$(5.9) \quad (M \cap N)^c = (M \cap N^c) \cup (M^c \cap N) \cup (M^c \cap N^c).$$

Let $A \subset \mathbb{R}^2$ be arbitrary. Since N is measurable, write (5.4) with $A \cap M$ and N replacing A and M , respectively, obtaining

$$\text{area}(A \cap M) = \text{area}(A \cap M \cap N) + \text{area}(A \cap M \cap N^c).$$

Now, write (5.4) with $A \cap M^c$ and N replacing A and M respectively, obtaining

$$\text{area}(A \cap M^c) = \text{area}(A \cap M^c \cap N) + \text{area}(A \cap M^c \cap N^c).$$

Now, insert the last two equalities in (5.4). By (5.9) and subadditivity, we obtain

$$\begin{aligned} \text{area}(A) &= \text{area}(A \cap M) + \text{area}(A \cap M^c) \\ &= \text{area}(A \cap (M \cap N)) + \text{area}(A \cap (M \cap N^c)) \\ &\quad + \text{area}(A \cap (M^c \cap N)) + \text{area}(A \cap (M^c \cap N^c)) \\ &\geq \text{area}(A \cap (M \cap N)) + \text{area}(A \cap (M \cap N)^c). \end{aligned}$$

Hence,

$$\text{area}(A) \geq \text{area}(A \cap (M \cap N)) + \text{area}(A \cap (M \cap N)^c).$$

Since the reverse inequality is an immediate consequence of subadditivity, we conclude that $M \cap N$ is measurable.

Now, let (M_n) be a sequence of measurable sets and set $N_n = \bigcap_{k=1}^n M_k, n \geq 1$. Then, $N_n, n \geq 1$, are measurable. Indeed, $N_1 = M_1$ is measurable. For the inductive step, suppose that N_n is measurable. Since $N_{n+1} = N_n \cap M_{n+1}$, we conclude that N_{n+1} is measurable. Hence, by induction, N_n is measurable for all $n \geq 1$. Now, $M_\infty = \bigcap_{n=1}^\infty M_n = \bigcap_{n=1}^\infty N_n$ and $N_1 \supset N_2 \supset \dots$. Hence, $N_1^c \subset N_2^c \subset \dots$, so, by the measurable Method, we obtain

$$\begin{aligned} \text{area}(A \cap M_\infty^c) &= \text{area} \left[A \cap \left(\bigcap_{n=1}^\infty N_n \right)^c \right] \\ &= \text{area} \left(A \cap \bigcup_{n=1}^\infty N_n^c \right) \\ (5.10) \quad &= \lim_{n \nearrow \infty} \text{area}(A \cap N_n^c). \end{aligned}$$

Here, we used De Morgan's law (§1.1). Now, for each $n \geq 1$,

$$\begin{aligned} \text{area}(A) &= \text{area}(A \cap N_n) + \text{area}(A \cap N_n^c) \\ (5.11) \quad &\geq \text{area}(A \cap M_\infty) + \text{area}(A \cap N_n^c). \end{aligned}$$

Sending $n \nearrow \infty$ in (5.11) and using (5.10) yields

$$\text{area}(A) \geq \text{area}(A \cap M_\infty) + \text{area}(A \cap M_\infty^c).$$

Since the reverse inequality follows from subadditivity, we conclude that $M_\infty = \bigcap_{n=1}^\infty M_n$ is measurable. \square

By choosing (M_n) in the theorem to consist of open sets, we see that every interopen set is measurable. Hence, we conclude that the Method is valid for interopen sets. *This completes stage two of the derivation of the Method.*

The third and final stage of the derivation of the Method is to establish it for an increasing sequence of arbitrary sets. To this end, let $A_1 \subset A_2 \subset \dots$ be an arbitrary increasing sequence of sets. For each $n \geq 1$, by Exercise 7, choose an interopen set I_n containing A_n and having the same area: $I_n \supset A_n$ and $\text{area}(I_n) = \text{area}(A_n)$. For each $n \geq 1$, let

$$J_n = I_n \cap I_{n+1} \cap I_{n+2} \cap \dots$$

Then, J_n is interopen, $A_n \subset J_n \subset I_n$, and $\text{area}(J_n) = \text{area}(A_n)$, for all $n \geq 1$. Moreover, $J_n = I_n \cap J_{n+1}$. Hence, (and this is the reason for introducing the sequence (J_n)), the sequence (J_n) is increasing. Thus, by applying the Method for interopen sets,

$$\begin{aligned} \lim_{n \nearrow \infty} \text{area}(A_n) &= \lim_{n \nearrow \infty} \text{area}(J_n) \\ (5.12) \qquad &= \text{area} \left(\bigcup_{n=1}^\infty J_n \right) \geq \text{area} \left(\bigcup_{n=1}^\infty A_n \right). \end{aligned}$$

On the other hand, by monotonicity, the sequence $(\text{area}(A_n))$ is increasing and bounded above by $\text{area}(\bigcup_{n=1}^\infty A_n)$. Hence,

$$\lim_{n \nearrow \infty} \text{area}(A_n) \leq \text{area} \left(\bigcup_{n=1}^\infty A_n \right).$$

Combining this with (5.12), we conclude that

$$\lim_{n \nearrow \infty} \text{area}(A_n) = \text{area} \left(\bigcup_{n=1}^\infty A_n \right).$$

This completes stage three, hence, the derivation of the Method. \square

We end by describing the connection between the areas of the inscribed and circumscribed polygons of the unit disk D , as the number of sides doubles. Let

$$P_k = \left(\frac{\cos(2\pi k/n)}{\cos(\pi/n)}, \frac{\sin(2\pi k/n)}{\cos(\pi/n)} \right), \quad 0 \leq k \leq n.$$

Then, the points P_k are evenly spaced about the circle $\{(x, y) : x^2 + y^2 = \sec^2(\pi/n)\}$, and $P_n = P_0$. Let D'_n denote the interior of the regular n -sided polygon obtained by joining the points P_0, \dots, P_n by line segments. Then, $D'_n \supset D$ and $D'_n = c \cdot D_n$ with $c = \sec(\pi/n)$. Hence, by dilation invariance, we obtain

$$\text{area}(D'_n) = c^2 \cdot \text{area}(D_n) = n \tan(\pi/n)$$

which also goes to π as $n \nearrow \infty$.

Let a_n, a'_n denote the areas of the inscribed and circumscribed n -sided polygons D_n, D'_n , respectively. Then, using trigonometry, one obtains (Exercise 11)

$$(5.13g) \quad a_{2n} = \sqrt{a_n a'_n}$$

and

$$(5.13h) \quad \frac{1}{a'_{2n}} = \frac{1}{2} \left(\frac{1}{a_{2n}} + \frac{1}{a'_n} \right).$$

Since $a_4 = 2$ and $a'_4 = 4$, we obtain $a_8 = 2\sqrt{2}$ and $a'_8 = 8(\sqrt{2} - 1)$. Thus,

$$2\sqrt{2} < \pi < 8(\sqrt{2} - 1).$$

Continuing in this manner, one obtains approximations to π . These identities are very similar to those leading to Gauss' *arithmetic-geometric mean*, which we discuss in §5.3.

Exercises 4.5

1. If Q is an open rectangle and $(x, y) \in Q$, then, $d((x, y), Q^c) > 0$.
2. Find a sequence (A_n) of open sets, such that $\bigcap_{n=1}^{\infty} A_n$ is not open.
3. ★ A set A is *closed* if A^c is open. Show that a compact rectangle is closed, an infinite intersection of closed sets is closed, and a finite union of closed sets is closed. Find a sequence (A_n) of closed sets, such that $\bigcup_{n=1}^{\infty} A_n$ is not closed. (You will need De Morgan's law (§1.1).)
4. ★ Given a real a , let L_a denote the vertical infinite line through a , $L_a = \{(x, y) : x = a, y \in \mathbb{R}\}$. Also set $L_{\infty} = L_{-\infty} = \emptyset$. Let f be nonnegative and continuous on (a, b) . Show that

$$C = \{(x, y) : a < x < b, 0 \leq y \leq f(x)\} \cup L_a \cup L_b$$

is a closed set and

$$\int_a^b f(x) dx = \text{area}(C).$$

This shows that the integral of a continuous nonnegative function is also the area of a closed set. (Compare C with the subgraph of $f(x) + \epsilon/(1 + x^2)$ for $\epsilon > 0$ small.)

5. ★ Show that C is closed iff

$$d((x, y), C) = 0 \quad \Longleftrightarrow \quad (x, y) \in C.$$

If C is closed and $G_n = \{(x, y) : d((x, y), C) < 1/n\}$, then, G_n is open and $C = \bigcap_{n=1}^{\infty} G_n$. Thus, every closed set is interopen.

6. Let $A \subset \mathbb{R}^2$ be arbitrary. Use the definition of area (A) to show: For all $\epsilon > 0$, there is an open superset G of A satisfying $\text{area}(G) \leq \text{area}(A) + \epsilon$. Conclude

that

$$\text{area}(A) = \inf\{\text{area}(G) : A \subset G, G \text{ open}\}.$$

7. Let $A \subset \mathbb{R}^2$ be arbitrary. Show that there is an interopen set I containing A and having the same area as A (use Exercise 6).

8. If (M_n) is a sequence of measurable sets, then, $\bigcup_{n=1}^{\infty} M_n$ is measurable.

9. ★ The Cantor set is closed.

10. Show that $D'_n \supset D$.

11. Derive (5.13g) and (5.13h).

12. If A and B are disjoint and A is measurable, then, $\text{area}(A \cup B) = \text{area}(A) + \text{area}(B)$.

13. ★ If (A_n) is a sequence of disjoint measurable sets, then,

$$\text{area}\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \text{area}(A_n).$$

14. If A and B are measurable, then, $\text{area}(A \cup B) = \text{area}(A) + \text{area}(B) - \text{area}(A \cap B)$.

15. ★ Let A, B, C, D be measurable subsets of \mathbb{R}^2 . Obtain expressions for $\text{area}(A \cup B \cup C)$ and $\text{area}(A \cup B \cup C \cup D)$ akin to the result in the previous Exercise.

16. ★ Show that M is measurable iff, for all $\epsilon > 0$, there is an open superset G of M , such that $\text{area}(G \setminus M) \leq \epsilon$.

17. ★ Let $A \subset \mathbb{R}^2$ be measurable. If $\text{area}(A) > 0$, there is an $\epsilon > 0$, such that $\text{area}(A \cap A') > 0$ for all translates $A' = A + (a, b)$ of A with $|a| < \epsilon$ and $|b| < \epsilon$. (Start with A a rectangle, and use Exercise 15 of §4.2.)

18. ★ If $A \subset \mathbb{R}^2$ is measurable and $\text{area}(A) > 0$, let

$$A - A = \{(x - x', y - y') : (x, y) \text{ and } (x', y') \in A\}$$

be the *set of differences*. Note that $A - A$ contains the origin. Then, for some $\epsilon > 0$, $A - A$ must contain the open rectangle $Q_\epsilon = (-\epsilon, \epsilon) \times (-\epsilon, \epsilon)$. (Use Exercise 17.)

5

Applications

§5.1 Euler's Gamma Function

In this section, we derive the beautiful formula

$$(1.1) \quad \int_0^1 \frac{dx}{x^x} = \sum_{n=1}^{\infty} \frac{1}{n^n} \\ = \frac{1}{1^1} + \frac{1}{2^2} + \frac{1}{3^3} + \dots$$

Along the way we will meet Euler's gamma function and the monotone convergence theorem, both of which play roles in subsequent sections.

The gamma function is defined by

$$\Gamma(x) = \int_0^{\infty} e^{-t} t^{x-1} dt, \quad x > 0.$$

Clearly $\Gamma(x)$ is positive for $x > 0$. Below we see that the gamma function is finite, and, in the next section, we see that it is continuous. Since

$$(1.2) \quad \int_0^{\infty} e^{-at} dt = \frac{1}{-a} e^{-at} \Big|_0^{\infty} = -\frac{1}{a} (e^{-a\infty} - e^{-a0}) = \frac{1}{a}, \quad a > 0,$$

we have $\Gamma(1) = 1$. Below, we use the convention $0! = 1$.

Theorem. *The gamma function $\Gamma(x)$ is positive, finite, and $\Gamma(x+1) = x\Gamma(x)$ for $x > 0$. Moreover, $\Gamma(n) = (n-1)!$ for $n \geq 1$.*

To derive the first identity, use integration by parts with $u = t^x$, $dv = e^{-t} dt$. Then, $v = -e^{-t}$, and $du = xt^{x-1} dt$. Hence, we obtain the following equality between primitives:

$$\int e^{-t} t^x dt = -e^{-t} t^x + x \int e^{-t} t^{x-1} dt.$$

Since $e^{-t}t^x$ vanishes at $t = 0$ and $t = \infty$ for $x > 0$ fixed, and the integrands are positive, by the fundamental theorem, we obtain

$$\begin{aligned}\Gamma(x+1) &= \int_0^\infty e^{-t}t^x dt \\ &= -e^{-t}t^x \Big|_0^\infty + x \int_0^\infty e^{-t}t^{x-1} dt \\ &= 0 + x\Gamma(x) = x\Gamma(x).\end{aligned}$$

Note that this identity is true whether or not $\Gamma(x)$ is finite. We derive $\Gamma(n) = (n-1)!$ by induction. The statement is true for $n = 1$ since $\Gamma(1) = 1$ from above. Assuming the statement is true for n , $\Gamma(n+1) = n\Gamma(n) = n(n-1)! = n!$. Hence, the statement is true for all $n \geq 1$. Now, we show that $\Gamma(x)$ is finite for all $x > 0$. Since the integral $\int_0^1 e^{-t}t^{x-1} dt \leq \int_0^1 t^{x-1} dt = 1/x$ is finite for $x > 0$, it is enough to verify integrability of $e^{-t}t^{x-1}$ over $(1, \infty)$. Over this interval, $e^{-t}t^{x-1}$ increases with x , hence, $\int_1^\infty e^{-t}t^{x-1} dt \leq \int_1^\infty e^{-t}t^{n-1} dt \leq \Gamma(n)$ for any natural $n \geq x$. But we already know that $\Gamma(n) = (n-1)! < \infty$, hence, the result. \square

Because of this result, we *define* $x! = \Gamma(x+1)$ for $x > -1$. For example, in Exercise 1 of §5.4, we obtain $(1/2)! = \sqrt{\pi}/2$.

We already know (linearity §4.4) that the integral of a finite sum of continuous functions is the sum of their integrals. To obtain linearity for infinite sums, we first derive the following.

Monotone Convergence Theorem (For Integrals). *Let $0 \leq f_1 \leq f_2 \leq f_3 \leq \dots$ be an increasing sequence of nonnegative functions, all defined on an interval (a, b) . If*

$$\lim_{n \nearrow \infty} f_n(x) = f(x), \quad a < x < b,$$

then,

$$\lim_{n \nearrow \infty} \int_a^b f_n(x) dx = \int_a^b \lim_{n \nearrow \infty} f_n(x) dx = \int_a^b f(x) dx.$$

To see this, let G_n denote the subgraph of f_n over (a, b) , G the subgraph of f over (a, b) . Then, $G_n \subset G_{n+1}$ since $y < f_n(x)$ implies $y < f_{n+1}(x)$. Moreover, $y < f(x)$ iff $y < f_n(x)$ for some $n \geq 1$, hence, $G = \bigcup_{n=1}^\infty G_n$. The result now follows from the method of exhaustion and the definition of the integral of a nonnegative function. \square

We caution that this result may be false when the sequence (f_n) is not increasing (Exercise 1). Nevertheless, one can still obtain roughly half this result for any sequence (f_n) of nonnegative functions (Exercise 2).

As an immediate application of the above, since $\Gamma(x) = \Gamma(x+1)/x$ for $x > 0$ and $\Gamma(1) = 1$, $\Gamma(0+) = \infty$. Also, for $x \geq n$, $\Gamma(x) \geq \int_1^\infty e^{-t}t^{x-1} dt \geq \int_1^\infty e^{-t}t^{n-1} dt \geq (n-1)! - 1/n$. Hence, $\Gamma(\infty) = \infty$. In the next section, we show that Γ is continuous whereas, in Exercise 7, we show that Γ is convex. Later

(§5.4), we show that Γ is strictly convex. Putting all of this together, we conclude that Γ has exactly one global positive minimum and the graph for $x > 0$ is as in Figure 5.1. Later (§5.8), we will extend the domain of Γ to negative reals.




FIGURE 5.1. The gamma function.

Now, we can derive linearity for infinite positive series of functions. Let (f_n) be a sequence of nonnegative functions. Then, the sequence of partial sums $s_n = f_1 + \cdots + f_n$, $n \geq 1$, is an increasing sequence, hence, the monotone convergence theorem applies. Since $s_n \nearrow \sum_{n=1}^{\infty} f_n$, we obtain the following.

Theorem (Summation Under the Integral Sign — Positive Case). *If (f_n) is a sequence of nonnegative continuous functions on (a, b) , then,*

$$\int_a^b \left[\sum_{n=1}^{\infty} f_n(x) \right] dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx.$$

For alternating series, we need a different version of this result (next section).

Since the functions are continuous, use linearity and the monotone convergence theorem to obtain

$$\begin{aligned} \int_a^b \left[\sum_{n=1}^{\infty} f_n(x) \right] dx &= \int_a^b \lim_{n \nearrow \infty} \left[\sum_{k=1}^n f_k(x) \right] dx \\ &= \lim_{n \nearrow \infty} \int_a^b \left[\sum_{k=1}^n f_k(x) \right] dx \\ &= \lim_{n \nearrow \infty} \sum_{k=1}^n \int_a^b f_k(x) dx \end{aligned}$$

$$= \sum_{n=1}^{\infty} \int_a^b f_n(x) dx. \quad \square$$

Now, we derive (1.1). To see this, we use the substitution $x = e^{-t}$ (Exercise 5 of §4.4), the exponential series, the previous theorem, shifting the index n by one, the substitution $nt = s$ (dilation invariance), and the property $\Gamma(n) = (n-1)!$:

$$\begin{aligned} \int_0^1 x^{-x} dx &= \int_0^{\infty} (e^{-t})^{-e^{-t}} e^{-t} dt = \int_0^{\infty} e^{te^{-t}} e^{-t} dt \\ &= \int_0^{\infty} \left[\sum_{n=0}^{\infty} \frac{1}{n!} t^n e^{-nt} \right] e^{-t} dt \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \int_0^{\infty} t^n e^{-(n+1)t} dt \\ &= \sum_{n=1}^{\infty} \frac{1}{(n-1)!} \int_0^{\infty} e^{-nt} t^{n-1} dt \\ &= \sum_{n=1}^{\infty} \frac{1}{(n-1)!} n^{-n} \int_0^{\infty} e^{-s} s^{n-1} ds \\ &= \sum_{n=1}^{\infty} \frac{1}{(n-1)!} n^{-n} \Gamma(n) = \sum_{n=1}^{\infty} n^{-n}. \quad \square \end{aligned}$$

We end with a special case of the monotone convergence theorem.

Monotone Convergence Theorem (For Series). *Let $(a_{nj}), n \geq 1$, be a sequence of sequences, and let (a_j) be a given sequence. Suppose that $0 \leq a_{1j} \leq a_{2j} \leq a_{3j} \leq \dots$ for all $j \geq 1$. If*

$$\lim_{n \nearrow \infty} a_{nj} = a_j, \quad j \geq 1,$$

then,

$$\lim_{n \nearrow \infty} \sum_{j=1}^{\infty} a_{nj} = \sum_{j=1}^{\infty} \lim_{n \nearrow \infty} a_{nj} = \sum_{j=1}^{\infty} a_j.$$

To see this, define piecewise constant functions $f_n(x) = a_{nj}$, $j-1 < x \leq j$, $j \geq 1$, and $f(x) = a_j$, $j-1 < x \leq j$, $j \geq 1$. Then, (f_n) is nonnegative on $(0, \infty)$ and increasing to f . Now, apply the monotone convergence theorem for integrals, and use Exercise 6 of §4.3. \square

Using this theorem, one can derive an analog of summation under the integral sign involving series ("summation under the summation sign"), rather than integrals. But we already did this in §1.7.

Exercises 5.1

1. Find a sequence $f_1 \geq f_2 \geq f_3 \geq \cdots \geq 0$ of nonnegative functions, such that $f_n(x) \rightarrow 0$ for all $x \in \mathbb{R}$, but $\int_{-\infty}^{\infty} f_n(x) dx = \infty$ for all $n \geq 1$. This shows that the monotone convergence theorem is false for decreasing sequences.
2. (Fatou's Lemma) Let f_n , $n \geq 1$, be nonnegative functions, all defined on (a, b) , and suppose that $f_n(x) \rightarrow f(x)$ for all x in (a, b) . Then, the lower limit of the sequence $\left(\int_a^b f_n(x) dx\right)$ is greater or equal to $\int_a^b f(x) dx$. (For each x , let $(g_n(x))$ equal the lower sequence (§1.5) of the sequence $(f_n(x))$.)
3. Let $f_0(x) = 1 - x^2$ for $|x| < 1$ and $f_0(x) = 0$ for $|x| \geq 1$, and let $f_n(x) = f_0(x - n)$ for $-\infty < x < \infty$ and $n \geq 1$. Compute $f(x) = \lim_{n \nearrow \infty} f_n(x)$, $-\infty < x < \infty$, $\int_{-\infty}^{\infty} f_n(x) dx$, $n \geq 1$, and $\int_{-\infty}^{\infty} f(x) dx$ (Figure 5.2). Conclude that, for this example, the inequality in Fatou's lemma is strict.




FIGURE 5.2. Exercise 3.

4. Show that

$$\Gamma(x) = \lim_{n \nearrow \infty} \int_0^n \left(1 - \frac{t}{n}\right)^n t^{x-1} dt, \quad x > 0.$$

(Use Exercise 4 of §3.2.)

5. Use substitution $t = ns$, and integrate by parts to get

$$\int_0^n \left(1 - \frac{t}{n}\right)^n t^{x-1} dt = \frac{n^x n!}{x(x+1) \cdots (x+n)}, \quad x > 0,$$

for $n \geq 1$. Conclude that

$$(1.3) \quad \Gamma(x) = \lim_{n \nearrow \infty} \frac{n^x n!}{x(x+1) \cdots (x+n)}, \quad x > 0.$$

6. ★ We say that a function $f : (a, b) \rightarrow \mathbb{R}^+$ is *log-convex* if $\log f$ is convex (§3.3). Show that the right side of (1.3) is log-convex on $(0, \infty)$. Suppose that $f_n : (a, b) \rightarrow \mathbb{R}$, $n \geq 1$, is a sequence of convex functions, and $f_n(x) \rightarrow f(x)$, as $n \nearrow \infty$, for all x in (a, b) . Show that f is convex on (a, b) . Conclude that the gamma function is log-convex on $(0, \infty)$.

7. ★ Show that Γ is convex on $(0, \infty)$. (Consider $\Gamma = \exp(\log \Gamma)$ and remember that e^x is convex.)

8. Let $s_n(t)$ denote the n th partial sum of

$$\frac{1}{e^t - 1} = e^{-t} + e^{-2t} + e^{-3t} + \dots, \quad t > 0.$$

Use $s_n(t)$ to derive

$$\int_0^\infty \frac{t^{x-1}}{e^t - 1} dt = \Gamma(x)\zeta(x), \quad x > 1,$$

where $\zeta(x) = \sum_{n=1}^\infty n^{-x}$, $x > 1$.

9. Let

$$\psi(t) = \sum_{n=1}^\infty e^{-n^2\pi t}, \quad t > 0.$$

Show that

$$\int_0^\infty \psi(t)t^{x/2-1} dt = \pi^{-x/2}\Gamma(x/2)\zeta(x), \quad x > 1,$$

where ζ is as in the previous exercise.

10. Show that $\int_0^1 t^{x-1}(-\log t)^{n-1} dt = \Gamma(n)/x^n$ for $x > 0$ and $n \geq 1$ (Exercise 5 of §4.4).

11. Show that

$$\int_0^\infty e^{-t} t^{x-1} |\log t|^{n-1} dt \leq \frac{\Gamma(n)}{x^n} + \Gamma(x+n-1)$$

for $x > 0$ and $n \geq 1$. (Break up the integral into $\int_0^1 + \int_1^\infty$ and use $\log t \leq t$ for $t \geq 1$.)

12. Use the monotone convergence theorem for series to compute $\zeta(1+)$ and $\psi(0+)$.

13. With $\tau(t) = t/(1 - e^{-t})$, show that

$$\int_0^\infty e^{-xt} \tau(t) dt = \sum_{n=0}^\infty \frac{1}{(x+n)^2}, \quad x > 0.$$

(Compare with Exercise 8.)

14. Use the monotone convergence theorem to derive continuity at the endpoints (§4.3): If $f : (a, b) \rightarrow \mathbb{R}$ is nonnegative and $a_n \searrow a$, $b_n \nearrow b$, then, $\int_{a_n}^{b_n} f(x) dx \rightarrow \int_a^b f(x) dx$.

§5.2 The Number π

In this section, we discuss several formulas for π , namely,

- the *Leibnitz series*

$$(2.1) \quad \frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots,$$

- the *Wallis product*

$$(2.2) \quad \frac{\pi}{2} = \lim_{n \rightarrow \infty} \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot \dots \cdot 2n \cdot 2n}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot \dots \cdot (2n-1) \cdot (2n+1)},$$

- the *Vieta formula*

$$(2.3) \quad \frac{2}{\pi} = \sqrt{\frac{1}{2}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}}} \cdot \dots,$$

- the *continued fraction expansion*

$$(2.4) \quad \frac{\pi}{4} = \frac{1}{1 + \frac{1}{2 + \frac{9}{2 + \frac{25}{2 + \frac{49}{2 + \dots}}}}},$$

- the *Bailey–Borwein–Plouffe series*

$$(2.5) \quad \pi = \sum_{n=0}^{\infty} \frac{1}{16^n} \left(\frac{4}{8n+1} - \frac{2}{8n+4} - \frac{1}{8n+5} - \frac{1}{8n+6} \right).$$

Along the way, we will meet the dominated convergence theorem, and we also compute the Laplace transform of the Bessel function of order zero.

It is one thing to derive these remarkable formulas and quite another to *discover* them. We begin by rederiving the Leibnitz series for π by an alternate method to that in §3.6.

Start with the power series expansion

$$(2.6) \quad \frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \dots, \quad 0 < x < 1.$$

We seek to integrate (2.6), term by term, as in §4.4. Since $\arctan 1 = \pi/4$ and $\arctan x$ is a primitive of $1/(1+x^2)$, we seek to integrate (2.6) over the interval $(0, 1)$. However, since the radius of convergence of (2.6) is 1, the result in §4.4 is not applicable. On the other hand, the theorem in §5.1 allows us to integrate, term by term, any series of nonnegative continuous functions. Since (2.6) is alternating, again this is not applicable.

If we let $s_n(x)$ denote the n th partial sum in (2.6), then, by the Leibnitz test (§1.7),

$$(2.7) \quad 0 < s_n(x) < 1, \quad 0 < x < 1, n \geq 1.$$

It turns out that (2.7) allows us to integrate (2.6) over the interval $(0,1)$, term by term. This is captured in the following theorem.

Dominated Convergence Theorem (for Integrals). *Let $f_n : (a, b) \rightarrow \mathbb{R}, n \geq 1$, be a sequence of continuous functions, and let $f : (a, b) \rightarrow \mathbb{R}$ be continuous. Suppose that there is an integrable positive continuous function $g : (a, b) \rightarrow \mathbb{R}$ satisfying $|f_n(x)| \leq g(x)$ for all x in (a, b) and all $n \geq 1$. If*

$$\lim_{n \nearrow \infty} f_n(x) = f(x), \quad a < x < b,$$

then,

$$(2.8) \quad \lim_{n \nearrow \infty} \int_a^b f_n(x) dx = \int_a^b \lim_{n \nearrow \infty} f_n(x) dx = \int_a^b f(x) dx.$$

Note that (2.8) says we can switch the limit and the integral, exactly as in the monotone convergence theorem. This theorem takes its name from the hypothesis $|f_n(x)| \leq g(x), a < x < b$, which is read *f_n is dominated by g over (a, b)* . The point of this hypothesis is the existence of a *single* continuous integrable g that dominates *all* the f_n 's.

The two results, the monotone and the dominated convergence theorems, are used throughout mathematics to justify the interchange of integrals and limits. Which theorem is applied when depends on which hypothesis is applicable to the problem at hand. When trigonometric or more general oscillatory functions are involved, the monotone convergence theorem is not applicable. Often, in these cases, it is the dominated convergence theorem that saves the day.

When $(a, b) = (0, \infty)$ and the functions $f_n, n \geq 1, f, g$, are piecewise constant, the dominated convergence theorem reduces to a theorem about series, which we discuss at the end of the section. Also one can allow the interval (a_n, b_n) to vary with $n \geq 1$ (Exercise 13). We defer the derivation of the dominated convergence theorem to the end of the section.

Going back to the derivation of (2.1), since the n th partial sum s_n converges to $f(x) = 1/(1+x^2)$ and $|s_n(x)| \leq 1$ by (2.7), we can choose $g(x) = 1$ which is integrable on $(0, 1)$. Hence, applying the fundamental theorem and the dominated convergence theorem yields

$$\begin{aligned} \frac{\pi}{4} &= \arctan 1 - \arctan 0 = \int_0^1 \frac{1}{1+x^2} dx = \int_0^1 \left[\sum_{n=0}^{\infty} (-1)^n x^{2n} \right] dx \\ &= \int_0^1 \lim_{N \nearrow \infty} s_N(x) dx = \lim_{N \nearrow \infty} \int_0^1 s_N(x) dx \\ &= \lim_{N \nearrow \infty} \sum_{n=0}^N (-1)^n \int_0^1 x^{2n} dx = \sum_{n=0}^{\infty} (-1)^n \int_0^1 x^{2n} dx \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots \end{aligned}$$

This completes the derivation of (2.1).

The idea behind this derivation of (2.1) can be carried out more generally.

Theorem (Summation Under the Integral Sign — Alternating Case). *Let $f_n : (a, b) \rightarrow \mathbf{R}, n \geq 1$, be a decreasing sequence of continuous positive functions on (a, b) , and suppose that f_1 is integrable. If $\sum_{n=1}^{\infty} (-1)^{n-1} f_n$ is continuous on (a, b) , then,*

$$\int_a^b \left[\sum_{n=1}^{\infty} (-1)^{n-1} f_n(x) \right] dx = \sum_{n=1}^{\infty} (-1)^{n-1} \int_a^b f_n(x) dx.$$

To derive this, we need only note that the n th partial sum $s_n(x)$ is nonnegative and no greater than $g(x) = f_1(x)$, which is continuous and integrable. Hence, we may apply the dominated convergence theorem, as above, to the sequence (s_n) of partial sums. \square

For example, using this theorem to integrate the geometric series $1/(1+x) = 1 - x + x^2 - x^3 + \dots$ over $(0, 1)$, we obtain

$$\log 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

Now, we discuss the general case.

Theorem (Summation Under the Integral Sign — Absolute Case). *Let $f_n : (a, b) \rightarrow \mathbf{R}, n \geq 1$, be a sequence of continuous functions, and suppose that there is an integrable, positive, continuous function $g : (a, b) \rightarrow \mathbf{R}$ satisfying $\sum_{n=1}^{\infty} |f_n(x)| \leq g(x)$ for all x in (a, b) and all $n \geq 1$. If $\sum_{n=1}^{\infty} f_n(x)$ is continuous on (a, b) , then,*

$$\int_a^b \left[\sum_{n=1}^{\infty} f_n(x) \right] dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx.$$

To derive this, we need only note that $|s_N(x)| \leq |f_1(x)| + \dots + |f_N(x)| \leq g(x)$, which is continuous and integrable. Hence, we may apply the dominated convergence theorem, as above, to the sequence (s_n) of partial sums. \square

The *Bessel function of order zero* is defined by

$$J_0(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{4^n (n!)^2}, \quad -\infty < x < \infty.$$

To check the convergence, rewrite the series using Exercise 13 of §3.4 obtaining

$$J_0(x) = \sum_{n=0}^{\infty} \binom{-1/2}{n} \frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$$

Now, use the definition of $\binom{\nu}{n}$ (§3.4) to check the inequality $|\binom{-1/2}{n}| \leq 1$ for all $n \geq 0$. Hence,

$$(2.9) \quad |J_0(x)| \leq \sum_{n=0}^{\infty} \left| \binom{-1/2}{n} \frac{x^{2n}}{(2n)!} \right| \leq \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} \leq e^{|x|}, \quad -\infty < x < \infty.$$

This shows that the series J_0 converges absolutely for all x real. Since J_0 is a convergent power series, J_0 is a smooth function on \mathbb{R} . We wish to use summation under the integral sign to obtain

$$(2.10) \quad \int_0^{\infty} e^{-sx} J_0(x) dx = \frac{1}{\sqrt{1+s^2}}, \quad s > 1.$$

The left side of (2.10), by definition, is the *Laplace transform* of J_0 . Thus, (2.10) exhibits the Laplace transform of the Bessel function J_0 . In Exercise 1, you are asked to derive the Laplace transform of $\sin x/x$.

To obtain (2.10), fix $s > 1$, and set $f_n(x) = e^{-sx} \binom{-1/2}{n} x^{2n}/(2n)!$, $n \geq 0$. Then, by (2.9), we may apply summation under the integral sign with $g(x) = e^{-sx} e^x$, $x > 0$, which is positive, continuous, and integrable (since $s > 1$). Hence,

$$\int_0^{\infty} e^{-sx} J_0(x) dx = \sum_{n=0}^{\infty} \binom{-1/2}{n} \frac{1}{(2n)!} \int_0^{\infty} e^{-sx} x^{2n} dx.$$

Inserting the substitution $x = t/s$, $dx = dt/s$, and recalling Newton's generalization of the binomial theorem (§3.4) yields

$$\begin{aligned} \int_0^{\infty} e^{-sx} J_0(x) dx &= \sum_{n=0}^{\infty} \binom{-1/2}{n} \frac{1}{(2n)! s^{2n+1}} \int_0^{\infty} e^{-t} t^{2n} dt \\ &= \sum_{n=0}^{\infty} \binom{-1/2}{n} \frac{\Gamma(2n+1)}{(2n)! s^{2n+1}} \\ &= \frac{1}{s} \sum_{n=0}^{\infty} \binom{-1/2}{n} \left(\frac{1}{s^2}\right)^n = \frac{1}{s} \cdot \frac{1}{\sqrt{1+(1/s)^2}} = \frac{1}{\sqrt{1+s^2}}. \end{aligned}$$

This establishes (2.10).

Now, let

$$F(x) = \int_0^{\infty} \frac{\sin(xt)}{1+t^2} dt, \quad -\infty < x < \infty.$$

We use the dominated convergence theorem to show that F is continuous on \mathbb{R} . To show this, fix $x \in \mathbb{R}$, and let $x_n \rightarrow x$; we have to show that $F(x_n) \rightarrow F(x)$. Set $f_n(t) = \sin(x_n t)/(1+t^2)$, $f(t) = \sin(xt)/(1+t^2)$, and $g(t) = 1/(1+t^2)$. Then, $f_n(t)$, $n \geq 1$, $f(t)$, and $g(t)$ are continuous, all the $f_n(t)$'s are dominated by $g(t)$ over $(0, \infty)$, $f_n(t) \rightarrow f(t)$ for all $t > 0$, and $g(t)$ is integrable over $(0, \infty)$ since $\int_0^{\infty} g(t) dt = \pi/2$. Hence, the theorem applies, and

$$\lim_{n \nearrow \infty} F(x_n) = \lim_{n \nearrow \infty} \int_0^{\infty} \frac{\sin(x_n t)}{1+t^2} dt$$

$$\begin{aligned}
&= \int_0^\infty \lim_{n \nearrow \infty} \frac{\sin(x_n t)}{1+t^2} dt \\
&= \int_0^\infty \frac{\sin(xt)}{1+t^2} dt = F(x).
\end{aligned}$$

This establishes the continuity of F .

Similarly, one can establish the continuity of the gamma function on $(0, \infty)$. To this end, choose $0 < a < x < b < \infty$, and let $x_n \rightarrow x$ with $a < x_n < b$. We have to show $\Gamma(x_n) \rightarrow \Gamma(x)$. Now, $f_n(t) = e^{-t} t^{x_n-1}$ satisfies

$$|f_n(t)| \leq \begin{cases} e^{-t} t^{b-1}, & 1 \leq t < \infty, \\ e^{-t} t^{a-1}, & 0 < t \leq 1. \end{cases}$$

If we call the right side of this inequality $g(t)$, we see that $f_n(t)$, $n \geq 1$, are all dominated by $g(t)$ over $(0, \infty)$. Moreover, $g(t)$ is continuous (especially at $t = 1$) and integrable over $(0, \infty)$, since $\int_0^\infty g(t) dt \leq \Gamma(a) + \Gamma(b)$. Also, the functions $f_n(t)$, $n \geq 1$, and $f(t) = e^{-t} t^{x-1}$ are continuous, and $f_n(t) \rightarrow f(t)$ for all $t > 0$. Thus, the dominated convergence theorem applies. Hence,

$$\begin{aligned}
\lim_{n \nearrow \infty} \Gamma(x_n) &= \lim_{n \nearrow \infty} \int_0^\infty e^{-t} t^{x_n-1} dt \\
&= \int_0^\infty \lim_{n \nearrow \infty} e^{-t} t^{x_n-1} dt = \int_0^\infty e^{-t} t^{x-1} dt = \Gamma(x).
\end{aligned}$$

Hence, Γ is continuous on (a, b) . Since $0 < a < b$ are arbitrary, Γ is continuous on $(0, \infty)$.

Next, we derive Wallis' product (2.2). Begin with integrating by parts to obtain

$$(2.11) \quad \int \sin^n x \, dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x \, dx, \quad n \geq 2.$$

Evaluating at 0 and $\pi/2$ yields

$$\int_0^{\pi/2} \sin^n x \, dx = \frac{n-1}{n} \int_0^{\pi/2} \sin^{n-2} x \, dx, \quad n \geq 2.$$

Since $\int_0^{\pi/2} \sin^0 x \, dx = \pi/2$ and $\int_0^{\pi/2} \sin^1 x \, dx = 1$, by the last equation and induction,

$$I_{2n} = \int_0^{\pi/2} \sin^{2n} x \, dx = \frac{(2n-1) \cdot (2n-3) \cdot \dots \cdot 1}{2n \cdot (2n-2) \cdot \dots \cdot 2} \cdot \frac{\pi}{2},$$

and

$$I_{2n+1} = \int_0^{\pi/2} \sin^{2n+1} x \, dx = \frac{2n \cdot (2n-2) \cdot \dots \cdot 2}{(2n+1) \cdot (2n-1) \cdot \dots \cdot 3} \cdot 1,$$

for $n \geq 1$. Since $0 < \sin x < 1$ on $(0, \pi/2)$, the integrals I_n are decreasing in n . But, by the formula for I_n with n odd,

$$1 \leq \frac{I_{2n-1}}{I_{2n+1}} \leq 1 + \frac{1}{2n}, \quad n \geq 1.$$

Thus,

$$1 \leq \frac{I_{2n}}{I_{2n+1}} \leq \frac{I_{2n-1}}{I_{2n+1}} \leq 1 + \frac{1}{2n}, \quad n \geq 1,$$

or $I_{2n}/I_{2n+1} \rightarrow 1$, as $n \nearrow \infty$. Since

$$\frac{I_{2n}}{I_{2n+1}} = \frac{(2n+1) \cdot (2n-1) \cdot (2n-1) \cdot \dots \cdot 3 \cdot 3 \cdot 1}{2n \cdot 2n \cdot (2n-2) \cdot \dots \cdot 4 \cdot 2 \cdot 2} \cdot \frac{\pi}{2},$$

we obtain (2.2).

A derivation of Vieta's formula (2.3) starts with the identity

$$(2.12) \quad \frac{\sin \theta}{2^n \sin(\theta/2^n)} = \cos\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{2^2}\right) \dots \cos\left(\frac{\theta}{2^n}\right)$$

which follows by multiplying both sides by $\sin(\theta/2^n)$ and using the double-angle formula $\sin(2x) = 2 \sin x \cos x$ repeatedly. Now insert in (2.12) $\theta = \pi/2$, and use, repeatedly, the formula $\cos(\theta/2) = \sqrt{(1 + \cos \theta)/2}$. This yields

$$\begin{aligned} \frac{2}{\pi} \cdot \frac{\pi/2^{n+1}}{\sin(\pi/2^{n+1})} &= \sqrt{\frac{1}{2}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}}} \cdot \dots \\ &\quad \dots \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \dots}}}, \end{aligned}$$

where the last (n th) factor involves n square roots. Letting $n \nearrow \infty$ yields (2.3) since $\sin x/x \rightarrow 1$ as $x \rightarrow 0$.

To derive the continued fraction expansion (2.4), first, we must understand what it means. To this end, introduce the *convergents*

$$(2.13) \quad c_n = \frac{1}{1 + \frac{1}{2 + \frac{9}{2 + \frac{25}{2 + \frac{49}{\ddots \frac{(2n-1)^2}{2 + \frac{1}{2}}}}}}},$$

Then, we take (2.4) to mean that the sequence (c_n) converges to $\pi/4$. To derive this, it is enough to show that¹⁶ c_n equals the n th partial sum

$$s_n = 1 - \frac{1}{3} + \frac{1}{5} - \dots \pm \frac{1}{2n+1}$$

¹⁶Following Euler.

of the Leibnitz series (2.1) for all $n \geq 1$.

Given reals a_1, \dots, a_n , let

$$s_n^* = a_1 + a_1 a_2 + a_1 a_2 a_3 + \dots + a_1 a_2 \dots a_n.$$

Then, $s_n^* = s_n^*(a_1, \dots, a_n)$ is a function of the n variables a_1, \dots, a_n . Later, we will make a judicious choice of a_1, \dots, a_n . Note that

$$a_1 + a_1 s_n^*(a_2, \dots, a_{n+1}) = s_{n+1}^*(a_1, a_2, \dots, a_{n+1}).$$

Let $f(x, y) = x/(1 + x - y)$, and let

$$c_1^*(a_1) = f(a_1, 0) = \frac{a_1}{1 + a_1},$$

$$c_2^*(a_1, a_2) = f(a_1, f(a_2, 0)) = \frac{a_1}{1 + a_1 - \frac{a_2}{1 + a_2}},$$

$$c_3^*(a_1, a_2, a_3) = f(a_1, f(a_2, f(a_3, 0))) = \frac{a_1}{1 + a_1 - \frac{a_2}{1 + a_1 - \frac{a_3}{1 + a_2 - \frac{a_3}{1 + a_3}}}},$$

and so on. More systematically, define $c_n^*(a_1, \dots, a_n)$ inductively by setting $c_1^*(a_1) = a_1/(1 + a_1)$ and

$$c_{n+1}^*(a_1, a_2, \dots, a_{n+1}) = \frac{a_1}{1 + a_1 - c_n^*(a_2, \dots, a_{n+1})}, \quad n \geq 1.$$

We claim that: $c_n^* = s_n^*/(1 + s_n^*)$ for all $n \geq 1$, and we verify this by induction. Here $c_n^* = c_n^*(a_1, \dots, a_n)$ and $s_n^* = s_n^*(a_1, \dots, a_n)$. Clearly $c_1^* = s_1^*/(1 + s_1^*)$ since $s_1^* = a_1$. Now, assume $c_n^* = s_n^*/(1 + s_n^*)$. Replacing a_1, \dots, a_n by a_2, \dots, a_{n+1} yields

$$c_n^*(a_2, \dots, a_{n+1}) = s_n^*(a_2, \dots, a_{n+1})/[1 + s_n^*(a_2, \dots, a_{n+1})].$$

Then,

$$\begin{aligned} c_{n+1}^*(a_1, a_2, \dots, a_{n+1}) &= \frac{a_1}{1 + a_1 - c_n^*(a_2, \dots, a_{n+1})} \\ &= \frac{a_1}{1 + a_1 - \frac{s_n^*(a_2, \dots, a_{n+1})}{1 + s_n^*(a_2, \dots, a_{n+1})}} \\ &= \frac{s_{n+1}^*(a_1, a_2, \dots, a_{n+1})}{1 + s_{n+1}^*(a_1, a_2, \dots, a_{n+1})}. \end{aligned}$$

Thus, $c_{n+1}^* = s_{n+1}^*/(1 + s_{n+1}^*)$. Hence, by induction, the claim is true.

Now, choose

$$a_1 = -\frac{1}{3}, a_2 = -\frac{3}{5}, a_3 = -\frac{5}{7}, \dots, a_n = -\frac{2n-1}{2n+1}.$$

Then,

$$s_n^* = -\frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots \pm \frac{1}{2n+1} = s_n - 1,$$

and

$$(2.14) \quad c_n^* = \frac{-1/3}{1 - 1/3 + \frac{3/5}{1 - 3/5 + \frac{5/7}{\ddots \frac{1 - \frac{2n-3}{2n-1} + \frac{(2n-1)/(2n+1)}{1 - (2n-1)/(2n+1)}}}}.$$

Now, multiply the top and bottom of the first fraction by 3, then, the top and bottom of the second fraction by 5, then, the top and bottom of the third fraction by 7, and so on. (2.14) becomes

$$c_n^* = \frac{-1}{2 + \frac{9}{2 + \frac{25}{2 + \frac{49}{\ddots \frac{(2n-1)^2}{2}}}}},$$

which, when compared with (2.13), yields

$$c_n = \frac{1}{1 - c_n^*} = \frac{1}{1 - s_n^*/(1 + s_n^*)} = \frac{1}{1 - (s_n - 1)/s_n} = s_n, \quad n \geq 1.$$

Since $s_n \rightarrow \pi/4$, we conclude that $c_n \rightarrow \pi/4$. This completes the derivation of (2.4).

The series (2.5) is remarkable not only because of its rapid convergence, but because it can be used to compute specific digits in the hexadecimal (base 16, see §1.6) expansion of π , without computing all previous digits (the Bailey–Borwein–Plouffe paper explains this very clearly; see the references).

To obtain (2.5), check that

$$\frac{4\sqrt{2} - 8x^3 - 4\sqrt{2}x^4 - 8x^5}{1 - x^8} = \frac{4\sqrt{2} - 4x}{x^2 - \sqrt{2}x + 1} - \frac{4x}{1 - x^2}$$

using $x^8 - 1 = (x^4 - 1)(x^4 + 1)$ and $x^4 + 1 = (x^2 + \sqrt{2}x + 1)(x^2 - \sqrt{2}x + 1)$. Hence (Exercises 15 and 16 of §3.6),

$$\int \frac{4\sqrt{2} - 8x^3 - 4\sqrt{2}x^4 - 8x^5}{1 - x^8} dx$$

$$= 4 \arctan(\sqrt{2}x - 1) - 2 \log(x^2 - \sqrt{2}x + 1) + 2 \log(1 - x^2).$$

Evaluating at 0 and $1/\sqrt{2}$ yields

$$(2.15) \quad \pi = \int_0^{1/\sqrt{2}} \frac{4\sqrt{2} - 8x^3 - 4\sqrt{2}x^4 - 8x^5}{1 - x^8} dx.$$

To see the equivalence of (2.15) and (2.5), note that

$$\begin{aligned} \int_0^{1/\sqrt{2}} \frac{x^{k-1}}{1 - x^8} dx &= \int_0^{1/\sqrt{2}} \left(\sum_{n=0}^{\infty} x^{k-1+8n} \right) dx \\ &= \sum_{n=0}^{\infty} \int_0^{1/\sqrt{2}} x^{k-1+8n} dx \\ &= \sum_{n=0}^{\infty} \frac{1}{(k+8n)(\sqrt{2})^{k+8n}} = \frac{1}{\sqrt{2}^k} \sum_{n=0}^{\infty} \frac{1}{16^n(k+8n)}. \end{aligned}$$

Now, use this with k equal 1, 4, 5, and 6, and insert the resulting four series in (2.15). You obtain (2.5).

To derive the dominated convergence theorem, we will need Fatou's lemma, which is Exercise 2 of §5.1. This states that, for any sequence $f_n : (a, b) \rightarrow \mathbf{R}$, $n \geq 1$, of nonnegative functions satisfying $f_n(x) \rightarrow f(x)$ for all x in (a, b) , the lower limit of the sequence $\left(\int_a^b f_n(x) dx \right)$ is greater or equal to $\int_a^b f(x) dx$.

Now, we derive the dominated convergence theorem. Let I^* and I_* denote the upper and lower limits of the sequence $(I_n) = \left(\int_a^b f_n(x) dx \right)$, and let $I = \int_a^b f(x) dx$. It is enough to show that

$$(2.16) \quad I \leq I_* \leq I^* \leq I,$$

since this implies the convergence of (I_n) to I .

If f_n , $n \geq 1$, are as given, then, $\pm f_n(x) \leq g(x)$. Hence, $g(x) - f_n(x)$ and $g(x) + f_n(x)$, $n \geq 1$, are nonnegative and converge to $g(x) - f(x)$ and $g(x) + f(x)$, respectively, for all x in (a, b) .

Apply Fatou's lemma to the sequence $(g + f_n)$. Then, the lower limit of the sequence

$$\int_a^b [g(x) + f_n(x)] dx = \int_a^b g(x) dx + \int_a^b f_n(x) dx = \int_a^b g(x) dx + I_n, \quad n \geq 1,$$

is greater or equal to

$$\int_a^b [g(x) + f(x)] dx = \int_a^b g(x) dx + I.$$

In the last two equations, we are justified in using linearity since the functions f , g , and f_n , $n \geq 1$, are all continuous. But the lower limit of the sequence $\left(\int_a^b g(x) dx + I_n \right)$ equals $\int_a^b g(x) dx + I_*$. Subtracting $\int_a^b g(x) dx$, we conclude that $I_* \geq I$, which is half of (2.16).

Now apply Fatou's lemma to the sequence $(g - f_n)$. Then, the lower limit of the sequence

$$\int_a^b [g(x) - f_n(x)] dx = \int_a^b g(x) dx - \int_a^b f_n(x) dx = \int_a^b g(x) dx - I_n, \quad n \geq 1,$$

is greater than or equal to

$$\int_a^b [g(x) - f(x)] dx = \int_a^b g(x) dx - I.$$

Since the lower limit of $(-a_n)$ equals minus the upper limit of (a_n) , the lower limit of the sequence $\left(\int_a^b g(x) dx - I_n\right)$ equals $\int_a^b g(x) dx - I^*$. Subtracting $\int_a^b g(x) dx$, we conclude that $I^* \leq I$, which is the other half of (2.16). \square

Let $f : (a, b) \times (c, d) \rightarrow \mathbb{R}$ be a function of two variables (x, t) , and fix $a < x < b, c < t < d$. We say that f is *continuous*¹⁷ at (x, t) if $x_n \rightarrow x$ and $t_n \rightarrow t$ imply $f(x_n, t_n) \rightarrow f(x, t)$. We say that f is *continuous on* $(a, b) \times (c, d)$ if f is continuous at every (x, t) in $(a, b) \times (c, d)$.

A useful consequence of the dominated convergence theorem is the following.

Theorem (Continuity Under the Integral Sign). *Let $f : (a, b) \times (c, d) \rightarrow \mathbb{R}$ be continuous, and suppose that there is an integrable, positive, continuous $g : (c, d) \rightarrow \mathbb{R}$ satisfying $|f(x, t)| \leq g(t)$ for $a < x < b$ and $c < t < d$. If $F : (a, b) \rightarrow \mathbb{R}$ is defined by*

$$(2.17) \quad F(x) = \int_c^d f(x, t) dt, \quad a < x < b,$$

then, F is continuous.

Note that the domination hypothesis guarantees that F is well defined. To establish this, fix x in (a, b) and let $x_n \rightarrow x$. We have to show that $F(x_n) \rightarrow F(x)$. Let $k_n(t) = f(x_n, t)$, $c < t < d$, $n \geq 1$, and $k(t) = f(x, t)$, $c < t < d$. Then, $k_n(t)$ and $k(t)$ are continuous on (c, d) and $k_n(t) \rightarrow k(t)$ for $c < t < d$. By the domination hypothesis, $|k_n(t)| \leq g(t)$. Thus, the dominated convergence theorem applies, and

$$F(x_n) = \int_c^d k_n(t) dt \rightarrow \int_c^d k(t) dt = F(x). \quad \square$$

For example, the continuity of the gamma function on (a, b) , $0 < a < b < \infty$, follows by choosing, as in the beginning of the section,

$$g(t) = \begin{cases} e^{-t} t^{b-1}, & 1 \leq t < \infty, \\ e^{-t} t^{a-1}, & 0 < t \leq 1. \end{cases}$$

We also have to check that $f(x, t) = e^{-t} t^{x-1}$ is continuous on $(0, \infty) \times (0, \infty)$, so, let $x_n \rightarrow x$ and $t_n \rightarrow t$. Since e^{-t} is continuous, $e^{-t_n} \rightarrow e^{-t}$. Since \log

¹⁷In some texts, this is called *joint continuity*.

is continuous, $\log t_n \rightarrow \log t$. Hence, $(x_n - 1) \log t_n \rightarrow (x - 1) \log t$. Hence, $t_n^{x_n-1} = \exp[(x_n - 1) \log t_n] \rightarrow \exp[(x - 1) \log t] = t^{x-1}$. Hence, $e^{-t_n} t_n^{x_n-1} \rightarrow e^{-t} t^{x-1}$. This shows that f is continuous. Because continuity is established in the same manner in all our examples below, we will usually omit this step.

In fact, continuity under the integral sign is nothing but a packaging of the derivation of continuity of Γ , presented earlier.

Let us go back to the statement of the dominated convergence theorem. When $(a, b) = (0, \infty)$ and the functions (f_n) , f , and g are piecewise constant, the dominated convergence theorem reduces to the following.

Dominated Convergence Theorem (for Series). *Let $(a_{nj}), n \geq 1$, be a sequence of sequences, and let (a_j) be a given sequence. Also suppose that there is a convergent positive series $\sum_{j=1}^{\infty} g_j$ satisfying $|a_{nj}| \leq g_j$ for all $j \geq 1$ and $n \geq 1$. If*

$$\lim_{n \nearrow \infty} a_{nj} = a_j, \quad j \geq 1,$$

then,

$$\lim_{n \nearrow \infty} \sum_{j=1}^{\infty} a_{nj} = \sum_{j=1}^{\infty} \lim_{n \nearrow \infty} a_{nj} = \sum_{j=1}^{\infty} a_j.$$

To see this, for $j - 1 < x \leq j$ set $f_n(x) = a_{nj}$, $n \geq 1$, $f(x) = a_j$, and $g(x) = g_j$, $j = 1, 2, \dots$, and use Exercise 6 of §4.3. Although, strictly speaking, the dominated convergence theorem for integrals is not applicable since, here, the functions $f_n, n \geq 1, f, g$ are not continuous, if we go back to the derivation of the dominated convergence theorem, we see that continuity was used only to assure linearity. But linearity holds (§4.4) just as well for piecewise continuous functions and, in particular, for piecewise constant functions. Thus, (this extension of) the dominated convergence theorem for integrals is applicable, and we get the result for series. Alternatively, instead of extending the dominated convergence theorem to piecewise continuous functions, one can simply repeat the derivation of the dominated convergence theorem with series replacing integrals everywhere. \square

Let us use the dominated convergence theorem for series to show¹⁸

$$(2.18) \quad \lim_{x \rightarrow 1} \left(1 - \frac{1}{2^x} + \frac{1}{3^x} - \frac{1}{4^x} + \dots \right) = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots,$$

which sums to $\log 2$. For this, by the mean value theorem, $(2j - 1)^{-x} - (2j)^{-x} = x(2j - t)^{-x-1}$ for some $0 < t < 1$. Hence, $(2j - 1)^{-x} - (2j)^{-x} \leq 2(2j - 1)^{-3/2}$ when $1/2 < x < 2$. Now, let $x_n \rightarrow 1$ with $1/2 < x_n < 2$, and set $a_{nj} = (2j - 1)^{-x_n} - (2j)^{-x_n}$, $a_j = (2j - 1)^{-1} - (2j)^{-1}$, $g_j = 2(2j - 1)^{-3/2}$ for $j \geq 1, n \geq 1$. Then, $a_{nj} \rightarrow a_j$ and $|a_{nj}| \leq g_j$ for all $j \geq 1$. Hence, the theorem applies, and, since the sequence (x_n) is arbitrary, we obtain (2.18). Note

¹⁸This series converges for $x > 0$ by the Leibnitz test.

how, here, we are not choosing a_{nj} as the individual terms but as pairs of terms, producing an absolutely convergent series out of a conditionally convergent one (cf. the Dirichlet test (§1.7)).

Just as we used the dominated convergence theorem for integrals to obtain continuity under the integral sign, we can use the theorem for series to obtain the following.

Theorem (Continuity Under the Summation Sign). *Let (f_n) be a sequence of continuous functions defined on (a, b) , and suppose that there is a convergent positive series $\sum_{n=1}^{\infty} g_n$ of numbers satisfying $|f_n(x)| \leq g_n$ for $n \geq 1$ and $a < x < b$. If*

$$F(x) = \sum_{n=1}^{\infty} f_n(x), \quad a < x < b,$$

then, $F : (a, b) \rightarrow \mathbf{R}$ is continuous. \square

For example,

$$\zeta(x) = \sum_{n=1}^{\infty} \frac{1}{n^x}$$

is continuous on (a, ∞) for $a > 1$, since $1/n^x \leq 1/n^a$ for $x \geq a$ and $\sum g_n = \sum 1/n^a$ converges. Since $a > 1$ is arbitrary, ζ is continuous on $(1, \infty)$.

Exercises 5.2

1. Use

$$\frac{\sin x}{x} = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \dots$$

to derive the Laplace transform

$$\int_0^{\infty} e^{-sx} \frac{\sin x}{x} dx = \arctan\left(\frac{1}{s}\right), \quad s > 1.$$

2. Suppose that $f_n, n \geq 1$, f , and g are as in the dominated convergence theorem. Show that f is integrable over (a, b) .

3. Show that $x^2 J_0''(x) + x J_0'(x) + x^2 J_0(x) = 0$ for all x .

4. Derive (2.11) by integrating by parts.

5. Show that $\sin x/x = \cos(x/2) \cos(x/4) \cos(x/8) \dots$

6. This is an example where switching the integral and the series changes the answer. Show that

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \int_0^{\infty} e^{-x} x^n dx \neq \int_0^{\infty} e^{-x} \left[\sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right] dx.$$

7. Show that the *Fourier transform*

$$\int_0^\infty \frac{\sin(sx)}{e^x - 1} dx = \sum_{n=1}^\infty \frac{s}{n^2 + s^2}, \quad -\infty < s < \infty.$$

(Compare with Exercise 8 of §5.1.)

8. Show that

$$\int_0^\infty \frac{\sinh(sx)}{e^x - 1} dx = \sum_{n=1}^\infty \frac{s}{n^2 - s^2}, \quad |s| < 1.$$

9. ★ Let (p_n) denote the sequence

$$(4, 0, 0, -2, -1, -1, 0, 0, 4, 0, 0, -2, -1, -1, 0, 0, \dots),$$

where the block $(4, 0, 0, -2, -1, -1, 0, 0)$ repeats forever. Show that the Bailey–Borwein–Plouffe series (2.4) can be rewritten

$$\pi = \sum_{n=1}^\infty \frac{p_n}{16^{[n/8]} \cdot n},$$

where $[x]$ denotes the greatest integer $\leq x$.

10. Show that the ν th *Bessel function*

$$J_\nu(x) = \frac{1}{\pi} \int_0^\pi \cos(\nu t - x \sin t) dt, \quad -\infty < x < \infty,$$

is continuous. Here, ν is any real.

11. Show that $\psi(x) = \sum_{n=1}^\infty e^{-n^2\pi x}$, $x > 0$, is continuous.

12. Let $f_n, f, g : (a, b) \rightarrow \mathbb{R}$ be as in the dominated convergence theorem, and suppose that $a_n \rightarrow a+$ and $b_n \rightarrow b-$. Suppose that we have domination $|f_n(x)| \leq g(x)$ only on (a_n, b_n) , $n \geq 1$. Show that

$$\lim_{n \nearrow \infty} \int_{a_n}^{b_n} f_n(x) dx = \int_a^b f(x) dx.$$

13. Show that the J_0 in the text is the same as the J_ν in Exercise 10 with $\nu = 0$.

14. Use Exercise 18 of §4.4 to show that Euler's constant satisfies

$$\gamma = \lim_{n \nearrow \infty} \left[\int_0^1 \frac{1 - (1 - t/n)^n}{t} dt - \int_1^n \frac{(1 - t/n)^n}{t} dt \right].$$

Use the dominated convergence theorem to conclude that

$$\gamma = \int_0^1 \frac{1 - e^{-t}}{t} dt - \int_1^\infty \frac{e^{-t}}{t} dt.$$

(For the second part, first, use the mean value theorem to show that $0 \leq [1 - (1 - t/n)^n]/t \leq 1$.)

§5.3 Gauss' Arithmetic-Geometric Mean (AGM)

Given $a > b > 0$, their *arithmetic mean* is given by

$$a' = \frac{a+b}{2},$$

and their *geometric mean* by

$$b' = \sqrt{ab}.$$

Since

$$(3.1) \quad a' - b' = \frac{a+b}{2} - \sqrt{ab} = \frac{1}{2} (\sqrt{a} - \sqrt{b})^2 > 0,$$

these equations transform the pair (a, b) , $a > b > 0$, into a pair (a', b') , $a' > b' > 0$. Gauss discovered that iterating this transformation leads to a limit with striking properties.

To begin, since a is the larger of a and b and a' is their arithmetic mean, $a' < a$. Similarly, since b is the smaller of a and b , $b' > b$. Thus, $b < b' < a' < a$.

Set $a_0 = a$ and $b_0 = b$, and define the iteration

$$(3.2i) \quad \begin{aligned} a_{n+1} &= \frac{a_n + b_n}{2}, \\ b_{n+1} &= \sqrt{a_n b_n}, \end{aligned} \quad n \geq 0.$$

By the previous paragraph, for $a > b > 0$, this gives a strictly decreasing sequence (a_n) and a strictly increasing sequence (b_n) with all the a 's greater than all the b 's. Thus, both sequences converge (Figure 5.3) to finite positive limits a_* , b^* with $a_* \geq b^* > 0$.




FIGURE 5.3. The AGM iteration.

Letting $n \nearrow \infty$ in (3.2), we see that $a_* = (a_* + b^*)/2$ which yields $a_* = b^*$. Thus, both sequences converge to a common limit, the *arithmetic-geometric mean (AGM) of (a, b)* , which we denote

$$M(a, b) = \lim_{n \nearrow \infty} a_n = \lim_{n \nearrow \infty} b_n.$$

If (a'_n) , (b'_n) are the sequences associated with $a' = ta$ and $b' = tb$, then, from (3.2i), $a'_n = ta_n$, and $b'_n = tb_n$, $n \geq 1$, $t > 0$. This implies that M is homogeneous in (a, b) ,

$$M(ta, tb) = t \cdot M(a, b), \quad t > 0.$$

The convergence of the sequences (a_n) , (b_n) to the real $M(a, b)$ is *quadratic* in the following sense. The differences $a_n - M(a, b)$ and $M(a, b) - b_n$ are no more than $2c_{n+1}$, where

$$(3.2ii) \quad c_{n+1} = \frac{a_n - b_n}{2}, \quad n \geq 0.$$

By (3.1),

$$0 < c_{n+1} = \frac{1}{4} \left(\sqrt{a_{n-1}} - \sqrt{b_{n-1}} \right)^2 = \frac{c_n^2}{(\sqrt{a_{n-1}} + \sqrt{b_{n-1}})^2} \leq \frac{1}{4b} c_n^2.$$

Iterating the last inequality yields

$$(3.3) \quad 0 < a_n - b_n \leq 8b \left(\frac{a - b}{8b} \right)^{2^n}, \quad n \geq 1.$$

This shows that each additional iteration roughly doubles the number-of-decimal-place agreement, at least if $(a - b)/8b < 1$. For a general pair (a, b) , eventually, $(a_N - b_N)/8b_N < 1$. After this point, we have the rapid convergence (3.3). In (3.16) below, we improve (3.3) from an inequality to an asymptotic equality.

For future reference note that

$$a_n^2 = b_n^2 + c_n^2, \quad n \geq 1.$$

The following remarkable formula is due to Gauss.

Theorem. For $a > b > 0$,

$$\frac{1}{M(a, b)} = \frac{2}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta}}.$$

The proof of this is straightforward, once one knows the answer. Gauss was initially guided to this formula by noting that, for $(a, b) = (1, 1/\sqrt{2})$, both sides agreed to eleven decimal places. We compute $M(1, 1/\sqrt{2})$ explicitly in the next section (see (4.5)).

To derive the formula, call the right side $I(a, b)$. Note that, if $a = b = m$, then, $I(a, b) = I(m, m) = 1/m$. The main step is to show that

$$(3.4) \quad I(a, b) = I\left(\frac{a+b}{2}, \sqrt{ab}\right).$$

To see this, substitute $t = g(\theta) = b \tan \theta$ in $I(a, b)$. This maps $(0, \pi/2)$ to $(0, \infty)$. Since $dt = g'(\theta)d\theta = b \sec^2 \theta d\theta = (b + b \tan^2 \theta)d\theta$, $d\theta = bdt/(b^2 + t^2)$. Also

$$a^2 \cos^2 \theta + b^2 \sin^2 \theta = (a^2 + t^2) \cos^2 \theta = \frac{a^2 + t^2}{1 + \tan^2 \theta} = b^2 \cdot \frac{a^2 + t^2}{b^2 + t^2}.$$

Plugging into $I(a, b)$, we obtain the alternate formula

$$(3.5) \quad I(a, b) = \frac{2}{\pi} \int_0^\infty \frac{dt}{\sqrt{(a^2 + t^2)(b^2 + t^2)}}.$$

Now substitute $u = g(t) = (t - ab/t)/2$, and let $a' = (a + b)/2$, $b' = \sqrt{ab}$. This substitution takes $(0, \infty)$ to $(-\infty, \infty)$, and $g'(t) = (1 + ab/t^2)/2 = (t^2 + b'^2)/2t^2$. Thus,

$$du = \frac{(t^2 + b'^2) dt}{2t^2}.$$

Now,

$$b'^2 + u^2 = ab + \frac{1}{4} \left(t - \frac{ab}{t} \right)^2 = \frac{1}{4} \left(t + \frac{ab}{t} \right)^2 = \frac{(t^2 + b'^2)^2}{4t^2},$$

so,

$$(3.6) \quad \frac{du}{\sqrt{b'^2 + u^2}} = \frac{dt}{t}.$$

Also,

$$\begin{aligned} a'^2 + u^2 &= \left(\frac{a+b}{2} \right)^2 + \frac{1}{4} \left(t - \frac{ab}{t} \right)^2 \\ &= \frac{1}{4} \left(a^2 + b^2 + t^2 + \frac{a^2 b^2}{t^2} \right) \\ &= \frac{1}{4t^2} (a^2 + t^2)(b^2 + t^2). \end{aligned}$$

So,

$$(3.7) \quad \frac{1}{2\sqrt{a'^2 + u^2}} = \frac{t}{\sqrt{(a^2 + t^2)(b^2 + t^2)}}.$$

Multiplying (3.6) and (3.7),

$$\begin{aligned} I(a, b) &= \frac{2}{\pi} \int_0^\infty \frac{dt}{\sqrt{(a^2 + t^2)(b^2 + t^2)}} \\ &= \frac{2}{\pi} \int_{-\infty}^\infty \frac{du}{2\sqrt{(a'^2 + u^2)(b'^2 + u^2)}} \\ &= \frac{2}{\pi} \int_0^\infty \frac{du}{\sqrt{(a'^2 + u^2)(b'^2 + u^2)}} = I(a', b'), \end{aligned}$$

which is (3.4). Iterating (3.4), we obtain $I(a, b) = I(a_n, b_n)$ for all $n \geq 1$. Now, let $m = M(a, b)$. Since $I(a, b)$ is a continuous function of (a, b) (Exercise 1) and $a_n \rightarrow m$, $b_n \rightarrow m$, passing to the limit,

$$I(a, b) = \lim_{n \nearrow \infty} I(a_n, b_n) = I(m, m).$$

But $I(m, m) = 1/m$, hence, the result. \square

Next, we look at the behavior of $M(1, x)$, as $x \rightarrow 0+$. When $a_0 = 1$ and $b_0 = 0$, the arithmetic-geometric iteration yields $a_n = 2^{-n}$ and $b_n = 0$ for all

$n \geq 1$. Hence, $M(1, 0) = 0$. This leads us to believe that $M(1, x) \rightarrow 0$, as $x \rightarrow 0+$, or, what is the same, $1/M(1, x) \rightarrow \infty$, as $x \rightarrow 0+$. Exactly at what speed this happens leads us to another formula for π .

Theorem.

$$(3.8) \quad \lim_{x \rightarrow 0+} \left[\frac{\pi/2}{M(1, x)} - \log \left(\frac{4}{x} \right) \right] = 0.$$

To derive this, from (3.5),

$$(3.9) \quad \frac{\pi/2}{M(1, x)} = \int_0^\infty \frac{dt}{\sqrt{(1+t^2)(x^2+t^2)}}.$$

By Exercise 4, this equals

$$(3.10) \quad \frac{\pi/2}{M(1, x)} = 2 \int_0^{1/\sqrt{x}} \frac{dt}{\sqrt{(1+t^2)(1+(xt)^2)}}.$$

Now call the right side of (3.10) $I(x)$. Thus, the result will follow if we show that

$$(3.11) \quad \lim_{x \rightarrow 0+} \left[I(x) - \log \left(\frac{4}{x} \right) \right] = 0.$$

To derive (3.11), note that

$$\begin{aligned} J(x) &= 2 \int_0^{1/\sqrt{x}} \frac{dt}{\sqrt{1+t^2}} = 2 \log \left(t + \sqrt{1+t^2} \right) \Big|_0^{1/\sqrt{x}} \\ &= 2 \log \left(1 + \sqrt{x+1} \right) + \log \left(\frac{1}{x} \right) \\ &= 2 \log \left(\frac{1}{2} + \frac{1}{2} \sqrt{x+1} \right) + \log \left(\frac{4}{x} \right) \end{aligned}$$

and, so, $\log(4/x) - J(x) \rightarrow 0$ as $x \rightarrow 0+$. Thus, it is enough to show that

$$(3.12) \quad \lim_{x \rightarrow 0+} [I(x) - J(x)] = 0.$$

But, for $xt > 0$,

$$0 \leq 1 - \frac{1}{\sqrt{1+(xt)^2}} \leq 1 - \frac{1}{1+xt} = \frac{xt}{1+xt} \leq xt.$$

So,

$$\begin{aligned} 0 \leq J(x) - I(x) &= 2 \int_0^{1/\sqrt{x}} \frac{1}{\sqrt{1+t^2}} \left[1 - \frac{1}{\sqrt{1+(xt)^2}} \right] dt \\ &\leq 2 \int_0^{1/\sqrt{x}} \frac{xt}{\sqrt{1+t^2}} dt \end{aligned}$$

$$= 2x\sqrt{1+t^2}\Big|_0^{1/\sqrt{x}} = 2\sqrt{x(1+x)} - 2x,$$

which clearly goes to zero as $x \rightarrow 0+$. \square

Our next topic is the functional equation. First and foremost, since the AGM limit starting from (a_0, b_0) is the same as that starting from (a_1, b_1) ,

$$M(a, b) = M\left(\frac{a+b}{2}, \sqrt{ab}\right).$$

Below, given $0 < x < 1$ we let $x' = \sqrt{1-x^2}$ be the *complementary variable*. For example, $(x')' = x$ and $k = 2\sqrt{x}/(1+x)$ implies $k' = (1-x)/(1+x)$ since

$$\left(\frac{2\sqrt{x}}{1+x}\right)^2 + \left(\frac{1-x}{1+x}\right)^2 = 1.$$

Also, with $a_n, b_n, c_n, n \geq 1$, as above, $(b_n/a_n)' = c_n/a_n$. The functional equation we are after is best expressed in terms of the function

$$Q(x) = \frac{M(1, x)}{M(1, x')}, \quad 0 < x < 1.$$

Note that $Q(x') = 1/Q(x)$.

Theorem (AGM Functional Equation).

$$(3.13) \quad Q(x) = 2Q\left(\frac{1-x'}{1+x'}\right), \quad 0 < x < 1.$$

To see this, note that $M(1+x', 1-x') = M(1, x)$. So,

$$(3.14) \quad M(1, x) = M(1+x', 1-x') = (1+x')M\left(1, \frac{1-x'}{1+x'}\right).$$

Here, we used homogeneity of M . On the other hand,

$$(3.15) \quad \begin{aligned} M(1, x') &= M\left[(1+x')/2, \sqrt{x'}\right] = \frac{(1+x')}{2}M\left(1, \frac{2\sqrt{x'}}{1+x'}\right) \\ &= \frac{(1+x')}{2}M\left[1, \left(\frac{1-x'}{1+x'}\right)'\right]. \end{aligned}$$

Here, again, we used homogeneity of M . Dividing (3.14) by (3.15), the result follows. \square

If (a_n) and (b_n) are positive sequences, we say that (a_n) and (b_n) are *asymptotically equal*, and we write $a_n \sim b_n$, as $n \nearrow \infty$, if $a_n/b_n \rightarrow 1$, as $n \nearrow \infty$. Note that (a_n) and (b_n) are asymptotically equal iff $\log(a_n) - \log(b_n) \rightarrow 0$. Now, we combine the last two results to obtain the following improvement of (3.3).

Theorem. Let $a > b > 0$, and let $a_n, b_n, n \geq 1$, be as in (3.2i). Then,

$$(3.16) \quad a_n - b_n \sim 8M(a, b) \cdot q^{2^n}, \quad n \nearrow \infty,$$

where $q = e^{-\pi Q(b/a)}$.

To derive this, use (3.2ii) and $a_n \rightarrow M(a, b)$ to check that (3.16) is equivalent to

$$(3.17) \quad \frac{c_n}{4a_n} \sim \left(e^{-\pi Q(b/a)/2}\right)^{2^n}, \quad n \nearrow \infty.$$

Now, let $x_n = c_n/a_n$. Then, $x_n \rightarrow 0$, as $n \nearrow \infty$. By taking the log of (3.17), it is enough to show that

$$(3.18) \quad \log\left(\frac{4}{x_n}\right) - 2^n \frac{\pi}{2} Q\left(\frac{b}{a}\right) \rightarrow 0, \quad n \nearrow \infty.$$

By (3.8), (3.18) is implied by

$$(3.19) \quad \frac{1}{M(1, x_n)} - 2^n Q\left(\frac{b}{a}\right) \rightarrow 0, \quad n \nearrow \infty.$$

By Exercise 6, (3.19) is implied by

$$(3.20) \quad \frac{1}{Q(x_n)} - 2^n Q\left(\frac{b}{a}\right) \rightarrow 0, \quad n \nearrow \infty.$$

In fact, we will show that the left side of (3.20) is zero for all $n \geq 1$. To this end, since $c_n/a_n = (b_n/a_n)'$,

$$\begin{aligned} \frac{c_{n+1}}{a_{n+1}} &= \frac{a_n - b_n}{a_n + b_n} \\ &= \frac{1 - (b_n/a_n)}{1 + (b_n/a_n)} \\ &= \frac{1 - (c_n/a_n)'}{1 + (c_n/a_n)'}. \end{aligned}$$

Hence, by the functional equation,

$$Q(c_{n+1}/a_{n+1}) = Q\left(\frac{1 - (c_n/a_n)'}{1 + (c_n/a_n)'}\right) = \frac{1}{2} Q(c_n/a_n).$$

Iterating this down to $n = 1$, we obtain

$$\begin{aligned} Q(c_n/a_n) &= 2^{-1} Q(c_{n-1}/a_{n-1}) = 2^{-2} Q(c_{n-2}/a_{n-2}) = \dots \\ &\dots = 2^{-(n-1)} Q(c_1/a_1) = 2^{-n} Q((b/a)') = 2^{-n} / Q(b/a), \quad n \geq 1. \end{aligned}$$

This shows that $1/Q(x_n) = 2^n Q(b/a)$. \square

Dividing by 2^n in (3.18), we obtain

$$(3.21) \quad \lim_{n \nearrow \infty} 2^{-n} \log\left(\frac{a_n}{c_n}\right) = \frac{\pi}{2} Q\left(\frac{b}{a}\right),$$

which we will need in §5.7. Note that we have discarded the 4 since $2^{-n} \log 4 \rightarrow 0$.

Exercises 5.3

1. Use the dominated convergence theorem to show that $a_n \rightarrow a$ and $b_n \rightarrow b$, $a > b > 0$, implies $I(a_n, b_n) \rightarrow I(a, b)$.

2. Show that

$$\frac{1}{M(1+x, 1-x)} = \frac{2}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-x^2 \sin^2 \theta}}, \quad 0 < x < 1.$$

3. Show that

$$\frac{1}{M(1+x, 1-x)} = \sum_{n=0}^{\infty} \binom{2n}{n}^2 \frac{x^{2n}}{16^n}, \quad 0 < x < 1,$$

by using the binomial theorem to expand the square root and integrating term by term.

4. Show that

$$\frac{1}{M(1, x)} = \frac{4}{\pi} \int_0^{\sqrt{x}} \frac{dt}{\sqrt{(1+t^2)(x^2+t^2)}} = \frac{4}{\pi} \int_0^{1/\sqrt{x}} \frac{dt}{\sqrt{(1+(xt)^2)(1+t^2)}}.$$

(Break (3.9) into $\int_0^{\sqrt{x}} + \int_{\sqrt{x}}^{\infty}$, and substitute $t = x/s$ in the second piece.)

5. With $x' = \sqrt{1-x^2}$, show that $M(1+x, 1-x) = M((1+x')/2, \sqrt{x'})$.

6. Show that

$$\left| \frac{1}{M(1, x)} - \frac{1}{Q(x)} \right| \leq x, \quad 0 < x < 1.$$

7. Show that

$$Q(x) = \frac{1}{2} Q\left(\frac{2\sqrt{x}}{1+x}\right), \quad 0 < x < 1.$$

8. Show that $M(1, \cdot) : (0, 1) \rightarrow (0, 1)$ and $Q : (0, 1) \rightarrow (0, \infty)$ are strictly increasing, continuous bijections.

9. Show that for each $a > 1$, there exists a unique $1 > b = f(a) > 0$, such that $M(a, b) = 1$.

10. ★ With f as in the previous exercise, use (3.8) to show

$$f(a) \sim 4ae^{-\pi a/2}, \quad a \rightarrow \infty.$$

(Let $x = b/a = f(a)/a$ and take logs of both sides.)

§5.4 The Bell-Shaped Curve

In this section, we derive the *Gaussian integral*

$$(4.1) \quad \int_{-\infty}^{\infty} e^{-x^2/2} dx = \sqrt{2\pi}.$$

This formula is remarkable because the primitive of $e^{-x^2/2}$ cannot be expressed in terms of the elementary functions (i.e., the functions studied in Chapter 3). Nevertheless the area (Figure 5.4) of the (total) subgraph of $e^{-x^2/2}$ is explicitly computable. The *Gaussian* function $g(x) = e^{-x^2/2}$ is also called “the bell-shaped curve”.




FIGURE 5.4. The Gaussian.

The usual derivation of (4.1) involves changing variables from cartesian coordinates (x, y) to polar coordinates (r, θ) (§3.5) in a double integral. How to do this is a two-variable result. Here, we give an elementary derivation that uses only the one-variable material we have studied so far. To derive (4.1), we will, however, need to know how to “differentiate under an integral sign.”

To explain this, consider the integral

$$(4.2) \quad F(x) = \int_c^d f(x, t) dt, \quad a < x < b,$$

where $f(x, t) = 3(2x + t)^2$ and $a < b, c < d$ are reals. We wish to differentiate F . There are two ways we can do this. The first method is to evaluate the integral obtaining $F(x) = (2x + d)^3 - (2x + c)^3$ and, then, to differentiate to get $F'(x) = 6(2x + d)^2 - 6(2x + c)^2$. The second method is to differentiate the integrand $f(x, t) = 3(2x + t)^2$ with respect to x , obtaining $12(2x + t)$ and, then, to evaluate the integral $\int_c^d 12(2x + t) dt$, obtaining $6(2x + d)^2 - 6(2x + c)^2$. Since both methods yield the same result, for $f(x, t) = 3(2x + t)^2$, we conclude that

$$(4.3) \quad F'(x) = \int_c^d \frac{\partial f}{\partial x}(x, t) dt, \quad a < x < b,$$

where the *partial derivative* $\partial f / \partial x(x, t)$ is the derivative with respect to x ,

$$\frac{\partial f}{\partial x}(x, t) = \lim_{x' \rightarrow x} \frac{f(x', t) - f(x, t)}{x' - x}, \quad a < x < b.$$

It turns out that (4.2) implies (4.3) in a wide variety of cases.

Theorem (Differentiation Under the Integral Sign). *Let $f : (a, b) \times (c, d) \rightarrow \mathbf{R}$ be a continuous function, such that $\partial f / \partial x : (a, b) \times (c, d) \rightarrow \mathbf{R}$ exists and is continuous. Suppose that there is an integrable positive continuous function $g : (c, d) \rightarrow \mathbf{R}$, such that*

$$|f(x, t)| + \left| \frac{\partial f}{\partial x}(x, t) \right| \leq g(t), \quad a < x < b, c < t < d.$$

If $F : (a, b) \rightarrow \mathbf{R}$ is defined by (4.2), then, F is differentiable on (a, b) , $F' : (a, b) \rightarrow \mathbf{R}$ is continuous, and (4.3) holds.

Note that the domination hypothesis guarantees that $F(x)$ and the right side of (4.3) are well defined. Let us apply the theorem right away to obtain (4.1).

To this end, let $I = \int_0^\infty e^{-s^2/2} ds$ be half the integral in (4.1). Since $(s-1)^2 \geq 0$, $-s^2/2 \leq (1/2) - s$. Hence, $I \leq \int_0^\infty e^{(1/2)-s} ds = \sqrt{e}$. Thus, I is finite and

$$(4.4) \quad I^2 = I \int_0^\infty e^{-t^2/2} dt = \int_0^\infty e^{-t^2/2} I dt.$$

Now, set

$$f(x, t) = e^{-t^2/2} \int_0^{t \cdot \tan x} e^{-s^2/2} ds, \quad 0 < t < \infty, 0 < x < \pi/2.$$

Since $\tan(\pi/2-) = \infty$, by continuity at the endpoints, $f(\pi/2-, t) = e^{-t^2/2} I$ for all $t > 0$. Now, let

$$F(x) = \int_0^\infty f(x, t) dt, \quad 0 < x < \pi/2.$$

Since $f(x, t) \leq I e^{-t^2/2}$ and $g(t) = I e^{-t^2/2}$ is integrable by (4.4), by the dominated convergence theorem, we obtain

$$F(\pi/2-) = \lim_{x \rightarrow \pi/2-} \int_0^\infty f(x, t) dt = \int_0^\infty f(\pi/2-, t) dt = I^2.$$

Thus, to evaluate I^2 , we need to compute $F(x)$. Although $F(x)$ is not directly computable from its definition, it turns out that $F'(x)$ is, using differentiation under the integral sign.

To motivate where the formula for F comes from, note that the formula for I^2 can be thought of as a double integral over the first quadrant $0 < s < \infty$, $0 < t < \infty$, in the st -plane, and the formula for $F(x)$ can be thought of as a double integral over the triangular sector $0 < s < t \cdot \tan x$, $0 < t < \infty$, in the st -plane. As the angle x opens up to $\pi/2$, the triangular sector fills the quadrant. Of course, we do not actually use double integrals in the derivation of (4.1).

Now, by the fundamental theorem and the chain rule,

$$\frac{\partial f}{\partial x}(x, t) = e^{-t^2(1+\tan^2 x)/2} t \sec^2 x = e^{-t^2 \sec^2 x/2} t \sec^2 x.$$

We verify the hypotheses of the theorem on $(0, b) \times (0, \infty)$, where $0 < b < \pi/2$ is fixed. Note, first, that $f(x, t)$ and $\partial f/\partial x$ are continuous in (x, t) . Moreover, $0 \leq f(x, t) \leq I e^{-t^2/2}$ and $0 \leq \partial f/\partial x(x, t) \leq e^{-t^2/2} t \sec^2 b$ ($\sec^2 x \geq 1$ is increasing on $(0, \pi/2)$). So, we may take $g(t) = e^{-t^2/2}(I + t \sec^2 b)$, which is integrable.¹⁹ This verifies all the hypotheses. Applying the theorem yields

$$F'(x) = \int_0^\infty e^{-t^2 \sec^2 x/2} t \sec^2 x dt = \int_0^\infty e^{-u} du = 1, \quad 0 < x < b.$$

¹⁹ $\int_0^\infty g(t) dt = I^2 + \sec^2 b$.

Here, we used the substitution $u = t^2 \sec^2 x/2$, $du = t \sec^2 x dt$. Since $0 < b < \pi/2$ is arbitrary, $F'(x) = 1$ is valid on $(0, \pi/2)$.

Thus, $F(x) = x + \text{constant}$ on $(0, \pi/2)$. To evaluate the constant, note that $f(0+, t) = 0$ for all $0 < t < \infty$, by continuity at the endpoints. Then, since $f(x, t) \leq I e^{-t^2/2}$, we can apply the dominated convergence theorem to get

$$F(0+) = \lim_{x \rightarrow 0+} \int_0^\infty f(x, t) dt = \int_0^\infty f(0+, t) dt = 0.$$

This shows that $F(x) = x$, so, $F(\pi/2-) = \pi/2$. Hence, $I^2 = \pi/2$. Since I is half the integral in (4.1), this derives (4.1). \square

Let us apply the theorem to the gamma function

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt, \quad x > 0.$$

To this end, fix $0 < a < b < \infty$. We show that Γ is differentiable on (a, b) . With $f(x, t) = e^{-t} t^{x-1}$,

$$\frac{\partial f}{\partial x}(x, t) = e^{-t} t^{x-1} \log t, \quad 0 < t < \infty, 0 < x < \infty.$$

Then, f and $\partial f/\partial x$ are continuous on $(a, b) \times (0, \infty)$. Since $|f| + |\partial f/\partial x| \leq g(t)$ on $(a, b) \times (0, \infty)$, where

$$g(t) = \begin{cases} e^{-t} t^{a-1} (|\log t| + 1), & 0 < t \leq 1, \\ e^{-t} t^{b-1} (|\log t| + 1), & 1 \leq t, \end{cases}$$

and g is continuous and integrable over $(0, \infty)$ (Exercise 11 of §5.1), the domination hypothesis of the theorem is verified. Thus, we can apply the theorem to obtain

$$\Gamma'(x) = \int_0^\infty e^{-t} t^{x-1} \log t dt, \quad a < x < b.$$

Since $0 < a < b$ are arbitrary, this shows that Γ is differentiable on $(0, \infty)$. Since this argument can be repeated,

$$\Gamma''(x) = \int_0^\infty e^{-t} t^{x-1} (\log t)^2 dt, \quad x > 0.$$

Since this last quantity is positive, we see that Γ is strictly convex on $(0, \infty)$ (§3.3). Differentiating repeatedly we obtain $\Gamma^{(n)}(x)$ for all $n \geq 1$. Hence, the gamma function is smooth.

In Exercise 1 of §5.2, the Laplace transform

$$F(x) = \int_0^\infty e^{-xt} \frac{\sin t}{t} dt = \arctan\left(\frac{1}{x}\right)$$

is computed for $x > 1$ by expanding $\sin t/t$ in a series. Now, we compute $F(x)$ for $x > 0$ by using differentiation under the integral sign. In Exercise 12, we need to know this for $x > 0$. $x > 1$ is not enough. Note that, to compute $F(x)$ for $x > 0$, it is enough to compute $F(x)$ for $x > a$, where $a > 0$ is arbitrarily small.

First, $\partial f/\partial x = -e^{-xt} \sin t$, so, f and $\partial f/\partial x$ are continuous on $(a, \infty) \times (0, \infty)$. Since $\sin t$ and $\sin t/t$ are bounded by 1, $|f(x, t)| + |\partial f/\partial x|$ is dominated by $2e^{-at}$ on $(a, \infty) \times (0, \infty)$. Applying the theorem and Exercise 7 of §4.4 yields

$$F'(x) = - \int_0^\infty e^{-xt} \sin t \, dt = -\frac{1}{1+x^2}, \quad x > a.$$

Now, by the dominated convergence theorem, $F(\infty) = \lim_{x \rightarrow \infty} F(x) = 0$. So,

$$\begin{aligned} F(x) &= F(x) - F(\infty) = - \int_x^\infty F'(t) \, dt \\ &= \arctan t \Big|_x^\infty = \pi/2 - \arctan x = \arctan \left(\frac{1}{x} \right), \quad x > a. \end{aligned}$$

Since $a > 0$ is arbitrarily small, this is what we wanted to show.

Now, we derive the theorem. To this end, fix $a < x < b$, and let $x_n \rightarrow x$, with $x_n \neq x$ for all $n \geq 1$. We have to show that

$$\frac{F(x_n) - F(x)}{x_n - x} \rightarrow \int_c^d \frac{\partial f}{\partial x}(x, t) \, dt.$$

Let

$$k_n(t) = \frac{f(x_n, t) - f(x, t)}{x_n - x}, \quad c < t < d, n \geq 1,$$

and

$$k(t) = \frac{\partial f}{\partial x}(x, t), \quad c < t < d.$$

Then, $k_n(t)$ and $k(t)$ are continuous on (c, d) . By the mean value theorem²⁰

$$k_n(t) = \frac{\partial f}{\partial x}(x'_n, t), \quad c < t < d, n \geq 1,$$

for some x'_n between x_n and x . By the domination hypothesis, we see that $|k_n(t)| \leq g(t)$. Thus, we can apply the dominated convergence theorem, which yields

$$\frac{F(x_n) - F(x)}{x_n - x} = \int_c^d k_n(t) \, dt \rightarrow \int_c^d k(t) \, dt = \int_c^d \frac{\partial f}{\partial x}(x, t) \, dt.$$

This establishes (4.3). By continuity under the integral sign, (4.3) implies that F' is continuous. \square

Now, we compute $M(1, 1/\sqrt{2})$.

Theorem.

$$(4.5) \quad M\left(1, \frac{1}{\sqrt{2}}\right) = \frac{\Gamma(3/4)}{\Gamma(1/4)} \sqrt{2\pi}.$$

²⁰ x'_n also depends on t .

To this end, bring in the *beta function*²¹

$$(4.6) \quad B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad x > 0, y > 0.$$

The next result shows that $1/B(x, y)$ extends the binomial coefficient $\binom{x+y}{x}$ to nonnatural x and y .

Theorem. For all $a > 0$ and $b > 0$,

$$(4.7) \quad B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

We derive this following the method used to obtain (4.1). First, write

$$(4.8) \quad \Gamma(b)\Gamma(a) = \int_0^\infty \Gamma(b)e^{-t}t^{a-1} dt$$

and

$$\begin{aligned} \Gamma(b)e^{-t}t^{a-1} &= e^{-t}t^{a-1} \int_0^\infty e^{-s}s^{b-1} ds \\ &= \int_0^\infty e^{-s-t}s^{b-1}t^{a-1} ds \\ &= \int_t^\infty e^{-r}(r-t)^{b-1}t^{a-1} dr, \quad t > 0. \end{aligned}$$

Here, we substituted $r = s + t$, $dr = ds$. Now, set $h(t, r) = e^{-r}(r-t)^{b-1}t^{a-1}$,

$$f(x, t) = \int_{t/x}^\infty h(t, r) dr, \quad t > 0, 0 < x < 1,$$

and

$$F(x) = \int_0^\infty f(x, t) dt, \quad 0 < x < 1.$$

By continuity at the endpoints (the integrand is nonnegative), $f(1-, t) = \int_t^\infty h(t, r) dr = e^{-t}t^{a-1}\Gamma(b)$. Then, (4.8) says that $\int_0^\infty f(1-, t) dt = \Gamma(a)\Gamma(b)$. Since $f(x, t) \leq f(1-, t)$ for $0 < x < 1$ and $f(1-, t)$ is integrable, the dominated convergence theorem applies, and we conclude that $F(1-) = \Gamma(a)\Gamma(b)$.

Moreover, $F(0+) = 0$. To see this, note, by continuity at the endpoints (the integrand is integrable), that we have $f(0+, t) = 0$ for all $t > 0$. By the dominated convergence theorem, again, it follows that $F(0+) = 0$.

Now, by the fundamental theorem and the chain rule,

$$\frac{\partial f}{\partial x}(x, t) = -e^{-t/x}t^{a-1}(t/x - t)^{b-1} \left(-\frac{t}{x^2}\right)$$

²¹ $B(x, y)$ is finite by (4.7).

$$(4.9) \quad = \left(\frac{t}{x}\right)^{a+b-1} e^{-t/x} x^{a-1} (1-x)^{b-1} \cdot \frac{1}{x}.$$

Fix $0 < \epsilon < 1$. Then, by (4.9), $\partial f / \partial x$ is dominated by $e^{-t/\epsilon} (t/\epsilon)^{a+b-1} / \epsilon$ for $t > 0$ and $\epsilon \leq x < 1$. Thus, the domination hypothesis is verified with²² $g(t) = f(1-, t) + e^{-t/\epsilon} (t/\epsilon)^{a+b-1} / \epsilon$ on $(\epsilon, 1) \times (0, \infty)$. Differentiating under the integral sign and substituting $t/x = u$, $dt/x = du$,

$$\begin{aligned} F'(x) &= \int_0^\infty \left(\frac{t}{x}\right)^{a+b-1} e^{-t/x} x^{a-1} (1-x)^{b-1} \cdot \frac{1}{x} dt \\ &= \int_0^\infty u^{a+b-1} e^{-u} x^{a-1} (1-x)^{b-1} du = x^{a-1} (1-x)^{b-1} \Gamma(a+b), \end{aligned}$$

valid on $(\epsilon, 1)$. Since $\epsilon > 0$ is arbitrary, we obtain

$$F'(x) = x^{a-1} (1-x)^{b-1} \Gamma(a+b), \quad 0 < x < 1.$$

Integrating, we arrive at

$$\begin{aligned} \Gamma(a)\Gamma(b) &= F(1-) - F(0+) = \int_0^1 F'(x) dx \\ &= \Gamma(a+b) \int_0^1 x^{a-1} (1-x)^{b-1} dx = \Gamma(a+b) B(a, b), \end{aligned}$$

which is (4.7). \square

To derive (4.5), we use (4.7) and a sequence of substitutions. From §5.3,

$$\frac{\pi/2}{M(1, 1/\sqrt{2})} = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}}.$$

Substituting $\sin \theta = t$, we obtain

$$\frac{\pi/2}{M(1, 1/\sqrt{2})} = \sqrt{2} \int_0^1 \frac{dt}{\sqrt{(1-t^2)(2-t^2)}}.$$

Now, substitute $x^2 = t^2/(2-t^2)$ to obtain

$$\frac{\pi/2}{M(1, 1/\sqrt{2})} = \sqrt{2} \int_0^1 \frac{dx}{\sqrt{1-x^4}}.$$

Now, substitute $u = x^4$ to get

$$\frac{\pi/2}{M(1, 1/\sqrt{2})} = \frac{\sqrt{2}}{4} \int_0^1 u^{1/4-1} (1-u)^{1/2-1} du = \frac{\sqrt{2}}{4} B\left(\frac{1}{4}, \frac{1}{2}\right).$$

Since

$$B\left(\frac{1}{4}, \frac{1}{2}\right) = \frac{\Gamma(1/4)\Gamma(1/2)}{\Gamma(3/4)}$$

²² $\int_0^\infty g(t) dt = \Gamma(a)\Gamma(b) + \Gamma(a+b)/\epsilon^{a+b}$.

and (Exercise 1) $\Gamma(1/2) = \sqrt{\pi}$, we obtain (4.5). \square

We end the section with an important special case of the theorem. Suppose that $(c, d) = (0, \infty)$ and $f(x, t)$ is piecewise constant in t , i.e., suppose that $f(x, t) = f_n(x)$, $a < x < b$, $n - 1 < t \leq n$, $n \geq 1$. Then, the integral in (4.2) reduces to an infinite series. Hence, the theorem takes the following form.

Theorem (Differentiation Under the Summation Sign). *Let $f_n : (a, b) \rightarrow \mathbf{R}$, $n \geq 1$, be a sequence of differentiable functions with $f'_n : (a, b) \rightarrow \mathbf{R}$, $n \geq 1$, continuous. Suppose that there is a convergent positive series $\sum g_n$ of numbers, such that*

$$|f_n(x)| + |f'_n(x)| \leq g_n, \quad a < x < b, n \geq 1.$$

If

$$F(x) = \sum_{n=1}^{\infty} f_n(x), \quad a < x < b,$$

then, F is differentiable on (a, b) , $F' : (a, b) \rightarrow \mathbf{R}$ is continuous, and

$$F'(x) = \sum_{n=1}^{\infty} f'_n(x), \quad a < x < b.$$

To derive this, one, of course, applies the dominated convergence theorem for series instead of the theorem for integrals. \square

Let $f : (a, b) \times (c, d) \rightarrow \mathbf{R}$ be a function of two variables (x, y) , and suppose that $\partial f / \partial x$ exists. If $\partial f / \partial x$ is differentiable with respect to x , we denote its derivative by

$$\frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2}.$$

Similarly the derivative of $\partial f / \partial x$ with respect to y is denoted $\partial^2 f / \partial y \partial x$, the derivative of $\partial^2 f / \partial y \partial x$ with respect to x is denoted $\partial^3 f / \partial x \partial y \partial x$, and so on. Although we do not discuss this here, it is often, but not always, true that $\partial^2 f / \partial x \partial y = \partial^2 f / \partial y \partial x$.

Exercises 5.4

1. Use the substitution $x = \sqrt{2t}$ in (4.1) to obtain $\Gamma(1/2) = \sqrt{\pi}$. Conclude that $(1/2)! = \sqrt{\pi}/2$.
2. Show that the Laplace transform

$$L(s) = \int_{-\infty}^{\infty} e^{sx} e^{-x^2/2} dx, \quad -\infty < s < \infty,$$

is given by $L(s) = \sqrt{2\pi} e^{s^2/2}$. (Complete the square in the exponent, and use translation invariance.)

3. Compute $L^{(2n)}(0)$ with L as in the previous exercise, to obtain

$$\int_{-\infty}^{\infty} e^{-x^2/2} x^{2n} dx = \sqrt{2\pi} \cdot \frac{(2n)!}{2^n n!}, \quad n \geq 0.$$

(Writing out the power series of L yields $L^{(2n)}(0)$.)

4. Show that the Fourier transform

$$F(s) = \int_{-\infty}^{\infty} e^{-x^2/2} \cos(sx) dx, \quad -\infty < s < \infty,$$

is finite and differentiable on $(-\infty, \infty)$. Differentiate under the integral sign, and integrate by parts to show that $F'(s)/F(s) = -s$ for all s . Integrate this equation over $(0, s)$, and use $F(0) = \sqrt{2\pi}$ to obtain

$$F(s) = \sqrt{2\pi} e^{-s^2/2}.$$

5. ★ Derive the *Hecke integral*

$$(4.7) \quad H(a) = \int_0^{\infty} e^{-x-a/x} \frac{dx}{\sqrt{x}} = \sqrt{\pi} e^{-2\sqrt{a}}, \quad a > 0,$$

by differentiating under the integral sign and substituting $x = a/t$ to obtain $H'(a)/H(a) = -1/\sqrt{a}$. Integrate this equation over $(0, a)$, and use $H(0) = \Gamma(1/2) = \sqrt{\pi}$ to obtain (4.7).

6. Show that

$$\int_{-\infty}^{\infty} e^{-x^2/2q} dx = \sqrt{2\pi q}, \quad q > 0.$$

7. ★ Let $\psi(t) = \sum_{n=1}^{\infty} e^{-n^2\pi t}$, $t > 0$. Use the integral test (Exercise 8 of §4.3) to show that

$$\lim_{t \rightarrow 0+} \sqrt{t} \cdot \psi(t) = \frac{1}{2}.$$

8. Show that $\zeta(s) = \sum_{n=1}^{\infty} 1/n^s$, $s > 1$, is smooth. (Differentiation under the summation sign.)

9. Show that $\psi(t) = \sum_{n=1}^{\infty} e^{-n^2\pi t}$, $t > 0$, is smooth.

10. ★ Show that the Bessel function J_ν (Exercise 10 of §5.2) is smooth. If ν is an integer, show that J_ν satisfies *Bessel's equation*

$$x^2 J_\nu''(x) + x J_\nu'(x) + (x^2 - \nu^2) J_\nu(x) = 0, \quad -\infty < x < \infty.$$

(Differentiation under the integral sign and integration by parts.)

11. ★ Suppose that $f: \mathbb{R} \rightarrow \mathbb{R}$ is nonnegative, superlinear, and continuous, and let

$$F(s) = \int_{-\infty}^{\infty} e^{sx} e^{-f(x)} dx, \quad -\infty < s < \infty,$$

denote the Laplace transform of e^{-f} (Exercise 11 of §4.3). Show that F is smooth, and compute $(\log F)''$. Use the Cauchy–Schwarz inequality (Exercise 17 of §4.4) to conclude that $\log F$ is convex.

12. ★ Let $F(b) = \int_0^b \sin x/x \, dx$, $b > 0$. Integrate by parts to show that

$$\int_0^b e^{-sx} \frac{\sin x}{x} \, dx = e^{-sb} F(b) + s \int_0^b e^{-sx} F(x) \, dx, \quad s > 0.$$

Let $b \rightarrow \infty$, change variables on the right, and let $s \rightarrow 0+$ to get

$$\lim_{s \rightarrow 0+} \int_0^\infty e^{-sx} \frac{\sin x}{x} \, dx = \lim_{b \rightarrow \infty} \int_0^b \frac{\sin x}{x} \, dx.$$

Conclude that $F(\infty) = \pi/2$.

§5.5 Stirling's Approximation of $n!$

The main purpose of this section is to derive Stirling's approximation to $n!$. If (a_n) and (b_n) are positive sequences, we say that (a_n) and (b_n) are *asymptotically equal*, and we write $a_n \sim b_n$ as $n \nearrow \infty$, if $a_n/b_n \rightarrow 1$ as $n \nearrow \infty$. Note that $a_n \sim b_n$ as $n \nearrow \infty$ iff $\log a_n - \log b_n \rightarrow 0$ as $n \nearrow \infty$.

Theorem. *If x is any real, then,*

$$(5.1) \quad \Gamma(x+n) \sim n^{x+n-1/2} e^{-n} \sqrt{2\pi}, \quad n \nearrow \infty.$$

In particular, if $x = 1$, we have Stirling's approximation

$$n! \sim n^{n+1/2} e^{-n} \sqrt{2\pi}, \quad n \nearrow \infty.$$

Note that $\Gamma(x+n)$ is defined, as soon as $n > -x$. By taking the log of both sides, (5.1) is equivalent to

$$\lim_{n \nearrow \infty} \log \Gamma(x+n) - \left[\left(x+n - \frac{1}{2} \right) \log n - n \right] = \frac{1}{2} \log(2\pi).$$

To derive (5.1), recall that

$$(5.2) \quad \Gamma(x+n) = \int_0^\infty e^{-t} t^{x+n-1} \, dt, \quad x > 0.$$

Since this integral is the area of the subgraph of $e^{-t} t^{x+n-1}$ and all we want is an approximation, not an exact evaluation, of this integral, let us check where the integrand is maximized, as this will tell us where the greatest contribution to the area is located. A simple computation shows that the integrand is maximized at $t = x+n-1$, which goes to infinity with n . To get a handle on this region of maximum area, perform the change of variable $t = ns$. This leads to

$$(5.3) \quad \Gamma(x+n) = n^{x+n} \int_0^\infty e^{-ns} s^{x+n-1} \, ds = n^{x+n} \int_0^\infty e^{nf(s)} s^{x-1} \, ds,$$

where

$$f(s) = \log s - s, \quad s > 0.$$

Now the varying part $e^{nf(s)}$ of the integrand is maximized at the maximum of f , which occurs at $s = 1$, since $f(0+) = -\infty$, $f(\infty) = -\infty$. Since the maximum value of f at $s = 1$ is -1 , the maximum value of the integrand is roughly $e^{nf(1)} = e^{-n}$. By analogy with sums (Exercise 1), we expect the limiting behavior of the integral in (5.3) to involve the maximum value of the integrand. Let us pause in the derivation of Stirling's formula, and turn to the study of the limiting behavior of such integrals, in general.

Theorem. Suppose that $f : (a, b) \rightarrow \mathbf{R}$ is continuous and bounded above on a bounded interval (a, b) . Then,

$$(5.4) \quad \lim_{n \nearrow \infty} \frac{1}{n} \log \left[\int_a^b e^{nf(x)} dx \right] = \sup\{f(x) : a < x < b\}.$$

To see this (Figure 5.5), let I_n denote the integral, and let $M = \sup\{f(x) : a < x < b\}$. Then, M is finite since f is bounded above. Given $\epsilon > 0$, choose $c \in (a, b)$ with $f(c) > M - \epsilon$, and, by continuity, choose $\delta > 0$, such that $f(x) > f(c) - \epsilon$ on $(c - \delta, c + \delta)$. Then, $f(x) > M - 2\epsilon$ on $(c - \delta, c + \delta)$, and

$$(b - a)e^{nM} \geq I_n \geq \int_{c-\delta}^{c+\delta} e^{nf(x)} dx \geq \int_{c-\delta}^{c+\delta} e^{n(M-2\epsilon)} dx = 2\delta e^{n(M-2\epsilon)}.$$

Now, take the log of this last inequality, and divide by n to obtain

$$\frac{1}{n} \log(b - a) + M \geq \frac{1}{n} \log(I_n) \geq \frac{1}{n} \log(2\delta) + M - 2\epsilon.$$

Sending $n \nearrow \infty$, the upper and lower limits of the sequence $((1/n) \log(I_n))$ lie between M and $M - 2\epsilon$. Since $\epsilon > 0$ is arbitrary, $(1/n) \log(I_n) \rightarrow M$. \square




FIGURE 5.5. The global max is what counts.

Although a good start, this result is not quite enough to obtain Stirling's approximation. The exact form of the limiting behavior, due to Laplace, is given by the following.

Laplace's Theorem. Let $f : (a, b) \rightarrow \mathbf{R}$ be differentiable, f' continuous, and f concave. Suppose that f has a global maximum at $c \in (a, b)$ with f twice

differentiable at c and $f''(c) < 0$. Suppose that $g : (a, b) \rightarrow \mathbf{R}$ is continuous with polynomial growth and $g(c) > 0$. Then,

$$(5.5) \quad I_n = \int_a^b e^{nf(x)} g(x) dx \sim e^{nf(c)} g(c) \sqrt{\frac{2\pi}{-nf''(c)}}, \quad n \nearrow \infty.$$

By *polynomial growth*, we mean that $|g(x)| \leq A + B|x|^p$, $a < x < b$, for some constants A, B, p . Before we derive this theorem, let us apply it to obtain the asymptotic behavior of (5.3) to complete the derivation of (5.1).

In the case of (5.3), $f'(s) = 1/s - 1$, and $f''(s) = -1/s^2$, so, f is strictly concave, has a global maximum $f(1) = -1$ at $c = 1$, and $f''(1) = -1 < 0$. Since $g(s) = s^{x-1}$ has polynomial growth (in s), the integral in (5.3) is asymptotic to $e^{-n} \sqrt{2\pi/n}$, which yields (5.1). \square

Now, we derive Laplace's theorem. We write $I_n = I_n^- + I_n^0 + I_n^+$, where

$$\begin{aligned} I_n^- &= \int_a^{c-\delta} e^{nf(x)} g(x) dx, \\ I_n^0 &= \int_{c-\delta}^{c+\delta} e^{nf(x)} g(x) dx, \\ I_n^+ &= \int_{c+\delta}^b e^{nf(x)} g(x) dx. \end{aligned}$$

Since c is a maximum, $f'(c) = 0$. Since $f''(c)$ exists, by Taylor's theorem (§3.4), there is a continuous function $h : (a, b) \rightarrow \mathbf{R}$ satisfying $h(c) = f''(c)$, and

$$(5.6) \quad f(x) = f(c) + f'(c)(x - c) + \frac{1}{2}h(x)(x - c)^2 = f(c) + \frac{1}{2}h(x)(x - c)^2.$$

If we let $\mu_c(\delta)$ denote the modulus of continuity of h at c and let $\epsilon = \mu_c(\delta)$, then, $\epsilon \rightarrow 0$ as $\delta \rightarrow 0$. Thus, we can choose $\delta > 0$, such that $h(x) \leq f''(c) + \mu_c(\delta) = f''(c) + \epsilon < 0$ and $g(x) > 0$ on $(c - \delta, c + \delta)$. Now, substituting $x = c + t/\sqrt{n}$ in I_n^0 , $dx = dt/\sqrt{n}$, and inserting (5.6),

$$\begin{aligned} I_n^0 &= \int_{-\delta}^{\delta} e^{nf(c) + nh(x)(x-c)^2/2} g(x) dt \\ &= \frac{e^{nf(c)}}{\sqrt{n}} \int_{-\delta\sqrt{n}}^{\delta\sqrt{n}} e^{h(c+t/\sqrt{n})t^2/2} g(c + t/\sqrt{n}) dt. \end{aligned}$$

But $g(x)$ is bounded on $(c - \delta, c + \delta)$. Hence,

$$e^{h(c+t/\sqrt{n})t^2/2} |g(c + t/\sqrt{n})| \leq C e^{(f''(c)+\epsilon)t^2/2}, \quad |t| < \delta/\sqrt{n},$$

which is integrable²³ over $(-\infty, \infty)$. Thus, the dominated convergence theorem applies, and²⁴

$$\begin{aligned} e^{-nf(c)} I_n^0 \sqrt{n} &= \int_{-\delta\sqrt{n}}^{\delta\sqrt{n}} e^{h(c+t/\sqrt{n})t^2/2} g(c+t/\sqrt{n}) dt \\ &\rightarrow \int_{-\infty}^{\infty} e^{f''(c)t^2/2} g(c) dt = g(c) \sqrt{\frac{2\pi}{-f''(c)}}, \end{aligned}$$

by Exercise 6 of §5.4.

We conclude that

$$(5.7) \quad I_n^0 \sim e^{nf(c)} g(c) \sqrt{\frac{2\pi}{-nf''(c)}}.$$

To finish the derivation, it is enough to show that $I_n \sim I_n^0$ as $n \nearrow \infty$.

To derive $I_n \sim I_n^0$, it is enough to obtain $I_n^+/I_n^0 \rightarrow 0$ and $I_n^-/I_n^0 \rightarrow 0$, since

$$\frac{I_n}{I_n^0} = \frac{I_n^-}{I_n^0} + 1 + \frac{I_n^+}{I_n^0}, \quad n \geq 1.$$

To obtain $I_n^+/I_n^0 \rightarrow 0$, we use convexity. Since f is concave, $-f$ is convex. Hence, the graph of $-f$ on $(c+\delta, b)$ lies above its tangent line at $c+\delta$ (Exercise 3 of §3.3). Thus,

$$f(x) \leq f(c+\delta) + f'(c+\delta)(x-c-\delta), \quad c+\delta < x < b.$$

Since f is strictly concave at c , $f'(c+\delta) < 0$ and $f(c+\delta) < f(c)$. Inserting this in the definition for I_n^+ and substituting $x = t + c + \delta$,

$$\begin{aligned} |I_n^+| &\leq e^{nf(c+\delta)} \int_{c+\delta}^b e^{nf'(c+\delta)(x-c-\delta)} |g(x)| dx \\ &\leq e^{nf(c+\delta)} \int_0^{b-c-\delta} e^{nf'(c+\delta)t} (A + B|t+c+\delta|^p) dt \\ (5.8) \quad &\leq e^{nf(c+\delta)} \int_0^\infty e^{f'(c+\delta)t} (A + B|t+c+\delta|^p) dt \\ &= Ce^{nf(c+\delta)}, \end{aligned}$$

where C denotes the (finite) integral in (5.8). Now, divide this last expression by the expression in (5.7), obtaining

$$\begin{aligned} \left| \frac{I_n^+}{I_n^0} \right| &\leq \frac{Ce^{nf(c+\delta)}}{I_n^0} \sim \frac{Ce^{nf(c+\delta)}}{e^{nf(c)} g(c) \sqrt{\frac{2\pi}{-nf''(c)}}} \\ &= \text{constant} \cdot \sqrt{n} \cdot e^{-n(f(c)-f(c+\delta))}, \end{aligned}$$

²³The integral is $C\sqrt{2\pi/(-f''(c) - \epsilon)}$.

²⁴by Exercise 12 of §5.2.

which goes to zero as $n \nearrow \infty$ since the exponent is negative. Since I_n^-/I_n^0 is similar, this completes the derivation. \square

Since Stirling's approximation provides a manageable expression for $n!$, it is natural to use it to derive the asymptotics of the binomial coefficient

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}, \quad 0 \leq k \leq n.$$

Actually, of more interest is the binomial coefficient divided by 2^n , since this is the probability of obtaining k heads in n tosses of a fair coin. To this end, suppose that the number of heads in n tosses is k_n , and suppose that $k_n/n = t_n$ approaches a fixed limit $t \in (0, 1)$, as $n \nearrow \infty$. Applying Stirling to $n!$, $k_n! = (t_n n)!$, and $(n - k_n)! = ((1 - t_n)n)!$ and simplifying, we obtain the following:

Theorem. Fix $0 < t < 1$. If the number of heads $k = k_n$ in n tosses is such that the ratio $k_n/n \rightarrow t$, as $n \nearrow \infty$, then,

$$\binom{n}{k} 2^{-n} \sim \frac{1}{\sqrt{2\pi n}} \cdot \frac{1}{\sqrt{t(1-t)}} \cdot e^{-nH(t)}, \quad n \nearrow \infty,$$

where

$$H(t) = t \log(2t) + (1-t) \log[2(1-t)], \quad 0 < t < 1. \quad \square$$

Because the binomial coefficients are so basic, the function H which governs their asymptotic decay, must be important. The function H , called the *entropy*, controls the rate of decay of the binomial coefficients. Note that H is convex (Figure 5.6) on $(0, 1)$ and has a global minimum of zero at $t = 1/2$ with $H''(1/2) = 4$.




FIGURE 5.6. The entropy $H(x)$.

We end the section with an application of (5.1) to the following formula for the gamma function.

Theorem (The Duplication Formula). For $s > 0$,

$$2^{2s} \cdot \frac{\Gamma(s)\Gamma(s+1/2)}{\Gamma(2s)} = 2\sqrt{\pi}.$$

To derive this, let $f(s)$ denote the left side. Then, using $\Gamma(s+1) = s\Gamma(s)$, check that f is periodic of period 1, i.e., $f(s+1) = f(s)$. Hence, $f(s+n) = f(s)$

for all $n \geq 1$. Now, inserting the asymptotic (5.1) (three times) in the expression for $f(s+n)$ yields $f(s+n) \sim 2\sqrt{\pi}$, as $n \nearrow \infty$. Hence, $f(s) = 2\sqrt{\pi}$, which is the duplication formula. \square

Exercises 5.5

1. Show that

$$\lim_{n \nearrow \infty} (a^n + b^n + c^n)^{1/n} = \max(a, b, c), \quad a, b, c > 0,$$

and

$$\lim_{n \nearrow \infty} \frac{1}{n} \log (e^{na} + e^{nb} + e^{nc}) = \max(a, b, c), \quad a, b, c \in \mathbb{R}.$$

2. Use a computer to obtain $100!$ and its Stirling approximation s . Compute the relative error $|100! - s|/100!$.

3. Show that $\binom{2n}{n} 2^{-2n} \sim 1/\sqrt{\pi n}$ as $n \nearrow \infty$.

4. Apply Stirling to $n!$, $k!$, and $(n-k)!$ to derive the asymptotic for $\binom{n}{k} 2^{-n}$ given above.

5. Let $0 < p < 1$. Graph

$$H(t, p) = t \log(t/p) + (1-t) \log[(1-t)/(1-p)], \quad 0 < t < 1.$$

6. Suppose that, in tossing a flawed coin, the probability of obtaining heads in a single toss is p , where $0 < p < 1$. Show that the probability of obtaining $k = k_n$ heads in n tosses is

$$\binom{n}{k} p^k (1-p)^{n-k} \sim \frac{1}{\sqrt{2\pi n}} \cdot \frac{1}{\sqrt{t(1-t)}} \cdot e^{-nH(t,p)}, \quad n \nearrow \infty,$$

with $k_n/n \rightarrow t$, as $n \nearrow \infty$.

7. For $0 < q < 1$ and $0 < a < b < \infty$, let $f(q) = \int_a^b q^{x^2} dx$. Compute

$$\lim_{n \nearrow \infty} \frac{1}{n} \log f(q^n).$$

8. ★ Show that

$$3^{3s} \cdot \frac{\Gamma(s)\Gamma(s+1/3)\Gamma(s+2/3)}{\Gamma(3s)}$$

is periodic of period 1, hence, is a constant. Determine the constant. Generalize.

9. ★ Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be superlinear and continuous. Consider the Laplace transforms

$$L_n(s) = \int_{-\infty}^{\infty} e^{sx} e^{-nf(x)} dx, \quad n \geq 1.$$

Show that

$$\lim_{n \nearrow \infty} \frac{1}{n} \log [L_n(ns)] = \max_{-\infty < x < \infty} [sx - f(x)].$$

(Break $L(ns)$ into three pieces, as in Exercise 11 of §4.3, and use the second part of Exercise 1.)

10. Differentiate the log of the duplication formula to obtain

$$\frac{\Gamma'(1)}{\Gamma(1)} - \frac{\Gamma'(1/2)}{\Gamma(1/2)} = 2 \log 2.$$

11. ★ Use the duplication formula to get $\Gamma(1/4)\Gamma(3/4) = \pi\sqrt{2}$. Hence,

$$M\left(1, \frac{1}{\sqrt{2}}\right) = \frac{2\pi^{3/2}}{\Gamma^2(1/4)}.$$

§5.6 Infinite Products

Given a sequence (a_n) , let $p_n = (1 + a_1)(1 + a_2) \dots (1 + a_n)$ denote the n th partial product, $n \geq 1$. We say that the *infinite product* $\prod_{n=1}^{\infty} (1 + a_n)$ *converges* if there is a *finite* L , such that $p_n \rightarrow L$. In this case, we write

$$L = \prod_{n=1}^{\infty} (1 + a_n).$$

For example, by induction, check that, for $n \geq 1$,

$$(1+x)(1+x^2)(1+x^4) \dots (1+x^{2^{n-1}}) = 1 + x + x^2 + x^3 + \dots + x^{2^n-1}.$$

If $|x| < 1$, the sum converges. Hence, the product converges to

$$\prod_{n=0}^{\infty} (1 + x^{2^n}) = 1 + x + x^2 + x^3 + \dots = \frac{1}{1-x}, \quad |x| < 1.$$

If $\prod_{n=1}^{\infty} (1 + a_n)$ converges and $L \neq 0$, then, $1 + a_n = p_n/p_{n-1} \rightarrow L/L = 1$. Hence, a *necessary condition for convergence*, when $L \neq 0$, is $a_n \rightarrow 0$.

Theorem. For $x \neq 0$,

$$(6.1) \quad \frac{\sinh(\pi x)}{\pi x} = \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{n^2}\right).$$

To see this, we need the factorization

$$(N) \quad X^{2N} - 1 = (X^2 - 1) \cdot \prod_{n=1}^{N-1} (X^2 - 2X \cos(n\pi/N) + 1).$$

This factorization, trivial for $N = 2$, is most easily derived using complex numbers. However, by replacing X by X^2 in (N) and using the double-angle formula, $(2N)$ is easily derived as a consequence of (N) (Exercise 14 of §3.5). Hence, without recourse to complex numbers, one obtains (N) for $N = 2, 4, 8, \dots$. In fact, this is all we need to derive (6.1).

Insert $X = a/b$ in (N) and multiply through by b^{2N} , obtaining

$$a^{2N} - b^{2N} = (a^2 - b^2) \cdot \prod_{n=1}^{N-1} \left[a^2 - 2ab \cos\left(\frac{n\pi}{N}\right) + b^2 \right].$$

Now, because $(1 + a/n)^n \rightarrow e^a$ and $\sinh x = (e^x - e^{-x})/2$, it makes sense to insert

$$a = \left(1 + \frac{\pi x}{2N}\right), \quad b = \left(1 - \frac{\pi x}{2N}\right).$$

Simplifying and dividing by $2\pi x$, we obtain

$$\begin{aligned} & \frac{\left(1 + \frac{\pi x}{2N}\right)^{2N} - \left(1 - \frac{\pi x}{2N}\right)^{2N}}{2\pi x} \\ &= \frac{1}{N} \cdot \prod_{n=1}^{N-1} \left\{ 2 \left[1 - \cos\left(\frac{n\pi}{N}\right) \right] + \frac{\pi^2 x^2}{2N^2} \left[1 + \cos\left(\frac{n\pi}{N}\right) \right] \right\} \end{aligned}$$

(6.2)

$$= \frac{1}{N} \cdot \prod_{n=1}^{N-1} \left[4 \sin^2\left(\frac{n\pi}{2N}\right) + \frac{\pi^2 x^2}{N^2} \cos^2\left(\frac{n\pi}{2N}\right) \right],$$

where we used the double-angle formula, again. Taking the limit of both sides as $x \rightarrow 0$ using l'Hopital's rule (§3.2), we obtain

$$(6.3) \quad 1 = \frac{1}{N} \cdot \prod_{n=1}^{N-1} \left[4 \sin^2\left(\frac{n\pi}{2N}\right) \right].$$

Now divide (6.2) by (6.3), factor by factor, obtaining

$$\frac{\left(1 + \frac{\pi x}{2N}\right)^{2N} - \left(1 - \frac{\pi x}{2N}\right)^{2N}}{2\pi x} = \prod_{n=1}^{N-1} \left[1 + \frac{x^2}{n^2} \cdot f\left(\frac{n\pi}{2N}\right) \right],$$

where $f(x) = x^2 \cot^2 x$. To obtain (6.1), we wish to take the limit $N \nearrow \infty$. But $(\tan x)' = \sec^2 x \geq 1$. So, $\tan x \geq x$, so, $f(x) \leq 1$ on $(0, \pi/2)$. Thus,

$$\frac{\left(1 + \frac{\pi x}{2N}\right)^{2N} - \left(1 - \frac{\pi x}{2N}\right)^{2N}}{2\pi x} \leq \prod_{n=1}^{N-1} \left(1 + \frac{x^2}{n^2} \right).$$

Sending $N \nearrow \infty$ through powers of 2, we obtain

$$\frac{\sinh(\pi x)}{\pi x} = \frac{e^{\pi x} - e^{-\pi x}}{2\pi x} \leq \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{n^2} \right),$$

which is half of (6.1). On the other hand, for $M \leq N$,

$$\frac{\left(1 + \frac{\pi x}{2N}\right)^{2N} - \left(1 - \frac{\pi x}{2N}\right)^{2N}}{2\pi x} \geq \prod_{n=1}^{M-1} \left[1 + \frac{x^2}{n^2} \cdot f\left(\frac{n\pi}{2N}\right)\right].$$

Since $\lim_{x \rightarrow 0} f(x) = 1$, sending $N \nearrow \infty$ through powers of 2 in this last equation, we obtain

$$\frac{\sinh(\pi x)}{\pi x} \geq \prod_{n=1}^{M-1} \left(1 + \frac{x^2}{n^2}\right).$$

Now, let $M \nearrow \infty$, obtaining the other half of (6.1). \square

To give an example of the power of (6.1), take the log of both sides to get

$$(6.4) \quad \log\left(\frac{\sinh(\pi x)}{\pi x}\right) = \sum_{n=1}^{\infty} \log\left(1 + \frac{x^2}{n^2}\right), \quad x \neq 0.$$

Now, differentiate under the summation sign to obtain

$$(6.5) \quad \pi \coth(\pi x) - \frac{1}{x} = \sum_{n=1}^{\infty} \frac{2x}{n^2 + x^2}, \quad x \neq 0.$$

Here, $\coth = \cosh / \sinh$ is the *hyperbolic cotangent*. To justify this, note that $\log(1+t) = \int_0^t ds/(1+s) \leq \int_0^t ds = t$. Hence, $\log(1+t) \leq t$ for $t \geq 0$. Thus, with $f_n(x) = \log(1+x^2/n^2)$, $|f_n(x)| + |f'_n(x)| \leq (2b+b^2)/n^2 = g_n$ on $|x| < b$, and $\sum g_n < \infty$. Thus, (6.5) is valid on $0 < |x| < b$, hence, on $x \neq 0$. Now, dividing (6.5) by $2x$, letting $x \searrow 0$, and setting $t = \pi x$ yields²⁵

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^2} &= \lim_{x \searrow 0} \sum_{n=1}^{\infty} \frac{1}{n^2 + x^2} \\ &= \lim_{x \searrow 0} \frac{\pi x \coth(\pi x) - 1}{2x^2} = \pi^2 \cdot \lim_{t \searrow 0} \frac{t \coth t - 1}{2t^2}. \end{aligned}$$

But this last limit can be evaluated as follows. Since

$$\frac{\sinh t}{t} = 1 + \frac{t^2}{3!} + \frac{t^4}{5!} + \dots,$$

and

$$\cosh t = 1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \dots,$$

it follows that

$$\frac{t \coth t - 1}{2t^2} = \frac{1}{2t^2} \left(\frac{\cosh t}{\sinh t/t} - 1 \right) = \frac{1}{2t^2} \left(\frac{1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \dots}{1 + \frac{t^2}{3!} + \frac{t^4}{5!} + \dots} - 1 \right)$$

²⁵by the monotone convergence theorem for series.

$$\begin{aligned}
 &= \frac{1}{2t^2} \cdot \frac{\left(1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \frac{t^6}{6!} + \dots\right) - \left(1 + \frac{t^2}{3!} + \frac{t^4}{5!} + \frac{t^6}{7!} + \dots\right)}{1 + \frac{t^2}{3!} + \frac{t^4}{5!} + \dots} \\
 &= \frac{1}{2t^2} \cdot \frac{\frac{t^2}{3} + \frac{t^4}{3!5} + \frac{t^6}{5!7} + \dots}{1 + \frac{t}{3!} + \frac{t^2}{5!} + \dots} \\
 &= \frac{\frac{1}{6} + \frac{t^2}{60} + \frac{t^4}{1680} + \dots}{1 + \frac{t}{3!} + \frac{t^2}{5!} + \dots}.
 \end{aligned}$$

Now, take the limit, as $t \searrow 0$, obtaining the following:

Theorem.

$$\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \quad \square$$

Recalling the zeta function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad s > 1,$$

this result says that $\zeta(2) = \pi^2/6$, a result due to Euler. In fact, Euler used (6.5) to compute $\zeta(2n)$ for all $n \geq 1$. This computation involves certain rational numbers first studied by Bernoulli.

The *Bernoulli function* is defined by

$$\tau(x) = \begin{cases} \frac{x}{1 - e^{-x}}, & x \neq 0, \\ 1, & x = 0. \end{cases}$$

Clearly, τ is a smooth function on $x \neq 0$. The *Bernoulli numbers* B_n , $n \geq 0$, are defined by the *Bernoulli series*

$$(6.6) \quad \tau(x) = B_0 + B_1x + \frac{B_2}{2!}x^2 + \frac{B_3}{3!}x^3 + \dots$$

Since $1 - e^{-x} = x - x^2/2! + x^3/3! - x^4/4! + \dots$, $(1 - e^{-x})/x = 1 - x/2! + x^2/3! - x^3/4! + \dots$. Hence, to obtain the B_n 's, one computes the reciprocal of this last series, which is obtained by setting the Cauchy product (§1.7)

$$\left(1 - \frac{x}{2!} + \frac{x^2}{3!} - \frac{x^3}{4!} + \dots\right) \left(B_0 + B_1x + \frac{B_2}{2!}x^2 + \frac{B_3}{3!}x^3 + \dots\right) = 1.$$

Multiplying, this leads to $B_0 = 1$ and the recursion formula

$$(6.7) \quad \frac{B_{n-1}}{(n-1)!1!} - \frac{B_{n-2}}{(n-2)!2!} + \dots + (-1)^{n-1} \frac{B_0}{0!n!} = 0, \quad n \geq 2.$$

Computing, we see that each B_n is a rational number with

$$\begin{aligned} B_1 &= \frac{1}{2}, & B_2 &= \frac{1}{6}, & B_4 &= -\frac{1}{30}, \\ B_6 &= \frac{1}{42}, & B_8 &= -\frac{1}{30}, & B_{10} &= \frac{5}{66}, & \dots \end{aligned}$$

It turns out (Exercise 2) that $|B_n| \leq 2^n n!$. Hence, by the root test, the Bernoulli series (6.6) converges, at least, for $|x| < 1/2$. In particular, this shows that τ is smooth near zero. Hence, τ is smooth on \mathbb{R} . Let $2\pi\beta > 0$ denote the radius²⁶ of convergence of (6.6). Then, (6.6) holds for $|x| < 2\pi\beta$. Since

$$\frac{x}{1-e^{-x}} - \frac{x}{2} = \frac{x}{2} \cdot \frac{1+e^{-x}}{1-e^{-x}} = \frac{x}{2} \cdot \frac{e^{x/2} + e^{-x/2}}{e^{x/2} - e^{-x/2}} = \frac{x}{2} \coth\left(\frac{x}{2}\right),$$

subtracting $x/2 = B_1x$ from both sides of (6.6), we obtain

$$\frac{x}{2} \coth\left(\frac{x}{2}\right) = 1 + \sum_{n=2}^{\infty} \frac{B_n}{n!} x^n, \quad 0 < |x| < 2\pi\beta.$$

But $(x/2) \coth(x/2)$ is even. Hence, $B_3 = B_5 = B_7 = \dots = 0$, and

$$\frac{x}{2} \coth\left(\frac{x}{2}\right) - 1 = \sum_{n=1}^{\infty} \frac{B_{2n}}{(2n)!} x^{2n}, \quad 0 < |x| < 2\pi\beta.$$

Now, replacing x by $2\pi\sqrt{x}$ and dividing by x ,

$$\frac{\pi\sqrt{x} \coth(\pi\sqrt{x}) - 1}{x} = \sum_{n=1}^{\infty} \frac{B_{2n}}{(2n)!} (2\pi)^{2n} x^{n-1}, \quad 0 < x < \beta^2.$$

Thus, from (6.5), we conclude that

$$\sum_{n=1}^{\infty} \frac{B_{2n}}{(2n)!} 2^{2n-1} \pi^{2n} x^{n-1} = \sum_{n=1}^{\infty} \frac{1}{n^2 + x}, \quad 0 < x < \beta^2.$$

Since the left side is a power series, we may differentiate it term by term (§3.4). On the other hand, the right side may²⁷ be differentiated under the summation sign. Differentiating both sides $r-1$ times,

$$\sum_{n=r}^{\infty} \frac{B_{2n}}{(2n)!} 2^{2n-1} \pi^{2n} \cdot \frac{(n-1)!}{(n-r)!} x^{n-r} = \sum_{n=1}^{\infty} \frac{(-1)^{r-1} (r-1)!}{(n^2 + x)^r}, \quad 0 < x < \beta^2.$$

Sending $x \rightarrow 0+$, the right side becomes $(-1)^{r-1} (r-1)! \zeta(2r)$, whereas the left side reduces to the first coefficient (that corresponding to $n=r$). We have derived the following.

²⁶In fact, although we do not show it here, $\beta = 1$, and the radius is 2π .

²⁷With $f_n(x) = 1/(n^2 + x)$ and $g_n = r!/n^2$, $|f_n(x)| + |f'_n(x)| + \dots + |f_n^{(r-1)}(x)| \leq g_n$ and $\sum g_n = r! \zeta(2)$.

Theorem. For all $n \geq 1$,

$$\zeta(2n) = (-1)^{n-1} \frac{B_{2n}}{(2n)!} 2^{2n-1} \cdot \pi^{2n}. \quad \square$$

Above, we saw that relating an infinite series to an infinite product led to some nice results. In particular, we derived the infinite product for $\sinh \pi x / \pi x$, which we rewrite, now, as

$$(6.8) \quad 1 + \frac{\pi^2 x^2}{3!} + \frac{\pi^4 x^4}{5!} + \cdots = \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{n^2}\right), \quad x \neq 0.$$

We wish to derive the analog of this result for the sine function, i.e., we want to obtain (6.8) with $-x^2$ replacing x^2 .

To this end, consider the following identity

$$(6.9) \quad 1 + b_1 x + b_2 x^2 + \cdots = \prod_{n=1}^{\infty} (1 + a_n x), \quad 0 < x < R.$$

We seek the relations between (a_n) and (b_n) . As a special case, if we suppose that $a_n = 0$ for all $n \geq 3$, (6.9) reduces to

$$1 + b_1 x + b_2 x^2 = (1 + a_1 x)(1 + a_2 x)$$

which implies $b_1 = a_1 + a_2$ and $b_2 = a_1 a_2$. Similarly, if we suppose that $a_n = 0$ for all $n \geq 4$, (6.9) reduces to

$$1 + b_1 x + b_2 x^2 + b_3 x^3 = (1 + a_1 x)(1 + a_2 x)(1 + a_3 x)$$

which implies

$$b_1 = a_1 + a_2 + a_3 = \sum_i a_i,$$

$$b_2 = a_1 a_2 + a_1 a_3 + a_2 a_3 = \sum_{i < j} a_i a_j,$$

and

$$b_3 = a_1 a_2 a_3 = \sum_{i < j < k} a_i a_j a_k.$$

Theorem. Suppose that (a_n) and (b_n) are positive sequences and the series in (6.9) converges on $(-R, R)$. If (6.9) holds, then,

$$(6.10) \quad 1 - b_1 x + b_2 x^2 - \cdots = \prod_{n=1}^{\infty} (1 - a_n x), \quad 0 < x < R.$$

We call (6.10) the *alternating version* of (6.9). Let us immediately apply this theorem to derive the infinite product for the sine. Replacing x by \sqrt{x} in (6.8),

we obtain

$$(6.11) \quad 1 + \frac{\pi^2 x}{3!} + \frac{\pi^4 x^2}{5!} + \cdots = \prod_{n=1}^{\infty} \left(1 + \frac{x}{n^2}\right), \quad x > 0.$$

Now, the alternating version of (6.11) is given by

$$1 - \frac{\pi^2 x}{3!} + \frac{\pi^4 x^2}{5!} - \cdots = \prod_{n=1}^{\infty} \left(1 - \frac{x}{n^2}\right), \quad x > 0.$$

Replacing x by x^2 in the last equation leads to

$$1 - \frac{\pi^2 x^2}{3!} + \frac{\pi^4 x^4}{5!} - \cdots = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2}\right), \quad x \neq 0.$$

But this last series is the series for $\sin(\pi x)/\pi x$.

Theorem. For $x \neq 0$,

$$(6.12) \quad \frac{\sin(\pi x)}{\pi x} = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2}\right). \quad \square$$

This is the alternating version of (6.1).

To derive (6.10) from (6.9), write the finite version of (6.9),

$$(6.13) \quad 1 + b_1^{(N)}x + b_2^{(N)}x^2 + \cdots + b_N^{(N)}x^N = \prod_{n=1}^N (1 + a_n x),$$

and let $b_1^{(N)}, b_2^{(N)}, \dots, b_N^{(N)}$ denote the coefficients obtained by expanding the right side. Then,

$$b_1^{(N)} = \sum_{i=1}^N a_i \nearrow \sum_{i=1}^{\infty} a_i = b_1^{(\infty)},$$

as $N \nearrow \infty$,

$$b_2^{(N)} = \sum_{1 \leq i < j \leq N} a_i a_j \nearrow \sum_{1 \leq i < j < \infty} a_i a_j = b_2^{(\infty)},$$

as $N \nearrow \infty$, and so on. Here, $b_1^{(\infty)}, b_2^{(\infty)}, \dots$, are *defined* as the positive infinite sums $\sum_i a_i, \sum_{i < j} a_i a_j, \dots$. We want to show that $(b_n^{(\infty)})$ equals the given sequence (b_n) . For this, let $N \nearrow \infty$ in (6.13). Since x is positive, there is no problem with the limits (everything is increasing) and we get

$$1 + b_1^{(\infty)}x + b_2^{(\infty)}x^2 + \cdots = \prod_{n=1}^{\infty} (1 + a_n x), \quad 0 < x < R.$$

Since the coefficients of a power series are unique (Exercise 7 of §3.4), this and (6.9) yield $b_n = b_n^{(\infty)}$ for $n \geq 1$. Hence, $b_n^{(N)} \nearrow b_n$ for all $n \geq 1$, as $N \nearrow \infty$.

Now, replace x by $-x$ in (6.13) to get

$$1 - b_1^{(N)}x + b_2^{(N)}x^2 - \cdots + (-1)^N b_N^{(N)}x^N = \prod_{n=1}^N (1 - a_n x), \quad 0 < x < R.$$

Clearly, as $N \nearrow \infty$, the right side of this last equation decreases to the right side of (6.10). Thus, to derive the theorem, it is enough to show that

$$\sum_{n=1}^N (-1)^n b_n^{(N)} x^n \rightarrow \sum_{n=1}^{\infty} (-1)^n b_n x^n, \quad N \nearrow \infty.$$

But, for $0 < x < R$,

$$\left| \sum_{n=1}^N (-1)^n b_n^{(N)} x^n - \sum_{n=1}^{\infty} (-1)^n b_n x^n \right| \leq \sum_{n=1}^N [b_n - b_n^{(N)}] x^n + \sum_{n=N+1}^{\infty} b_n x^n.$$

Now, the second sum on the right is the tail (§1.6) of a convergent series, hence, goes to zero, as $N \nearrow \infty$, whereas the first sum on the right goes to zero by the dominated convergence theorem for series. Indeed, the terms in the first sum on the right are no greater than $g_n = b_n x^n$ with $\sum g_n$ finite by assumption. Thus, we arrive at (6.10). \square

Exercises 5.6

1. Compute $\zeta(4)$, $\zeta(6)$, $\zeta(8)$.
2. Use the recursion (6.7) to derive $|B_n| \leq 2^n n!$, $n \geq 1$. Conclude that the Bernoulli series (6.6) converges on $(-1/2, 1/2)$. Also show that the numbers (B_2, B_4, B_6, \dots) , form an alternating sequence $(+, -, +, \dots)$.
3. If (a_n) is a positive sequence, then,

$$(6.14) \quad \sum_{n=1}^{\infty} a_n \leq \prod_{n=1}^{\infty} (1 + a_n) \leq \exp \left(\sum_{n=1}^{\infty} a_n \right).$$

Conclude that $\sum a_n < \infty$ iff $\prod (1 + a_n) < \infty$.

4. Use Exercise 5 of §5.1 (Eq. (1.3)) to show that

$$\Gamma(x) = \frac{e^{-\gamma x}}{x} \prod_{n=1}^{\infty} \left[\frac{e^{x/n}}{1 + \frac{x}{n}} \right], \quad x > 0,$$

where γ is Euler's constant (Exercise 14 of §4.4). (Use $1 + 1/2 + \cdots + 1/n - \log n \rightarrow \gamma$ and $n^x = e^{x \log n}$.)

5. Use Exercise 5 of §5.1 applied to $\Gamma(x)$ and $\Gamma(1 - x)$ to show that

$$\frac{\pi}{\Gamma(x)\Gamma(1-x)} = \sin(\pi x), \quad 0 < x < 1.$$

6. ★ Let

$$B(x) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{B_{2n}}{(2n)!} x^{2n}, \quad |x| < 2\pi\beta.$$

Use Exercise 8 of §1.7 to show that $(x/2) \cos(x/2) = B(x) \sin(x/2)$ for $|x| < 2\pi\beta$. Conclude that

$$\frac{x}{2} \cot\left(\frac{x}{2}\right) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{B_{2n}}{(2n)!} x^{2n}, \quad 0 < |x| < \min(2\pi, 2\pi\beta).$$

7. ★ Use Exercise 6 to conclude that $\beta \leq 1$, i.e., the radius of convergence of the Bernoulli series (6.6) is no more than 2π .

8. ★ Use (6.12) and modify the development leading up to (6.5) to obtain

$$\pi \cot(\pi x) - \frac{1}{x} = \sum_{n=1}^{\infty} \frac{2x}{x^2 - n^2}, \quad 0 < |x| < 1.$$

9. ★ Use Exercise 6 above and Exercise 13 of §3.5 to obtain

$$\tan x = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{B_{2n}}{(2n)!} 2^{2n} (2^{2n} - 1) x^{2n-1}, \quad |x| < \pi\beta/2.$$

10. Use Exercise 4, and differentiate under the summation sign to get

$$\frac{d}{dx} \log \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)} = -\gamma - \frac{1}{x} + \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+x} \right), \quad x > 0.$$

11. Use the previous exercise to show that $\Gamma'(1) = -\gamma$ and $\Gamma'(2) = 1 - \gamma$. Conclude that the global minimum of $\Gamma(x)$, $x > 0$, lies in the interval $(1, 2)$.

12. Differentiate under the summation sign to obtain the Laplace transform of τ ,

$$\frac{d^2}{dx^2} \log \Gamma(x) = \int_0^{\infty} e^{-xt} \tau(t) dt, \quad x > 0.$$

(Exercise 10 above and Exercise 13 of §5.1.)

13. Use Exercise 10 to show

$$\lim_{x \rightarrow 1-} \left\{ -\frac{1}{2} \frac{\Gamma'[(1-x)/2]}{\Gamma[(1-x)/2]} + \frac{1}{x-1} \right\} = \frac{1}{2}\gamma.$$

§5.7 Jacobi's Theta Functions

The *theta function* is defined by

$$\theta(s) = \sum_{n=-\infty}^{\infty} e^{-n^2\pi s} = 1 + 2e^{-\pi s} + 2e^{-4\pi s} + 2e^{-9\pi s} + \dots, \quad s > 0.$$

This positive sum, over all integers n (positive and negative and zero), converges for $s > 0$ since

$$\begin{aligned}\sum_{-\infty}^{\infty} e^{-n^2\pi s} &\leq \sum_{-\infty}^{\infty} e^{-|n|\pi s} \\ &= 1 + 2 \sum_{n=1}^{\infty} e^{-n\pi s} = 1 + \frac{2e^{-\pi s}}{1 - e^{-\pi s}} < \infty.\end{aligned}$$

Recall (Exercise 9 of §5.1) the function $\psi(s) = \sum_{n=1}^{\infty} e^{-n^2\pi s}$. This is related to θ by $\theta = 1 + 2\psi$. The main result in this section is the following remarkable identity, which we need in the next section.

Theorem (Theta Functional Equation). *For all $s > 0$,*

$$(7.1) \quad \theta(1/s) = \sqrt{s}\theta(s), \quad s > 0,$$

which can be rewritten as

$$(7.1) \quad \sum_{-\infty}^{\infty} e^{-n^2\pi/s} = \sqrt{s} \sum_{-\infty}^{\infty} e^{-n^2\pi s}, \quad s > 0.$$

As one indication of the power of (7.1), plug in $s = .01$. Then, the series for $\theta(.01)$ converges slowly (the tenth term is $1/e^\pi$), whereas the series for $\theta(100)$ converges quickly. In fact, the sum $\theta(100)$ of the series on the left differs from its zeroth term 1 by less than 10^{-100} .

In terms of ψ , the functional equation becomes

$$(7.2) \quad 1 + 2\psi(1/s) = \sqrt{s}[1 + 2\psi(s)].$$

To derive (7.1), we need to introduce three power series, *Jacobi's theta functions*, and relate them to the arithmetic-geometric mean of §5.3. These functions' most striking property, double-periodicity, does not appear unless one embraces the complex plane. Nevertheless, within the confines of the real line, we shall be able to get somewhere.

The Jacobi theta functions, defined for $|q| < 1$, are defined by

$$\begin{aligned}\theta_0(q) &= \sum_{-\infty}^{\infty} q^{n^2} = 1 + 2q + 2q^4 + 2q^9 + \dots, \\ \theta_+(q) &= \sum_{-\infty}^{\infty} q^{(n+1/2)^2} = 2q^{1/4} + 2q^{9/4} + 2q^{25/4} + \dots,\end{aligned}$$

and

$$\theta_-(q) = \sum_{-\infty}^{\infty} (-1)^n q^{n^2} = 1 - 2q + 2q^4 - 2q^9 + \dots.$$

By comparing these series with the geometric series, we see that they all converge for $|q| < 1$.

The simplest properties of these functions depend on parity properties of integers. For example, because n is odd iff n^2 is odd, $\theta_0(-q) = \theta_-(q)$, since

$$\theta_0(-q) = \sum_{-\infty}^{\infty} (-1)^{n^2} q^{n^2} = \sum_{-\infty}^{\infty} (-1)^n q^{n^2} = \theta_-(q).$$

Similarly, since $\theta_-(q)$ is the alternating version (§1.7) of $\theta_0(q)$,

$$(7.3) \quad \theta_0(q) + \theta_-(q) = 2 \sum_{n \text{ even}} q^{n^2} = 2 \sum_{-\infty}^{\infty} q^{(2n)^2} = 2 \sum_{-\infty}^{\infty} (q^4)^{n^2} = 2\theta_0(q^4).$$

In the remainder of the section, we restrict q to lie in the interval $(0, 1)$. In this case, $\theta_0(q) \geq 1$, $\theta_-(q)$ is bounded in absolute value by 1 by the Leibnitz test, and, hence, $\theta_-^2(q) \leq \theta_0^2(q)$.

For $n \geq 0$, let $\sigma(n)$ be the number of ways of writing n as a sum of squares, $n = i^2 + j^2$, with $i, j \in \mathbb{Z}$, where permutations and signs are taken into account. Thus,

$$\begin{aligned} \sigma(0) &= 1 & \text{because} & \quad 0 = 0^2 + 0^2, \\ \sigma(1) &= 4 & \text{because} & \quad 1 = (\pm 1)^2 + 0^2 = 0^2 + (\pm 1)^2, \\ \sigma(2) &= 4 & \text{because} & \quad 2 = (\pm 1)^2 + (\pm 1)^2, \\ \sigma(3) &= 0, \\ \sigma(4) &= 4 & \text{because} & \quad 4 = (\pm 2)^2 + 0^2 = 0^2 + (\pm 2)^2, \\ \sigma(5) &= 8 & \text{because} & \quad 5 = (\pm 2)^2 + (\pm 1)^2 = (\pm 1)^2 + (\pm 2)^2, \\ \sigma(6) &= \sigma(7) = 0, \\ \sigma(8) &= 4 & \text{because} & \quad 8 = (\pm 2)^2 + (\pm 2)^2, \\ \sigma(9) &= 4 & \text{because} & \quad 9 = (\pm 3)^2 + 0^2 = 0^2 + (\pm 3)^2, \\ \sigma(10) &= 8 & \text{because} & \quad 10 = (\pm 1)^2 + (\pm 3)^2 = (\pm 3)^2 + (\pm 1)^2, \\ & \text{etc.} \end{aligned}$$

Then,

$$\begin{aligned} \theta_0^2(q) &= \left(\sum_{-\infty}^{\infty} q^{n^2} \right)^2 = \left(\sum_{-\infty}^{\infty} q^{i^2} \right) \left(\sum_{-\infty}^{\infty} q^{j^2} \right) \\ (7.4) \quad &= \sum_{i, j \in \mathbb{Z}} q^{i^2 + j^2} = \sum_{n=0}^{\infty} \left(\sum_{i^2 + j^2 = n} q^n \right) = \sum_{n=0}^{\infty} \sigma(n) q^n \\ &= 1 + 4q + 4q^2 + 4q^4 + 8q^5 + 4q^8 + 4q^9 + 8q^{10} + \dots \end{aligned}$$

Similarly, since n is even iff n^2 is even,

$$(7.5) \quad \theta_-^2(q) = \sum_{n=0}^{\infty} (-1)^n \sigma(n) q^n.$$

Now, if $n = i^2 + j^2$, then, $2n = (i + j)^2 + (i - j)^2 = k^2 + \ell^2$. Conversely, if $2n = k^2 + \ell^2$, then, $n = ((k + \ell)/2)^2 + (k - \ell)/2)^2 = i^2 + j^2$. Thus,

$$\sigma(2n) = \sigma(n), \quad n \geq 1.$$

For example, $\sigma(1) = \sigma(2) = \sigma(4) = \sigma(8)$. Here, we used the fact that $k^2 + \ell^2$ is even iff $k + \ell$ is even iff $k - \ell$ is even. Since the series (7.5) is the alternating version of the series (7.4),

$$\begin{aligned} \theta_0^2(q) + \theta_-^2(q) &= 2 \sum_{n \text{ even}} \sigma(n) q^n \\ (7.6) \quad &= 2 \sum_{n=0}^{\infty} \sigma(2n) q^{2n} = 2 \sum_{n=0}^{\infty} \sigma(n) (q^2)^n = 2\theta_0^2(q^2). \end{aligned}$$

Now, subtract (7.6) from the square of (7.3). You obtain

$$\begin{aligned} 2\theta_0(q)\theta_-(q) &= [\theta_0(q) + \theta_-(q)]^2 - [\theta_0^2(q) + \theta_-^2(q)] \\ (7.7) \quad &= 4\theta_0^2(q^4) - 2\theta_0^2(q^2) = 2\theta_-^2(q^2), \end{aligned}$$

where the last equality is by (7.6) again. Rewriting (7.6) and (7.7), we have arrived at the AGM iteration

$$\begin{aligned} (7.8) \quad &\frac{\theta_0^2(q) + \theta_-^2(q)}{2} = \theta_0^2(q^2), \\ &\sqrt{\theta_0^2(q)\theta_-^2(q)} = \theta_-^2(q^2). \end{aligned}$$

Setting $a_0 = \theta_0^2(q)$ and $b_0 = \theta_-^2(q)$, let $(a_n), (b_n)$, be the AGM iteration (3.2i). Iterating (7.8), we obtain

$$a_n = \theta_0^2(q^{2^n}),$$

and

$$b_n = \theta_-^2(q^{2^n}).$$

$n \geq 0$. Since $\theta_0(0) = 1 = \theta_-(0)$, $q^{2^n} \rightarrow 0$, and $a_n \rightarrow M(a_0, b_0)$, $b_n \rightarrow M(a_0, b_0)$, we arrive at $M(a_0, b_0) = 1$ or $M(\theta_0^2(q), \theta_-^2(q)) = 1$.

Theorem. Suppose that (a, b) lies in the first quadrant of the ab -plane with $a > 1 > b$. Then, (a, b) lies on the AGM curve $M(a, b) = 1$ iff $(a, b) = (\theta_0^2(q), \theta_-^2(q))$ for a unique $0 < q < 1$. In particular,

$$(7.9) \quad M(\theta_0^2(q), \theta_-^2(q)) = 1, \quad 0 < q < 1.$$

Above we derived (7.9). To get the rest, suppose that $a > 1 > b > 0$ and $M(a, b) = 1$. Since $\theta_0^2 : (0, 1) \rightarrow (1, \infty)$ is a bijection (Exercise 1), there is a unique q in $(0, 1)$ satisfying $a = \theta_0^2(q)$. Then, by (7.9), $M[a, \theta_-^2(q)] = 1 = M(a, b)$. Since $b \mapsto M(a, b)$ is strictly increasing, we must have $b = \theta_-^2(q)$. \square

Now let $c_n = \sqrt{a_n^2 - b_n^2}$, $n \geq 0$, be given by (3.2ii). We show that

$$(7.10) \quad c_n = \theta_+^2(q^{2^n}), \quad n \geq 0.$$

To this end, compute

$$\begin{aligned} \theta_0^2(q) - \theta_0^2(q^2) &= \sum_{n=0}^{\infty} \sigma(n)q^n - \sum_{n=0}^{\infty} \sigma(2n)q^{2n} \\ &= \sum_{n \text{ odd}} \sigma(n)q^n \\ &= \sum_{\substack{i, j \in \mathbb{Z} \\ i^2 + j^2 \text{ odd}}} q^{i^2 + j^2}. \end{aligned}$$

Now, $i^2 + j^2$ is odd iff $i + j$ is odd iff $i - j$ is odd, in which case $k = (j + i - 1)/2$ and $\ell = (j - i - 1)/2$ are integers. Solving, since $i = k - \ell$, and $j = k + \ell + 1$, the last sum equals

$$\begin{aligned} \theta_0^2(q) - \theta_0^2(q^2) &= \sum_{k, \ell \in \mathbb{Z}} q^{(k-\ell)^2 + (k+\ell+1)^2} \\ &= \sum_{k, \ell \in \mathbb{Z}} (q^2)^{(k^2+k+1/4) + (\ell^2+\ell+1/4)} \\ &= \left[\sum_{-\infty}^{\infty} (q^2)^{k^2+k+1/4} \right]^2 = \theta_+^2(q^2). \end{aligned}$$

Hence,

$$(7.11) \quad \theta_0^2(q^2) + \theta_+^2(q^2) = \theta_0^2(q).$$

Adding (7.6) and (7.11) leads to

$$\theta_0^2(q^2) - \theta_+^2(q^2) = \theta_-^2(q).$$

Multiplying the last two equations and recalling (7.7) leads to

$$(7.12) \quad \theta_0^4(q^2) = \theta_-^4(q^2) + \theta_+^4(q^2).$$

Now replacing q^2 by q^{2^n} in (7.12) leads to (7.10) since $a_n^2 = b_n^2 + c_n^2$. This establishes (7.10). \square

From §5.3, we know that $c_n \rightarrow 0$. Let us compute the rate at which this happens. It turns out that the decay rate is exponential, in the sense that

$$(7.13) \quad \lim_{n \nearrow \infty} \frac{1}{2^n} \log(c_n) = \frac{1}{2} \log q.$$

To see this, let us denote, for clarity, $2^n = N$. Then, by (7.10),

$$\frac{1}{N} \log(c_n) = \frac{1}{N} \log(\theta_+^2(q^N))$$

$$\begin{aligned}
&= \frac{2}{N} \log (2q^{N/4} + 2q^{9N/4} + 2q^{25N/4} + \dots) \\
&= \frac{2}{N} [\log 2 + (N/4) \log q + \log (1 + q^{2N} + q^{6N} + \dots)].
\end{aligned}$$

Hence, since $q^N \rightarrow 0$, we obtain (7.13). This computation should be compared with Exercise 7 of §5.5. Now, (3.21) says

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} \log \left(\frac{a_n}{c_n} \right) = \frac{\pi}{2} Q \left(\frac{b}{a} \right).$$

Inserting $a_n \rightarrow \theta_0(0) = 1$ and (7.13) into this equation and recalling $a = \theta_0^2(q)$, $b = \theta_-^2(q)$, we obtain

$$(7.14) \quad -\frac{1}{\pi} \log q = Q \left(\frac{\theta_-^2(q)}{\theta_0^2(q)} \right).$$

Here, $Q(x) = M(1, x)/M(1, x')$. Solving for q , we obtain the following sharpening of the previous theorem.

Theorem. Suppose that (a, b) satisfies $M(a, b) = 1$, $a > 1 > b > 0$, and let $q \in (0, 1)$ be such that $(a, b) = (\theta_0^2(q), \theta_-^2(q))$. Then,²⁸

$$q = e^{-\pi Q(b/a)}. \quad \square$$

Now, let $q = e^{-\pi s}$, $s > 0$, and set $\theta_0(s) = \theta_0(e^{-\pi s})$, $\theta_{\pm}(s) = \theta_{\pm}(e^{-\pi s})$. Then, (7.14) can be written

$$(7.15) \quad s = Q \left(\frac{\theta_-^2(s)}{\theta_0^2(s)} \right) = \frac{M(1, \theta_-^2(s)/\theta_0^2(s))}{M(1, (\theta_-^2(s)/\theta_0^2(s))')}, \quad s > 0.$$

Replacing s by $1/s$ in this last equation and using $1/Q(x) = Q(x')$, we obtain

$$s = Q \left(\left(\frac{\theta_-^2(1/s)}{\theta_0^2(1/s)} \right)' \right) = Q \left(\frac{\theta_+^2(1/s)}{\theta_0^2(1/s)} \right), \quad s > 0.$$

Here, we use (7.12) to show that $(\theta_-^2/\theta_0^2)' = \theta_+^2/\theta_0^2$. Equating the last two expressions for s and using the strict monotonicity of Q (Exercise 8 of §5.3), we arrive at

$$(7.16) \quad \frac{\theta_-^2(s)}{\theta_0^2(s)} = \frac{\theta_+^2(1/s)}{\theta_0^2(1/s)}, \quad s > 0.$$

Now, we can derive the theta functional equation (7.1), as follows.

$$s\theta_0^2(s) = \frac{s\theta_0^2(s)}{M(\theta_0^2(s), \theta_-^2(s))} \quad [(7.9)]$$

²⁸Now, go back and look at (3.16).

$$\begin{aligned}
&= \frac{s}{M(1, \theta_-^2(s)/\theta_0^2(s))} && \text{[homogeneity]} \\
&= \left[M\left(1, (\theta_-^2(s)/\theta_0^2(s))'\right) \right]^{-1} && [(7.15)] \\
&= \left[M\left(1, \theta_+^2(s)/\theta_0^2(s)\right) \right]^{-1} \\
&= \left[M\left(1, \theta_-^2(1/s)/\theta_0^2(1/s)\right) \right]^{-1} && [(7.16)] \\
&= \frac{\theta_0^2(1/s)}{M(\theta_0^2(1/s), \theta_-^2(1/s))} && \text{[homogeneity]} \\
&= \theta_0^2(1/s). && [(7.9)]
\end{aligned}$$

Since $\theta_0(s) = \theta(s)$, this completes the derivation of (7.1). \square Combining (7.1) with (7.16), we obtain the companion functional equation

$$(7.17) \quad \sqrt{s}\theta_-(e^{-\pi s}) = \theta_+(e^{-\pi/s}), \quad s > 0.$$

Exercises 5.7

1. Show that θ_0 and θ_+ are strictly increasing functions of $(0, 1)$ onto $(1, \infty)$.
2. Derive (7.17).
3. Show that θ_- is a strictly decreasing function of $(0, 1)$ onto $(0, 1)$. (Use (7.17) to compute $\theta_-(1-)$.)
4. Compute $\sigma(n)$ for $n = 11, 12, 13, 14, 15$. Show that $\sigma(4n-1) = 0$ for $n \geq 1$.
5. ★ Let $\psi(t, x) = \sum_{n=1}^{\infty} e^{-n^2\pi t} \cos(nx)$, $t > 0$, $x \in \mathbb{R}$. Show that²⁹ ψ satisfies the *heat equation*

$$\frac{\partial \psi}{\partial t} = \pi \frac{\partial^2 \psi}{\partial x^2}.$$

§5.8 Riemann's Zeta Function

In this section, we study the *Riemann zeta function*

$$\zeta(x) = \sum_{n=1}^{\infty} \frac{1}{n^x} = 1 + \frac{1}{2^x} + \frac{1}{3^x} + \dots, \quad x > 1,$$

and we discuss

- the behavior of ζ near $x = 1$,
- the extension of the domains of definition of Γ and ζ ,

²⁹The notation ψ for this function, which occurs in Riemann's 1859 paper, became so well known that the physicist Schrödinger, in deriving (1926) his famous quantum-mechanical analog of the heat equation, denoted the solution of his equation by ψ .

- the functional equation,
- the values of the zeta function at the nonpositive integers,
- the Euler product, and
- primes in arithmetic progressions.

Most of the results in this section are due to Euler. Nevertheless, ζ is associated with Riemann because, as Riemann showed, the subject really takes off only after x is allowed to range in the complex plane.

We already know that $\zeta(x)$ is smooth for $x > 1$ (Exercise 8 of §5.4) and $\zeta(1) = \zeta(1+) = \infty$ (Exercise 12 of §5.1). We say that f is *asymptotically equal* to g as $x \rightarrow a$, and we write $f(x) \sim g(x)$, as $x \rightarrow a$, if $f(x)/g(x) \rightarrow 1$, as $x \rightarrow a$ (compare with $a_n \sim b_n$ §5.5).

Theorem.

$$\zeta(x) \sim \frac{1}{x-1}, \quad x \rightarrow 1+.$$

We have to show that $(x-1)\zeta(x) \rightarrow 1$ as $x \rightarrow 1+$. Multiply $\zeta(x)$ by 2^{-x} to get

$$(8.1) \quad 2^{-x}\zeta(x) = \frac{1}{2^x} + \frac{1}{4^x} + \frac{1}{6^x} + \dots, \quad x > 1.$$

Then,

$$(8.2) \quad \left(1 - \frac{2}{2^x}\right)\zeta(x) = 1 - \frac{1}{2^x} + \frac{1}{3^x} - \frac{1}{4^x} + \dots, \quad x > 1.$$

Now, by the Leibnitz test, the series in (8.2) converges for $x > 0$, equals $\log 2$ at $x = 1$ (Exercise 17 of §3.6), and

$$\lim_{x \rightarrow 1} \left(1 - \frac{1}{2^x} + \frac{1}{3^x} - \frac{1}{4^x} + \dots\right) = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

by the dominated convergence theorem for series (see (2.18)). On the other hand, by l'Hopital's rule,

$$\lim_{x \rightarrow 1} \frac{1 - 2^{1-x}}{x-1} = \log 2.$$

Thus,

$$\begin{aligned} \lim_{x \rightarrow 1+} (x-1)\zeta(x) &= \lim_{x \rightarrow 1+} \frac{x-1}{1-2^{1-x}} \cdot \left(1 - \frac{2}{2^x}\right)\zeta(x) \\ &= \frac{1}{\log 2} \left(1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots\right) = 1. \quad \square \end{aligned}$$

Thus, $\zeta(x)$ and $1/(x-1)$ are asymptotically equal as $x \rightarrow 1+$. Nevertheless, it may be possible that the difference $\zeta(x) - 1/(x-1)$ still goes to infinity. For

example, x^2 and x^2+x are asymptotically equal as $x \rightarrow \infty$ but $(x^2+x)-x^2 \rightarrow \infty$, as $x \rightarrow \infty$. In fact, for $\zeta(x)$, we show that this does not happen.

Theorem.

$$(8.3) \quad \lim_{x \rightarrow 1+} \left[\zeta(x) - \frac{1}{x-1} \right] = \gamma,$$

where γ is Euler's constant.

To see this, use Exercise 8 of §5.1 and $\Gamma(x) = (x-1)\Gamma(x-1)$ to get, for $x > 1$,

$$\begin{aligned} \left[\zeta(x) - \frac{1}{x-1} \right] \Gamma(x) &= \zeta(x)\Gamma(x) - \Gamma(x-1) \\ &= \int_0^\infty \frac{t^{x-1}}{e^t - 1} dt - \int_0^\infty e^{-t} t^{x-2} dt \\ &= \int_0^\infty t^{x-1} \left(\frac{1}{e^t - 1} - \frac{1}{te^t} \right) dt. \end{aligned}$$

Applying the dominated convergence theorem (Exercise 1),

$$(8.4) \quad \lim_{x \rightarrow 1+} \left(\zeta(x) - \frac{1}{x-1} \right) \Gamma(x) = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{te^t} \right) dt.$$

But the integral in (8.4) is not easy to evaluate directly, so, we abandon this approach. Instead, we use the following identity.

Theorem (Sawtooth Formula). *Let $f : (1, \infty) \rightarrow \mathbf{R}$ be differentiable, decreasing, and nonnegative, and suppose that f' is continuous. Then,*

$$(8.5) \quad \sum_{n=1}^{\infty} f(n) = \int_1^{\infty} f(t) dt + \int_1^{\infty} (1 + [t] - t)[-f'(t)] dt.$$

Here $[t]$ is the greatest integer $\leq t$, and $0 \leq 1 + [t] - t \leq 1$ is the sawtooth function (Figure 2.6 in §2.3). To get (8.5), break up the following integrals of nonnegative functions and integrate by parts:

$$\begin{aligned} &\int_1^{\infty} f(t) dt + \int_1^{\infty} (1 + [t] - t)[-f'(t)] dt \\ &= \sum_{n=1}^{\infty} \left\{ \int_n^{n+1} f(t) dt + \int_n^{n+1} (1 + n - t)[-f'(t)] dt \right\} \\ &= \sum_{n=1}^{\infty} \left\{ \int_n^{n+1} f(t) dt + (1 + n - t)(-f(t)) \Big|_n^{n+1} - \int_n^{n+1} f(t) dt \right\} \\ &= \sum_{n=1}^{\infty} f(n). \quad \square \end{aligned}$$

Now, insert $f(t) = 1/t^x$ in (8.5), and evaluate the integral obtaining

$$(8.6) \quad \zeta(x) = \frac{1}{x-1} + x \int_1^\infty \frac{1 + [t] - t}{t^{x+1}} dt, \quad x > 1.$$

We wish to take the limit $x \rightarrow 1+$. Since the integrand is dominated by $1/t^2$ when $x > 1$, the dominated convergence theorem applies. Hence,

$$\lim_{x \rightarrow 1+} \left[\zeta(x) - \frac{1}{x-1} \right] = \int_1^\infty \frac{1 + [t] - t}{t^2} dt.$$

But

$$\begin{aligned} \int_1^\infty \frac{1 + [t] - t}{t^2} dt &= \lim_{N \nearrow \infty} \sum_{n=1}^N \int_n^{n+1} \frac{1 + n - t}{t^2} dt \\ &= \lim_{N \nearrow \infty} \sum_{n=1}^N \left(-\frac{n+1}{t} - \log t \right) \Big|_n^{n+1} \\ &= \lim_{N \nearrow \infty} \left[1 + \frac{1}{2} + \cdots + \frac{1}{N} - \log(N+1) \right] = \gamma. \end{aligned}$$

This completes the derivation of (8.3). \square

The series expression for $\zeta(x)$ is valid only when $x > 1$. Below we extend the domain of ζ to $x < 1$. To this end, we seek an alternate expression for ζ . Because the expression that we will find for ζ involves Γ , first, we extend $\Gamma(x)$ to $x < 0$.

Recall (§5.1) that the gamma function is smooth and positive on $(0, \infty)$. Hence, its reciprocal $L = 1/\Gamma$ is smooth there. Since $\Gamma(x+1) = x\Gamma(x)$, $L(x) = xL(x+1)$. But $xL(x+1)$ is smooth on $x > -1$. Hence, we can use this last equation to define $L(x)$ on $x > -1$ as a smooth function vanishing at $x = 0$. Similarly, we can use $L(x) = xL(x+1) = x(x+1)L(x+2)$ to define $L(x)$ on $x > -2$ as a smooth function, vanishing at $x = 0$ and $x = -1$. Continuing in this manner, the reciprocal $L = 1/\Gamma$ of the gamma function extends to a smooth function on \mathbb{R} , vanishing at $x = 0, -1, -2, \dots$. From this, it follows that Γ itself extends to a smooth function on $\mathbb{R} \setminus \{0, -1, -2, \dots\}$. Moreover (Exercise 3),

$$(8.7) \quad \Gamma(x) \sim \frac{(-1)^n}{n!} \cdot \frac{1}{x+n}, \quad x \rightarrow -n, \quad n \geq 0.$$

To obtain an alternate expression for ζ , start with

$$\psi(t) = \sum_{n=1}^{\infty} e^{-n^2\pi t}, \quad t > 0,$$

use Exercise 9 of §5.1, and substitute $1/t$ for t to get, for $x > 1$,

$$\begin{aligned} \pi^{-x/2} \Gamma(x/2) \zeta(x) &= \int_0^\infty \psi(t) t^{x/2-1} dt \\ &= \int_0^1 \psi(t) t^{x/2-1} dt + \int_1^\infty \psi(t) t^{x/2-1} dt \end{aligned}$$

$$(8.8) \quad = \int_1^\infty \left[\psi\left(\frac{1}{t}\right) t^{-x/2-1} + \psi(t) t^{x/2-1} \right] dt.$$

Now, by the theta functional equation (7.2),

$$\psi\left(\frac{1}{t}\right) = \frac{\sqrt{t}}{2} + \sqrt{t}\psi(t) - \frac{1}{2}.$$

So, (8.8) leads to

$$\begin{aligned} \pi^{-x/2} \Gamma(x/2) \zeta(x) &= \int_1^\infty t^{-x/2-1} \left(\frac{\sqrt{t}}{2} - \frac{1}{2} \right) dt \\ &\quad + \int_1^\infty \psi(t) [t^{(1-x)/2} + t^{x/2}] \frac{dt}{t}. \end{aligned}$$

Evaluating the first integral (recall $x > 1$), we obtain our alternate expression for ζ ,

$$(8.9) \quad \pi^{-x/2} \Gamma(x/2) \zeta(x) = \frac{1}{x(x-1)} + \int_1^\infty \psi(t) [t^{(1-x)/2} + t^{x/2}] \frac{dt}{t},$$

valid for $x > 1$.

Let us analyze (8.9). The integral on the right is a smooth function of x on \mathbb{R} (Exercise 4). Hence, the right side of (8.9) is a smooth function of $x \neq 0, 1$. On the other hand, $\pi^{-x/2}$ is smooth and positive, and $L(x/2) = 1/\Gamma(x/2)$ is smooth on all of \mathbb{R} . Thus, (8.9) can be used to define $\zeta(x)$ as a smooth function on $x \neq 0, 1$. Moreover, since $1/x\Gamma(x/2) = 1/2\Gamma(x/2+1)$, (8.9) can be used to define $\zeta(x)$ as a smooth function near $x = 0$. Now, by (8.7), $\Gamma(x/2)(x+2n) \rightarrow 2(-1)^n/n!$ as $x \rightarrow -2n$. So, multiplying (8.9) by $x+2n$ and sending $x \rightarrow -2n$ yields

$$(-1)^n \pi^n (2/n!) \lim_{x \rightarrow -2n} \zeta(x) = \begin{cases} 0 & \text{if } n > 0, \\ -1 & \text{if } n = 0. \end{cases}$$

Thus, $\zeta(-2n) = 0$ for $n > 0$ and $\zeta(0) = -1/2$. Now, the zeta function $\zeta(x)$ is defined for all $x \neq 1$. We summarize the results.

Theorem. *The zeta function can be defined, for all $x \neq 1$, as a smooth function. Moreover, $\zeta(-2n) = 0$ for $n \geq 1$, and $\zeta(0) = -1/2$. \square*

Now the right side of (8.9) is unchanged under the substitution $x \mapsto (1-x)$. This, immediately, leads to the following.

Theorem (Zeta Functional Equation). *Set $\xi(x) = \pi^{-x/2} \Gamma(x/2) \zeta(x)$. Then,*

$$(8.10) \quad \xi(x) = \xi(1-x), \quad x \neq \dots, -4, -2, 0, 1, 3, 5, \dots \quad \square$$

Since we obtained $\zeta(2n)$, $n \geq 1$, in §5.6, plugging in $x = 2n$, $n \geq 1$, into (8.10) leads us to the following.

Theorem. For all $n \geq 1$,

$$\zeta(1 - 2n) = -\frac{B_{2n}}{2n}.$$

For example, $\zeta(-1) = -1/12$. We leave the derivation of this as Exercise 5. \square Now, we know $\zeta(x)$ at all nonpositive integers and all positive even integers. Even though this result is over 200 years old, similar expressions for $\zeta(2n + 1)$, $n \geq 1$, have not yet been computed. In particular, very little is known about $\zeta(3)$.

We turn to our last topic, the prime numbers. Before proceeding, the reader may wish to review the Exercises in §1.3. That there is a connection between the zeta function and the prime numbers was discovered by Euler in 1737.

Theorem (Euler Product). For all $x > 1$,

$$\sum_{n=1}^{\infty} \frac{1}{n^x} = \prod_p \left(1 - \frac{1}{p^x}\right)^{-1}.$$

Here the product³⁰ is over all primes.

This follows from the fundamental theorem of arithmetic (Exercise 15 of §1.3). More specifically, from (8.1),

$$(8.11) \quad \zeta(x) \left(1 - \frac{1}{2^x}\right) = 1 + \frac{1}{3^x} + \frac{1}{5^x} + \cdots = 1 + \sum_{2 \nmid n} \frac{1}{n^x}, \quad x > 1,$$

where $2 \nmid n$ means 2 does not divide n and $n > 1$. Similarly, subtracting $1/3^x$ times (8.11) from (8.11) yields

$$\zeta(x) \left(1 - \frac{1}{2^x}\right) \left(1 - \frac{1}{3^x}\right) = 1 + \sum_{\substack{2 \nmid n \\ 3 \nmid n}} \frac{1}{n^x}, \quad x > 1.$$

Continuing in this manner,

$$(8.12) \quad \zeta(x) \prod_{n=1}^N \left(1 - \frac{1}{p_n^x}\right) = 1 + \sum_{p_1, p_2, \dots, p_N \nmid n} \frac{1}{n^x}, \quad x > 1,$$

where p_1, p_2, \dots, p_N are the first N primes. But $p_1, p_2, \dots, p_N \nmid n$ and $n > 1$ implies $n > N$. Hence, the series on the right side of (8.12) is no greater than $\sum_{n=N+1}^{\infty} 1/n^x$, which goes to zero as $N \nearrow \infty$. \square

Euler used his product to establish the infinitude of primes, as follows: Since (Exercise 9)

$$(8.13) \quad 0 < -\log(1 - a) \leq 2a, \quad 0 < a \leq 1/2,$$

³⁰This equality and its derivation are valid whether or not there are infinitely many primes.

it follows that

$$(8.14) \quad \log \zeta(x) = \sum_p -\log \left(1 - \frac{1}{p^x}\right) \leq 2 \sum_p \frac{1}{p^x}, \quad x > 1.$$

Now, as $x \rightarrow 1+$, $\zeta(x) \rightarrow \infty$, hence, $\log \zeta(x) \rightarrow \infty$. On the other hand, $\sum_p 1/p^x \rightarrow \sum_p 1/p$, as $x \searrow 1$, by the monotone convergence theorem. We have arrived at the following.

Theorem. *There are infinitely many primes. In fact, there are enough of them so that*

$$\sum_p \frac{1}{p} = \infty. \quad \square$$

Our last topic is the infinitude of primes in arithmetic progressions. Let a and b be naturals. An *arithmetic progression* is a subset of \mathbb{N} of the form $a\mathbb{N} + b = \{a + b, 2a + b, 3a + b, \dots\}$. Apart from 2, every prime is either in $4\mathbb{N} + 1$ or $4\mathbb{N} + 3$. Note that $p \in a\mathbb{N} + b$ iff a divides $p - b$, which we write as $a \mid p - b$. Here is Euler's result on primes in arithmetic progressions.

Theorem. *There are infinitely many primes in $4\mathbb{N} + 1$ and in $4\mathbb{N} + 3$. In fact, there are enough of them so that*

$$\sum_{4 \mid p-1} \frac{1}{p} = \infty$$

and

$$\sum_{4 \mid p-3} \frac{1}{p} = \infty.$$

We proceed by analogy with the preceding derivation. Instead of relating $\sum_p 1/p^x$ to $\log \zeta(x)$, now, we relate $\sum_{4 \mid p-1} 1/p^x$ and $\sum_{4 \mid p-3} 1/p^x$ to $\log L_1(x)$ and $\log L_3(x)$, where

$$L_1(x) = \sum_{4 \mid n-1} \frac{1}{n^x}, \quad x > 1,$$

and

$$L_3(x) = \sum_{4 \mid n-3} \frac{1}{n^x}, \quad x > 1.$$

By comparison with $\zeta(x)$, the series $L_1(x)$ and $L_3(x)$ are finite for $x > 1$. To make the analogy clearer, we define $\chi_1 : \mathbb{N} \rightarrow \mathbb{R}$ and $\chi_3 : \mathbb{N} \rightarrow \mathbb{R}$ by setting

$$\chi_1(n) = \begin{cases} 1, & n \in 4\mathbb{N} + 1, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$\chi_3(n) = \begin{cases} 1, & n \in 4\mathbb{N} + 3, \\ 0, & \text{otherwise.} \end{cases}$$

Then,

$$L_1(x) = \sum_{n=1}^{\infty} \frac{\chi_1(n)}{n^x}, \quad L_3(x) = \sum_{n=1}^{\infty} \frac{\chi_3(n)}{n^x}, \quad x > 1.$$

Proceeding further, the next step was to obtain an identity of the form

$$(8.15) \quad \sum_{n=1}^{\infty} \frac{\chi(n)}{n^x} = \prod_p \left[1 - \frac{\chi(p)}{p^x} \right]^{-1}, \quad x > 1.$$

Denote the series in (8.15) by $L(x, \chi)$. Thus, $L_1(x) = L(x, \chi_1)$ and $L_3(x) = L(x, \chi_3)$. When $\chi = \chi_1$ or $\chi = \chi_3$, however, (8.15) is false, and for a very good reason: χ_1 and χ_3 are not multiplicative.

Theorem. *Suppose that $\chi : \mathbb{N} \rightarrow \mathbb{R}$ is bounded and multiplicative, i.e., suppose that $\chi(mn) = \chi(m)\chi(n)$ for all $m, n \in \mathbb{N}$. Then, (8.15) holds.*

The derivation of this is completely analogous to the previous case and involves inserting factors of χ in (8.12). \square Having arrived at this point, Euler bypassed the failure of (8.15) for χ_1, χ_3 by considering, instead,

$$\chi_+ = \chi_1 + \chi_3,$$

and

$$\chi_- = \chi_1 - \chi_3.$$

Then, $\chi_+(n)$ is 1 or 0 according to whether n is odd or even, $L(x, \chi_+)$ is given by (8.11), χ_- is given by

$$\chi_-(n) = \begin{cases} 1, & 4 \mid n - 1, \\ -1, & 4 \mid n - 3, \\ 0, & n \text{ even,} \end{cases}$$

and

$$L(x, \chi_-) = 1 - \frac{1}{3^x} + \frac{1}{5^x} - \frac{1}{7^x} + \dots$$

But this is an alternating, hence, convergent series for $x > 0$ by the Leibnitz test, and $L(1, \chi_-) > 0$. Moreover (Exercise 11), by the dominated convergence theorem,

$$(8.16) \quad \lim_{x \rightarrow 1} L(x, \chi_-) = L(1, \chi_-) > 0.$$

Now, the key point is that χ_+ and χ_- are multiplicative (Exercise 10), and, hence, (8.15) holds with $\chi = \chi_{\pm}$.

Proceeding, as in (8.14), and taking the log of (8.15) with $\chi = \chi_+$, we obtain

$$\log L(x, \chi_+) \leq 2 \sum_p \frac{\chi_+(p)}{p^x}, \quad x > 1.$$

Since, by (8.11), $\lim_{x \rightarrow 1+} L(x, \chi_+) = \infty$, sending $x \rightarrow 1+$, we conclude that

$$(8.17) \quad \lim_{x \rightarrow 1+} \sum_p \frac{\chi_+(p)}{p^x} = \infty.$$

Turning to χ_- , we claim it is enough to show that $\sum_p \chi_-(p)p^{-x}$ remains bounded as $x \rightarrow 1+$. Indeed, assuming this claim, we have

$$\begin{aligned} \sum_{4 \nmid p-1} \frac{1}{p} &= \sum_p \frac{\chi_1(p)}{p} \\ &= \lim_{x \searrow 1} \sum_p \frac{\chi_1(p)}{p^x} \\ &= \lim_{x \searrow 1} \frac{1}{2} \left[\sum_p \frac{\chi_+(p)}{p^x} + \sum_p \frac{\chi_-(p)}{p^x} \right] = \infty \end{aligned}$$

by the monotone convergence theorem, the claim, and (8.17). This is the first half of the theorem. Similarly,

$$\begin{aligned} \sum_{4 \nmid p-3} \frac{1}{p} &= \sum_p \frac{\chi_3(p)}{p} \\ &= \lim_{x \searrow 1} \sum_p \frac{\chi_3(p)}{p^x} \\ &= \lim_{x \searrow 1} \frac{1}{2} \left[\sum_p \frac{\chi_+(p)}{p^x} - \sum_p \frac{\chi_-(p)}{p^x} \right] = \infty. \end{aligned}$$

This is the second half of the theorem.

To complete the derivation, we establish the claim using

$$(8.18) \quad |-\log(1-a) - a| \leq a^2, \quad |a| \leq 1/2,$$

which follows from the power series for log (Exercise 9). Taking the log of (8.15) and using (8.18) with $a = \chi_-(p)/p^x$, we obtain

$$\left| \log L(x, \chi_-) - \sum_p \frac{\chi_-(p)}{p^x} \right| \leq \sum_p \frac{1}{p^{2x}} \leq \sum_{n=1}^{\infty} \frac{1}{n^2}, \quad x > 1.$$

By (8.16), $\log L(x, \chi_-) \rightarrow \log L(1, \chi_-)$ and, so, remains bounded as $x \rightarrow 1+$. Since, by the last equation, $\sum_p \chi_-(p)p^{-x}$ differs from $\log L(x, \chi_-)$ by a bounded quantity, this establishes the claim. \square

One hundred years later (1840), Dirichlet showed³¹ there are infinitely many primes in any arithmetic progression $aN + b$, as long as a and b have no common factor.

Exercises 5.8

1. Use the dominated convergence theorem to derive (8.4). (Exercise 15 of §3.4.)
2. Show that

$$\gamma = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{te^t} \right) dt.$$

3. Derive (8.7).
4. Dominate ψ by a geometric series to obtain $\psi(t) \leq ce^{-\pi t}$, $t \geq 1$, where $c = 1/(1 - e^{-\pi})$. Use this to show that the integral in (8.9) is a smooth function of x in \mathbb{R} .
5. Use (8.10) and the values $\zeta(2n)$, $n \geq 1$, obtained in §5.6, to show that $\zeta(1 - 2n) = -B_{2n}/2n$, $n \geq 1$.
6. Let $I(x)$ denote the integral in (8.6). Show that $I(x)$ is finite, smooth for $x > 0$, and satisfies $I'(x) = -(x + 1)I(x + 1)$. Compute $I(2)$.
7. Use (8.9) to check that $(x - 1)\zeta(x)$ is smooth and positive on $(1 - \delta, 1 + \delta)$ for δ small enough. Then, differentiate $\log[(x - 1)\zeta(x)]$ for $1 < x < 1 + \delta$ using (8.6). Conclude that

$$\lim_{x \rightarrow 1} \left[\frac{\zeta'(x)}{\zeta(x)} + \frac{1}{x - 1} \right] = \gamma.$$

8. Differentiate the log of (8.10) to obtain $\zeta'(0) = -\frac{1}{2} \log(2\pi)$. (Use the previous exercise, Exercise 10 of §5.5, and Exercise 13 of §5.6.)
9. Derive (8.13) and (8.18) using the power series for $\log(1 + a)$.
10. Show that $\chi_\pm : \mathbb{N} \rightarrow \mathbb{R}$ are multiplicative.
11. Derive (8.16) using the dominated convergence theorem. (Group the terms in pairs, and use the mean value theorem to show that $a^{-x} - b^{-x} \leq x/a^{x+1}$, $b > a > 0$.)

§5.9 The Euler–Maclaurin Formula

Given a smooth function $f : \mathbb{R} \rightarrow \mathbb{R}$, can we find a smooth function $g : \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$(9.1) \quad f(x + 1) - f(x) = \int_x^{x+1} g(t) dt, \quad x \in \mathbb{R}?$$

³¹By replacing χ_\pm by the characters χ of the cyclic group $(\mathbb{Z}/a\mathbb{Z})^*$.

By the fundamental theorem, the answer is yes: $g = f'$, and g is also smooth. However, this is not the only solution because $g = f' + p'$ solves (9.1) for any smooth periodic p , i.e., for any smooth p satisfying $p(x+1) = p(x)$ for all $x \in \mathbb{R}$.

The starting point for the Euler–Maclaurin formula is to ask the same question but with the left side in (9.1) modified. More precisely, given a smooth function $f : \mathbb{R} \rightarrow \mathbb{R}$, can we find a smooth function $g : \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$(9.2) \quad f(x+1) = \int_x^{x+1} g(t) dt, \quad x \in \mathbb{R}?$$

Note that $g = 1$ works when $f = 1$. We call a g satisfying (9.2) an *Euler–Maclaurin derivative of f* .

It turns out the answer is yes and (9.2) is always solvable. To see this, let q denote a primitive of g . Then, (9.2) becomes $f(x) = q(x) - q(x-1)$. Conversely, suppose that

$$(9.3) \quad f(x) = q(x) - q(x-1), \quad x \in \mathbb{R},$$

for some smooth q . Then, it is easy to check that $g = q'$ works in (9.2). Thus, given f , (9.2) is solvable for some smooth g iff (9.3) is solvable for some smooth q .

In fact, it turns out that (9.3) is always solvable. Note, however, that q solves (9.3) iff $q+p$ solves (9.3), where p is any periodic smooth function, i.e., $p(x+1) = p(x)$, $x \in \mathbb{R}$. So, the solution is not unique.

To solve (9.3), assume, in addition, that $f(x) = 0$ for $x < -1$, and define q by

$$(9.4) \quad q(x) = \begin{cases} f(x), & x \leq 0, \\ f(x) + f(x-1), & 0 \leq x \leq 1, \\ f(x) + f(x-1) + f(x-2), & 1 \leq x \leq 2, \\ \text{and so on.} \end{cases}$$

Then, q is well defined and smooth on \mathbb{R} and (9.3) holds (Exercise 1). Thus, (9.3) is solvable when f vanishes on $(-\infty, -1)$. Similarly, (9.3) is solvable when f vanishes on $(1, \infty)$ (Exercise 2).

To obtain the general case, we write $f = f_+ + f_-$, where $f_+ = 0$ on $(-\infty, -1)$ and $f_- = 0$ on $(1, \infty)$. Then, $q = q_+ + q_-$ solves (9.3) for f if q_{\pm} solve (9.3) for f_{\pm} . Thus, to complete the solution of (9.3), all we need do is construct f_{\pm} .

Because we require f_+ , f_- to be smooth, it is not immediately clear this can be done. To this end, we deal, first, with the special case $f = 1$, i.e., we construct ϕ_{\pm} smooth and satisfying $\phi_+ = 0$ on $(-\infty, -1)$, $\phi_- = 0$ on $(1, \infty)$, and $1 = \phi_+ + \phi_-$ on \mathbb{R} .

To construct ϕ_{\pm} , let h denote the function in Exercise 3 of §3.4. Then, $h : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, $h = 0$ on \mathbb{R}^- and $h > 0$ on \mathbb{R}^+ . Set

$$\phi_+(x) = \frac{h(1+x)}{h(1-x) + h(1+x)}, \quad x \in \mathbb{R},$$

and

$$\phi_-(x) = \frac{h(1-x)}{h(1-x) + h(1+x)}, \quad x \in \mathbb{R}.$$

Since $h(1-x) + h(1+x) > 0$ on all of \mathbb{R} , ϕ_{\pm} are smooth with $\phi_+ = 0$ on $(-\infty, -1)$, $\phi_- = 0$ on $(1, \infty)$, and $\phi_+ + \phi_- = 1$ on all of \mathbb{R} .

Now, for smooth f , we may set $f_{\pm} = f\phi_{\pm}$, yielding $f = f_+ + f_-$ on all of \mathbb{R} . Thus, (9.3) is solvable for all smooth f . Hence, (9.2) is solvable for all smooth f .

Theorem. *Every smooth $f : \mathbb{R} \rightarrow \mathbb{R}$ has a smooth Euler–Maclaurin derivative $g : \mathbb{R} \rightarrow \mathbb{R}$. \square*

Our main interest is to obtain a useful formula for an Euler–Maclaurin derivative g of f . To explain this, we denote $f' = Df$, $f'' = D^2f$, $f''' = D^3f$, and so on. Then, any polynomial in D makes sense. For example, $D^3 + 2D^2 - D + 5$ is the *differential operator* that associates the smooth function f with the smooth function

$$(D^3 + 2D^2 - D + 5)f = f''' + 2f'' - f' + 5f.$$

More generally, we may consider infinite linear combinations of powers of D . For example,

$$\begin{aligned} (9.5) \quad e^{tD}f(c) &= \left(1 + tD + \frac{t^2D^2}{2!} + \frac{t^3D^3}{3!} + \dots\right)f(c) \\ &= f(c) + tf'(c) + \frac{t^2}{2!}f''(c) + \frac{t^3}{3!}f'''(c) + \dots \end{aligned}$$

may sum to $f(c+t)$, since this is the Taylor series, but, for general smooth f , diverges from $f(c+t)$. When f is a polynomial of degree d , (9.5) does sum to $f(c+t)$. Hence, $e^{tD}f(c) = f(c+t)$. In fact, in this case, any power series in D applied to f is another polynomial of degree d , as $D^n f = 0$ for $n > d$. For example, if B_n , $n \geq 0$, are the Bernoulli numbers (§5.6), then,

$$(9.6) \quad \tau(D) = 1 + B_1D + \frac{B_2}{2!}D^2 + \frac{B_4}{4!}D^4 + \dots$$

may be applied to any polynomial $f(x)$ of degree d . The result $\tau(D)f(x)$, then, obtained is another polynomial of, at most, the same degree.

If $\tau(D)$ is applied to $f(x) = e^{ax}$ for a real, we obtain

$$(9.7) \quad \tau(D)e^{ax} = \tau(a)e^{ax},$$

where $\tau(a)$ is the Bernoulli function of §5.6,

$$\tau(a) = 1 + B_1a + \frac{B_2}{2!}a^2 + \frac{B_4}{4!}a^4 + \dots$$

Thus, (9.7) is valid only on the interval of convergence of the power series for $\tau(a)$.

Let $c(a)$ be a power series. To compute the effect of $c(D)$ on a product $e^{ax} f(x)$, where f is a polynomial, note that

$$D[e^{ax}x] = axe^{ax} + e^{ax} = e^{ax}(ax + 1)$$

by the product rule. Repeating this with D^2, D^3, \dots ,

$$D^n[e^{ax}x] = e^{ax}(a^n x + na^{n-1}).$$

Taking linear combinations, we conclude that

$$c(D)(e^{ax}x) = e^{ax}[c(a)x + c'(a)] = \frac{\partial}{\partial a}[c(a)e^{ax}].$$

Thus, $c(D)(e^{ax}x)$ is well defined for a in the interval of convergence of $c(a)$. Similarly, one checks that $c(D)(e^{ax}x^n)$ is well defined for any $n \geq 1$ and a in the interval of convergence (Exercise 3) and

$$(9.8) \quad c(D)(e^{ax}x^n) = \frac{\partial^n}{\partial a^n}[c(a)e^{ax}].$$

We call a smooth function *elementary* if it is a product $e^{ax} f(x)$ of an exponential e^{ax} with a in the interval of convergence of $\tau(a)$ and a polynomial $f(x)$. In particular, any polynomial is elementary. Note that $\tau(D)f$ is elementary whenever f is elementary.

Theorem. *For all elementary functions f , $\tau(D)f$ is an Euler–Maclaurin derivative, i.e.,*

$$(9.9) \quad f(x+1) = \int_x^{x+1} \tau(D)f(t) dt, \quad x \in \mathbb{R}.$$

To derive (9.9), start with $f(x) = e^{ax}$. If $a = 0$, (9.9) is clearly true. If $a \neq 0$, then, by (9.7), (9.9) is equivalent to

$$e^{a(x+1)} = \int_x^{x+1} \tau(a)e^{at} dt = \tau(a) \cdot \frac{e^{a(x+1)} - e^{ax}}{a}$$

which is true since $\tau(a) = a/(1 - e^{-a})$. Thus,

$$e^{a(x+1)} = \int_x^{x+1} \tau(a)e^{at} dt, \quad a \in \mathbb{R}, x \in \mathbb{R}.$$

Now, apply $\partial^n/\partial a^n$ to both sides of this last equation, differentiate under the integral sign, and use (9.8). You obtain (9.9) with $f(x) = e^{ax}x^n$. By linearity, one obtains (9.9) for any elementary function f . \square

Given $a < b$ with $a, b \in \mathbb{Z}$, insert $x = a, a+1, a+2, \dots, b-1$ in (9.9), and sum the resulting equations to get the following.

Theorem (Euler–Maclaurin). *For $a < b$ in \mathbb{Z} and any elementary function f ,*

$$(9.10) \quad \sum_{a < n \leq b} f(n) = \int_a^b \tau(D)f(t) dt. \quad \square$$

The derivation of this is a triviality. The depth lies in the usefulness of the result. This arises from the fact that (9.10) equates a discrete sum of f on the left with a continuous sum of a related function $\tau(D)f$ on the right. Indeed, the reader may take it as an article of faith that any mathematical phenomenon³² relating the discrete and the continuous is important.

By inserting $a = 0$, $b = \infty$, and $f(t) = 1/(x+t)^2$, x fixed, in (9.10), one can derive a sharpening of Stirling's approximation (§5.5), the *Stirling series* for $\log \Gamma(x)$. Since this f is not elementary, here, one obtains a divergent series $\tau(D)f$. Instead of starting with (9.10), however, it will be quicker for us to derive the Stirling series from the identity (Exercise 12 of §5.6)

$$(9.11) \quad \frac{d^2}{dx^2} \log \Gamma(x) = \int_0^\infty e^{-xt} \tau(t) dt, \quad x > 0.$$

But, first, we discuss asymptotic expansions.

Let f and g be defined near $x = c$. We say that f is *big oh of* g , as $x \rightarrow c$, and we write $f(x) = O(g(x))$, as $x \rightarrow c$, if the ratio $f(x)/g(x)$ is bounded for $x \neq c$ in some interval about c . If $c = \infty$, then, we require that $f(x)/g(x)$ be bounded for x sufficiently large. For example, $f(x) \sim g(x)$, as $x \rightarrow c$, implies $f(x) = O(g(x))$ and $g(x) = O(f(x))$, as $x \rightarrow c$. We write $f(x) = g(x) + O(h(x))$ to mean $f(x) - g(x) = O(h(x))$. Note that $f(x) = O(h(x))$ and $g(x) = O(h(x))$ imply $f(x) + g(x) = O(h(x))$ or, what is the same, $O(h(x)) + O(h(x)) = O(h(x))$.

We say that

$$(9.12) \quad f(x) \approx a_0 + a_1x + a_2x^2 + \dots,$$

is an *asymptotic expansion of f at zero* if

$$(9.13) \quad f(x) = a_0 + a_1x + \dots + a_nx^n + O(x^{n+1}), \quad x \rightarrow 0,$$

for all $n \geq 0$. Here, there is no assumption regarding the convergence of the series in (9.12). Although the Taylor series of a smooth function may diverge, we have the following.

Theorem. *If f is smooth in an interval about 0, then,*

$$f(x) \approx f(0) + f'(0)x + \frac{1}{2!}f''(0)x^2 + \frac{1}{3!}f'''(0)x^3 + \frac{1}{4!}f^{(4)}(0)x^4 + \dots$$

is an asymptotic expansion at zero.

This follows from Taylor's theorem. If $a_n = f^{(n)}(0)/n!$, then, from §3.4,

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \frac{1}{(n+1)!}h_{n+1}(x)x^{n+1},$$

with h_{n+1} continuous on an interval about 0, hence, bounded near 0. \square

For example,

$$e^{-1/|x|} \approx 0, \quad x \rightarrow 0,$$

³²Is light composed of particles or waves?

since $t = 1/x$ implies $e^{-1/|x|}/x^n = e^{-|t|}t^n \rightarrow 0$ as $t \rightarrow \pm\infty$.

Actually, we will need asymptotic expansions at ∞ . Let f be defined near ∞ , i.e., for x sufficiently large. We say that

$$(9.14) \quad f(x) \approx a_0 + \frac{a_1}{x} + \frac{a_2}{x^2} + \dots,$$

is an *asymptotic expansion of f at infinity* if

$$f(x) = a_0 + \frac{a_1}{x} + \dots + \frac{a_n}{x^n} + O\left(\frac{1}{x^{n+1}}\right), \quad x \rightarrow \infty,$$

for all $n \geq 0$. For example, $e^{-x} \approx 0$ as $x \rightarrow \infty$, since $e^{-x} = O(x^{-n})$, as $x \rightarrow \infty$, for all $n \geq 0$. Here is the Stirling series.

Theorem (Stirling). As $x \rightarrow \infty$,

$$(9.15) \quad \begin{aligned} \log \Gamma(x) - \left[\left(x - \frac{1}{2} \right) \log x - x \right] \\ \approx \frac{1}{2} \log(2\pi) + \frac{B_2}{2x} + \frac{B_4}{4 \cdot 3x^3} + \frac{B_6}{6 \cdot 5x^5} + \dots \end{aligned}$$

Note that, ignoring the terms with Bernoulli numbers, this result reduces to Stirling's approximation §5.5. Moreover, note that, because this is an expression for $\log \Gamma(x)$ and not $\Gamma(x)$, the terms involving the Bernoulli numbers are measures of *relative error*. Thus, the principal error term $B_2/2x = 1/12x$ equals $1/1200$ for $x = 100$ which agrees with the relative error of .08% found in Exercise 2 of §5.5.

To derive (9.15), we will use (9.11) and replace $\tau(t)$ by its Bernoulli series to obtain

$$(9.16) \quad \frac{d^2}{dx^2} \log \Gamma(x) \approx \frac{1}{x} + \frac{1}{2x^2} + \frac{B_2}{x^3} + \frac{B_4}{x^5} + \dots, \quad x \rightarrow \infty.$$

Then, we integrate this twice to get (9.15).

First, we show that the portion of the integral in (9.11) over $(1, \infty)$ has no effect on the asymptotic expansion (9.16). Fix $n \geq 0$. To this end, note that

$$0 \leq \int_1^\infty e^{-xt} \tau(t) dt \leq \frac{1}{1 - e^{-1}} \int_1^\infty e^{-xt} t dt = \frac{1}{1 - e^{-1}} \cdot \frac{1 + x}{x^2} \cdot e^{-x}$$

for $x > 0$. Thus,

$$\int_1^\infty e^{-xt} \tau(t) dt = O\left(\frac{1}{x^{n+1}}\right), \quad x \rightarrow \infty.$$

Since τ is smooth at zero and the Bernoulli series is the Taylor series of τ , by Taylor's theorem (§3.4), there is a continuous $h_n : \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$\tau(t) = B_0 + \frac{B_1}{1!}t + \frac{B_2}{2!}t^2 + \dots + \frac{B_{n-1}}{(n-1)!}t^{n-1} + \frac{h_n(t)}{n!}t^n, \quad t \in \mathbb{R}.$$

Then (Exercise 4),

$$\int_0^1 e^{-xt} h_n(t) \frac{t^n}{n!} dt = \frac{1}{x^{n+1}} \int_0^x e^{-t} h_n(t/x) \frac{t^n}{n!} dt = O\left(\frac{1}{x^{n+1}}\right)$$

since $h_n(x)$ is bounded for $0 \leq x \leq 1$. Similarly (Exercise 5),

$$\int_1^\infty e^{-xt} \frac{t^k}{k!} dt = O\left(\frac{1}{x^{n+1}}\right), \quad x \rightarrow \infty, k \geq 0.$$

Now, insert all this into (9.11), and use

$$\int_0^\infty e^{-xt} \frac{t^n}{n!} dt = \frac{1}{x^{n+1}}$$

to get, for fixed $n \geq 0$,

$$\begin{aligned} \int_0^\infty e^{-xt} \tau(t) dt &= \int_0^1 e^{-xt} \tau(t) dt + \int_1^\infty e^{-xt} \tau(t) dt \\ &= \int_0^1 e^{-xt} \tau(t) dt + O\left(\frac{1}{x^{n+1}}\right) \\ &= \sum_{k=0}^{n-1} B_k \int_0^1 e^{-xt} \frac{t^k}{k!} dt + \int_0^1 e^{-xt} h_n(t) \frac{t^n}{n!} dt + O\left(\frac{1}{x^{n+1}}\right) \\ &= \sum_{k=0}^{n-1} B_k \int_0^1 e^{-xt} \frac{t^k}{k!} dt + O\left(\frac{1}{x^{n+1}}\right) \\ &= \sum_{k=0}^{n-1} \frac{B_k}{x^{k+1}} - \sum_{k=0}^{n-1} B_k \int_1^\infty e^{-xt} \frac{t^k}{k!} dt + O\left(\frac{1}{x^{n+1}}\right) \\ &= \sum_{k=0}^{n-1} \frac{B_k}{x^{k+1}} + O\left(\frac{1}{x^{n+1}}\right). \end{aligned}$$

Since $B_0 = 1$, $B_1 = 1/2$, and this is true for all $n \geq 0$, this derives (9.16).

To get (9.15), let $f(x) = (\log \Gamma(x))'' - (1/x) - (1/2x^2)$. Then, by (9.16),

$$(9.17) \quad f(x) = \frac{B_2}{x^3} + \frac{B_4}{x^5} + \cdots + \frac{B_n}{x^{n+1}} + O\left(\frac{1}{x^{n+2}}\right).$$

Since the right side of this last equation is integrable over (x, ∞) for any $x > 0$, so is f . Since $-\int_x^\infty f(t) dt$ is a primitive of f , we obtain

$$\int_x^\infty f(t) dt = -[\log \Gamma(x)]' + \log x - \frac{1}{2x} - A$$

for some constant A . So, integrating both sides of (9.17) over (x, ∞) leads to

$$-[\log \Gamma(x)]' + \log x - \frac{1}{2x} - A = \frac{B_2}{2x^2} + \frac{B_4}{4x^4} + \cdots + \frac{B_n}{nx^n} + O\left(\frac{1}{x^{n+1}}\right).$$

Similarly, integrating this last equation over (x, ∞) leads to

$$\begin{aligned} \log \Gamma(x) - x \log x + x + \frac{1}{2} \log x + Ax - B \\ = \frac{B_2}{2 \cdot 1x} + \frac{B_4}{4 \cdot 3x^3} + \cdots + \frac{B_n}{n \cdot (n-1)x^{n-1}} + O\left(\frac{1}{x^n}\right). \end{aligned}$$

Noting that the right side of this last equation vanishes as $x \rightarrow \infty$, inserting $x = n$ in the left side, and comparing with Stirling's approximation in §5.5, we conclude that $A = 0$ and $B = \log(2\pi)/2$, thus, obtaining (9.15). \square

Exercises 5.9

1. Show that q , as defined by (9.4), is well defined, smooth on \mathbf{R} , and satisfies (9.3), when f vanishes on $(-\infty, -1)$.
2. Find a smooth q solving (9.3), when f vanishes on $(1, \infty)$.
3. Let c be a power series with radius of convergence R . Show that $c(D)(e^{ax}x^n)$ is well defined for $|a| < R$ and $n \geq 0$ and satisfies

$$c(D)(e^{ax}x^n) = \frac{\partial^n}{\partial a^n}[c(a)e^{ax}].$$

4. Show that $\int_0^1 e^{-xt} f(t)t^n dt = O\left(\frac{1}{x^{n+1}}\right)$ for any continuous bounded $f : (0, 1) \rightarrow \mathbf{R}$ and $n \geq 0$.
5. For all $n \geq 0$ and $p > 0$, show that $\int_1^\infty e^{-xt} t^p dt \approx 0$, as $x \rightarrow \infty$.
6. Show that the Stirling series in (9.15) cannot converge anywhere. (If it did converge at $a \neq 0$, then, the Bernoulli series would converge on all of \mathbf{R} .)

Appendix

Solutions

Below, 1.1.1 means the solution to the first Exercise in the first section of Chapter 1.

A.1 Chapter 1

Exercises 1.1

1.1.1. Let A denote the set of all cats (alive or dead), and, for each $a \in A$, let $f(a)$ and $g(a)$ denote the mother and father of a , respectively. Then, $f : A \rightarrow A$ and $g : A \rightarrow A$, and $f(g(a))$ is a 's father's mother whereas $g(f(a))$ is a 's mother's father.

1.1.2. For the first equation in De Morgan's law, choose a in the right side $\bigcap_{n=1}^{\infty} A_n^c$. Then, $a \in A_n^c$ for all $n \geq 1$. Hence, $a \notin A_n$ for all $n \geq 1$. Hence, a is not in $\bigcup_{n=1}^{\infty} A_n$, i.e., a is in the left side. Thus,

$$\left(\bigcup_{n=1}^{\infty} A_n \right)^c \supset \bigcap_{n=1}^{\infty} A_n^c.$$

If a is in the left side $\left(\bigcup_{n=1}^{\infty} A_n \right)^c$, then, a is not in $\bigcup_{n=1}^{\infty} A_n$. Hence, a is not in A_n for any $n \geq 1$. Hence, $a \in A_n^c$ for every $n \geq 1$. Hence, a is in $\bigcap_{n=1}^{\infty} A_n^c$. Thus,

$$\left(\bigcup_{n=1}^{\infty} A_n \right)^c \subset \bigcap_{n=1}^{\infty} A_n^c.$$

We conclude that

$$\left(\bigcup_{n=1}^{\infty} A_n \right)^c = \bigcap_{n=1}^{\infty} A_n^c.$$

To obtain the second equation in De Morgan's law, replace A_n by A_n^c in the first equation. Since $(A^c)^c = A$, we obtain

$$\left(\bigcup_{n=1}^{\infty} A_n^c \right)^c = \bigcap_{n=1}^{\infty} A_n.$$

Taking the complement of each side yields the second equation

$$\bigcup_{n=1}^{\infty} A_n^c = \left(\bigcap_{n=1}^{\infty} A_n \right)^c.$$

1.1.3. Suppose that $f : A \rightarrow B$ is invertible. This means there is a $g : B \rightarrow A$ with $g(f(a)) = a$ for all a and $f(g(b)) = b$ for all $b \in B$. If $f(a) = f(a')$, then, $a = g(f(a)) = g(f(a')) = a'$. Hence, f is injective. If $b \in B$, then, $a = g(b)$ satisfies $f(a) = b$. Hence, f is surjective. We conclude that f is bijective. Conversely, if f is bijective, for each $b \in B$, let $g(b)$ denote the unique element $a \in A$ satisfying $f(a) = b$. Then, by construction, $f(g(b)) = b$. Moreover, since a is unique element of A mapping to $f(a)$, we also have $a = g(f(a))$. Thus, g is the inverse of f . Hence, f is invertible.

1.1.4. Suppose that $g_1 : B \rightarrow A$ and $g_2 : B \rightarrow A$ are inverses of f . Then, $f(g_1(b)) = b = f(g_2(b))$ for all $b \in B$. Since f is injective, this implies $g_1(b) = g_2(b)$ for all $b \in B$, i.e., $g_1 = g_2$.

Exercises 1.2

1.2.1. $a0 = a0 + (a - a) = (a0 + a) - a = (a0 + a1) - a = a(0 + 1) - a = a1 - a = a - a = 0$. This is not the only way.

1.2.2. The number 1 satisfies $1a = a1 = a$ for all a . If $1'$ also satisfied $1'b = b1' = b$ for all b , then, choosing $a = 1'$ and $b = 1$ yields $1 = 11' = 1'$. Hence, $1 = 1'$. Now, suppose that a has two negatives, one called $-a$ and one called b . Then, $a + b = 0$, so, $-a = -a + 0 = -a + (a + b) = (-a + a) + b = 0 + b = b$. Hence, $b = -a$, so, a has a unique negative. If $a \neq 0$ has two reciprocals, one called $1/a$ and one called b , $ab = 1$, so, $1/a = (1/a)1 = (1/a)(ab) = [(1/a)a]b = 1b = b$.

1.2.3. Since $a + (-a) = 0$, a is the (unique) negative of $-a$ which means $-(-a) = a$. Also $a + (-1)a = 1a + (-1)a = [1 + (-1)]a = 0a = 0$, so, $(-1)a$ is the negative of a or $-a = (-1)a$.

1.2.4. By the ordering property, $a > 0$, and $b > 0$ implies $ab > 0$. If $a < 0$ and $b > 0$, then, $-a > 0$. Hence, $(-a)b > 0$, so, $ab = [-(-a)]b = (-1)(-a)b = -((-a)b) < 0$. Thus, negative times positive is negative. If $a < 0$ and $b < 0$, then, $-b > 0$. Hence, $a(-b) < 0$. Hence, $ab = a(-(-b)) = a(-1)(-b) = (-1)a(-b) = -(a(-b)) > 0$. Thus, negative times negative is positive. Also, $1 = 11 > 0$ whether $1 > 0$ or $1 < 0$ ($1 \neq 0$ is part of the ordering property).

1.2.5. $a < b$ implies $b - a > 0$ which implies $(b + c) - (a + c) > 0$ or $b + c > a + c$. Also $c > 0$ implies $c(b - a) > 0$ or $bc - ac > 0$ or $bc > ac$. If $a < b$ and $b < c$, then, $b - a$ and $c - b$ are positive. Hence, $c - a = (c - b) + (b - a)$ is positive or $c > a$. Multiplying $a < b$ by $a > 0$ and by $b > 0$ yields $aa < ab$ and $ab < bb$. Hence, $aa < bb$.

1.2.6. If $0 \leq a \leq b$ we know, from above, that $aa \leq bb$. Conversely, if $aa \leq bb$ then, we cannot have $a > b$ because applying 5 with the roles of a, b reversed yields $aa > bb$, a contradiction. Hence, $a \leq b$ iff $aa \leq bb$.

1.2.7.

- By definition, $\inf A \leq x$ for all $x \in A$. Hence, $-\inf A \geq -x$ for all $x \in A$. Hence, $-\inf A \geq y$ for all $y \in -A$. Hence, $-\inf A \geq \sup(-A)$. Conversely, $\sup(-A) \geq y$ for all $y \in -A$, or $\sup(-A) \geq -x$ for all $x \in A$, or $-\sup(-A) \leq x$ for all $x \in A$. Hence, $-\sup(-A) \leq \inf A$ which implies $\sup(-A) \geq -\inf A$. Since we already know that $\sup(-A) \leq -\inf A$, we conclude that $\sup(-A) = -\inf A$. Now, replace A by $-A$ in the last equation. Since $-(-A) = A$, we obtain $\sup A = -\inf(-A)$ or $\inf(-A) = -\sup A$.
- Now, $\sup A \geq x$ for all $x \in A$, so, $(\sup A) + a \geq x + a$ for all $x \in A$. Hence, $(\sup A) + a \geq y$ for $y \in A + a$, so, $(\sup A) + a \geq \sup(A + a)$. In this last inequality, replace a by $-a$ and A by $A + a$ to obtain $[\sup(A + a)] - a \geq \sup(A + a - a)$, or $\sup(A + a) \geq (\sup A) + a$. Combining the two inequalities yields $\sup(A + a) = (\sup A) + a$. Replacing A and a by $-A$ and $-a$ yields $\inf(A + a) = (\inf A) + a$.
- Now, $\sup A \geq x$ for all $x \in A$. Since $c > 0$, $c \sup A \geq cx$ for all $x \in A$. Hence, $c \sup A \geq y$ for all $y \in cA$. Hence, $c \sup A \geq \sup(cA)$. Now, in this last inequality, replace c by $1/c$ and A by cA . We obtain $(1/c) \sup(cA) \geq \sup A$ or $\sup(cA) \geq c \sup A$. Combining the two inequalities yields $\sup(cA) = c \sup A$. Replacing A by $-A$ in this last equation yields $\inf(cA) = c \inf A$.

Exercises 1.3

1.3.1. Let S be the set of naturals n for which there are no naturals between n and $n + 1$. From the text, we know that $1 \in S$. Assume $n \in S$. Then, we claim $n + 1 \in S$. Indeed, suppose that $m \in \mathbb{N}$ satisfies $n + 1 < m < n + 2$. Then, $m \neq 1$, so, $m - 1 \in \mathbb{N}$ (see §1.3) satisfies $n < m - 1 < n + 1$, contradicting $n \in S$. Hence, $n + 1 \in S$, so, S is inductive. Since $S \subset \mathbb{N}$, we conclude that $S = \mathbb{N}$.

1.3.2. Fix $n \in \mathbb{N}$, and let $S = \{x \in \mathbb{R} : nx \in \mathbb{N}\}$. Then, $1 \in S$ since $n1 = n$. If $x \in S$, then, $nx \in \mathbb{N}$, so, $n(x + 1) = nx + n \in \mathbb{N}$ (since $\mathbb{N} + \mathbb{N} \subset \mathbb{N}$), so, $x + 1 \in S$. Hence, S is inductive. We conclude that $S \supset \mathbb{N}$ or $nm \in \mathbb{N}$ for all $m \in \mathbb{N}$.

1.3.3. Let S be the set of all naturals n such that the following holds: If $m > n$ and $m \in \mathbb{N}$, then, $m - n \in \mathbb{N}$. From the text, we know that $1 \in S$. Assume $n \in S$. We claim $n + 1 \in S$. Indeed, suppose that $m > n + 1$ and $m \in \mathbb{N}$. Then, $m - 1 > n$. Since $n \in S$, we conclude that $(m - 1) - n \in \mathbb{N}$ or $m - (n + 1) \in \mathbb{N}$. Hence, by definition of S , $n + 1 \in S$. Thus, S is inductive, so, $S = \mathbb{N}$. Thus, $m > n$ implies $m - n \in \mathbb{N}$. Since, for $m, n \in \mathbb{N}$, $m > n$, $m = n$, or $m < n$,

we conclude that $m - n \in \mathbf{Z}$, whenever $m, n \in \mathbf{N}$. If $n \in -\mathbf{N}$ and $m \in \mathbf{N}$, then, $-n \in \mathbf{N}$ and $m - n = m + (-n) \in \mathbf{N} \subset \mathbf{Z}$. If $m \in -\mathbf{N}$ and $n \in \mathbf{N}$, then, $-m \in \mathbf{N}$ and $m - n = -(n + (-m)) \in -\mathbf{N} \subset \mathbf{Z}$. If n and m are both in $-\mathbf{N}$, then, $-m$ and $-n$ are in \mathbf{N} . Hence, $m - n = -((-m) - (-n)) \in \mathbf{Z}$. If either m or n equals zero, then, $m - n \in \mathbf{Z}$. This shows that \mathbf{Z} is closed under subtraction.

1.3.4. If n is even and odd, then, $n + 1$ is even. Hence, $1 = (n + 1) - n$ is even, say $1 = 2k$ with $k \in \mathbf{N}$. But $k \geq 1$ implies $1 = 2k \geq 2$, which contradicts $1 < 2$. If $n = 2k$ is even and $m \in \mathbf{N}$, then, $nm = 2(km)$ is even. If $n = 2k - 1$ and $m = 2j - 1$ are odd, then, $nm = 2(2kj - k - j + 1) - 1$ is odd.

1.3.5. For all $n \geq 1$, we establish the claim: *If $m \in \mathbf{N}$ and there is a bijection between $\{1, \dots, n\}$ and $\{1, \dots, m\}$, then, $n = m$.* For $n = 1$, the claim is clearly true. Now, assume the claim is true for a particular n , and suppose that we have a bijection f between $\{1, \dots, n + 1\}$ and $\{1, \dots, m\}$ for some $m \in \mathbf{N}$. Then, by restricting f to $\{1, \dots, n\}$, we obtain a bijection g between $\{1, \dots, n\}$ and $\{1, \dots, k - 1, k + 1, \dots, m\}$, where $k = f(n + 1)$. Now, define $h(i) = i$ if $1 \leq i \leq k - 1$, and $h(i) = i - 1$ if $k + 1 \leq i \leq m$. Then, h is a bijection between $\{1, \dots, k - 1, k + 1, \dots, m\}$ and $\{1, \dots, m - 1\}$. Hence, $h \circ g$ is a bijection between $\{1, \dots, n\}$ and $\{1, \dots, m - 1\}$. By the inductive hypothesis, this forces $m - 1 = n$ or $m = n + 1$. Hence, the claim is true for $n + 1$. Hence, the claim is true, by induction, for all $n \geq 1$. Now, suppose that A is a set with n elements and with m elements. Then, there are bijections $f : A \rightarrow \{1, \dots, n\}$ and $g : A \rightarrow \{1, \dots, m\}$. Hence, $g \circ f^{-1}$ is a bijection from $\{1, \dots, n\}$ to $\{1, \dots, m\}$. Hence, $m = n$. This shows that the number of elements of a set is well defined. For the last part, suppose that A and B have n and m elements, respectively, and are disjoint. Let $f : A \rightarrow \{1, \dots, n\}$ and $g : B \rightarrow \{1, \dots, m\}$ be bijections, and let $h(i) = i + n$, $1 \leq i \leq m$. Then, $h \circ g : B \rightarrow \{n + 1, \dots, n + m\}$ is a bijection. Now, define $k : A \cup B \rightarrow \{1, \dots, n + m\}$ by setting $k(x) = f(x)$ if $x \in A$, $k(x) = h \circ g(x)$ if $x \in B$. Then, k is a bijection, establishing the number of elements of $A \cup B$ is $n + m$.

1.3.6. Let $A \subset \mathbf{R}$ be finite. By induction on the number of elements, we show that $\max A$ exists. If $A = \{a\}$, then, $a = \max A$. So, $\max A$ exists. Now, assume that every subset with n elements has a max. If A is a set with $n + 1$ elements and $a \in A$, then, $B = A \setminus \{a\}$ has n elements. Hence, $\max B$ exists. There are two cases: If $a \leq \max B$, then, $\max B = \max A$. Hence, $\max A$ exists. If $a > \max B$, then, $a = \max A$. Hence, $\max A$ exists. Since, in either case $\max A$ exists when $\#A = n + 1$, by induction, $\max A$ exists for all finite subsets A . Since $-A$ is finite whenever A is finite, $\min A$ exists by the reflection property.

1.3.7. Let $c = \sup S$. Since $c - 1$ is not an upper bound, choose $n \in S$ with $c - 1 < n \leq c$. If $c \in S$, then, $c = \max S$ and we are done. Otherwise, $c \notin S$, and $c - 1 < n < c$. Now, choose $m \in S$ with $c - 1 < n < m < c$ concluding that $m - n = (m - c) - (n - c)$ lies between 0 and 1, a contradiction. Thus, $c = \max S$.

1.3.8. If $yq \leq x$, then, $q \leq x/y$. So, $\{q \in \mathbb{N} : yq \leq x\}$ is nonempty and bounded above, hence, has a max, call it q . Let $r = x - yq$. Then, $r = 0$, or $r \in \mathbb{R}^+$. If $r \geq y$, then, $x - y(q+1) = r - y \geq 0$, so, $q+1 \in S$, contradicting the definition of q . Hence, $0 \leq r < y$.

1.3.9. Since $\mathbb{N} \times \mathbb{R}$ is an inductive subset of $\mathbb{R} \times \mathbb{R}$, $f \subset \mathbb{N} \times \mathbb{R}$. Now, $1 \in A$ since $(1, a) \in f$, and $x \in A$ implies $(x, y) \in f$ for some y implies $(x+1, ay) \in f$ implies $x+1 \in A$. Hence, A is an inductive subset of \mathbb{R} . We conclude that $A \supset \mathbb{N}$. Since $f \subset \mathbb{N} \times \mathbb{R}$, we obtain $A = \mathbb{N}$. To show that f is a function, we need to show that, for each n , there is a unique $y \in \mathbb{R}$ with $(n, y) \in f$. Suppose that this is not so, and let n be the smallest natural, such that (n, y) is in f for at least two distinct y 's. We use this n to discard a pair (n, y) from f obtaining a strictly smaller inductive subset \tilde{f} . To this end, if $n = 1$, let $\tilde{f} = f \setminus \{(1, y)\}$, where $y \neq a$. If $n > 1$, let $\tilde{f} = f \setminus \{(n, y)\}$ where $y \neq af(n-1)$. Since there are at least two y 's corresponding to n and n is the least natural with this property, $f(n-1)$ is uniquely determined, and there is at least one pair $(n, y) \in f$ with $y \neq af(n-1)$. Now, check that \tilde{f} is inductive, contradicting the fact that f was the smallest. Hence, f is a function. Moreover, by construction, $f(1) = a$, and $f(n+1) = af(n)$ for all n .

1.3.10. By construction, we know that $a^{n+1} = a^n a$ for all $n \geq 1$. Let S be the set of $m \in \mathbb{N}$, such that $a^{n+m} = a^n a^m$ for all $n \in \mathbb{N}$. Then, $1 \in S$. If $m \in S$, then, $a^{n+m} = a^n a^m$, so, $a^{n+(m+1)} = a^{(n+m)+1} = a^{n+m} a = a^n a^m a = a^n a^{m+1}$. Hence, $m+1 \in S$. Thus, S is inductive. Hence, $S = \mathbb{N}$. This shows that $a^{n+m} = a^n a^m$ for all $n, m \in \mathbb{N}$. If $n = 0$, then, $a^{n+m} = a^n a^m$ is clear, whereas $n < 0$ implies $a^{n+m} = a^{n+m} a^{-n} a^n = a^{n+m-n} a^n = a^m a^n$. This shows that $a^{n+m} = a^n a^m$ for $n \in \mathbb{Z}$ and $m \in \mathbb{N}$. Repeating this last argument with m , instead of n , we obtain $a^{n+m} = a^n a^m$ for all $n, m \in \mathbb{Z}$. We also establish the second part by induction on m with $n \in \mathbb{Z}$ fixed: If $m = 1$, the equation $(a^n)^m = a^{nm}$ is clear. So, assume it is true for m . Then, $(a^n)^{m+1} = (a^n)^m (a^n)^1 = a^{nm} a^n = a^{nm+n} = a^{n(m+1)}$. Hence, it is true for $m+1$, hence, by induction, for all $m \geq 1$. For $m = 0$, it is clearly true, whereas for $m < 0$, $(a^n)^m = 1/(a^n)^{-m} = 1/a^{-nm} = a^{nm}$.

1.3.11. Since $p \geq 2$, $1 < 2^n \leq p^n$ and $n < 2^n \leq p^n$ for all $n \geq 1$. If $p^k m = p^j q$ with $k < j$, then, $m = p^{j-k} q = p p^{j-k-1} q$ is divisible by p . On the other hand, if $k > j$, then, q is divisible by p . Hence, $p^k m = p^j q$ with m, q not divisible by p implies $k = j$. This establishes the uniqueness of the number of factors k . For existence, if n is not divisible by p , we take $k = 0$ and $m = n$. If n is divisible by p , then, $n_1 = n/p$ is a natural $< p^{n-1}$. If n_1 is not divisible by p , we take $k = 1$ and $m = n_1$. If n_1 is divisible by p , then, $n_2 = n_1/p$ is a natural $< p^{n-2}$. If n_2 is not divisible by p , we take $k = 2$ and $m = n_2$. If n_2 is divisible by p , we continue this procedure by dividing n_2 by p . Continuing in this manner, we obtain n_1, n_2, \dots naturals with $n_j < p^{n-j}$. Since this procedure ends in n steps at most, there is some k natural or 0 for which $m = n/p^k$ is not divisible by p and n/p^{k-1} is divisible by p .

1.3.12. We want to show that $S \supset \mathbb{N}$. If this is not so, then, $\mathbb{N} \setminus S$ would be nonempty, hence, would have a least element n . Thus, $k \in S$ for all naturals $k < n$. By the given property of S , we conclude that $n \in S$, contradicting $n \notin S$. Hence, $\mathbb{N} \setminus S$ is empty or $S \supset \mathbb{N}$.

1.3.13. Since $m \geq p$, $m = pq + r$ with $q \in \mathbb{N}$, $r \in \mathbb{N} \cup \{0\}$, and $0 \leq r < p$ (Exercise 8). If $r \neq 0$, multiplying by a yields $ra = ma - q(pa) \in \mathbb{N}$ since $ma \in \mathbb{N}$ and $pa \in \mathbb{N}$. Hence, $r \in S_a$ is less than p , contradicting $p = \min S_a$. Thus, $r = 0$, or p divides m .

1.3.14. With $a = n/p$, $p \in S_a$ since $pa = n \in \mathbb{N}$, and $m \in S_a$ since $ma = nm/p \in \mathbb{N}$. By the previous exercise, $\min S_a$ divides p . Since p is prime, $\min S_a = 1$, or $\min S_a = p$. In the first case, $1 \cdot a = a = n/p \in \mathbb{N}$, i.e., p divides n , whereas, in the second case, p divides m by the previous Exercise.

1.3.15. We use induction according to Exercise 12. For $n = 1$, the statement is true. Suppose that the statement is true for all naturals less than n . Then, either n is a prime or n is composite, $n = jk$ with $j > 1$ and $k > 1$. By the inductive hypothesis, j and k are products of primes, hence, so is n . Hence, in either case, n is a product of primes. To show that the decomposition for n is unique except for the order, suppose that $n = p_1 \dots p_r = q_1 \dots q_s$. By the previous Exercise, since p_1 divides the left, hence, the right side, p_1 divides one of the q_j 's. Since the q_j 's are prime, we conclude that p_1 equals q_j for some j . Hence, $n' = n/p_1 < n$ can be expressed as the product $p_2 \dots p_r$ and the product $q_1 \dots q_{j-1}q_{j+1} \dots q_s$. By the inductive hypothesis, these p 's and q 's must be identical except for the order. Hence, the result is true for n . By induction, the result is true for all naturals.

1.3.16. If the algorithm ends, then, $r_n = 0$ for some n . Solving backward, we see that $r_{n-1} \in \mathbb{Q}$, $r_{n-2} \in \mathbb{Q}$, etc. Hence, $x = r_0 \in \mathbb{Q}$. Conversely, if $x \in \mathbb{Q}$, then, all the remainders r_n are rationals. Now, given a rational $r = m/n$ with $n \in \mathbb{N}$ and $m \in \mathbb{Z}$ (in lowest terms), let $N(r) = m$ and $D(r) = n$. Then, $D(r) = D(r + n)$ for $n \in \mathbb{Z}$. Moreover, since $0 \leq r_n < 1$, $N(r_n) < D(r_n)$. Then,

$$N(r_n) = D\left(\frac{1}{r_n}\right) = D(q_{n+1} + r_{n+1}) = D(r_{n+1}) > N(r_{n+1}).$$

Thus, $N(r_0) > N(r_1) > N(r_2) > \dots$ is strictly decreasing. Hence, $N(r_n) = 1$ for some n , so, $r_{n+1} = 0$.

Exercises 1.4

1.4.1. Since $|x| = \max(x, -x)$, $x \leq |x|$. Also $-a < x < a$ is equivalent to $-x < a$ and $x < a$, hence, to $|x| = \max(x, -x) < a$. By definition of intersection, $x > -a$ and $x < a$ is equivalent to $\{x : x < a\} \cap \{x : x > -a\}$. Similarly, $|x| > a$ is equivalent to $x > a$ or $-x > a$, i.e., x lies in the union of $\{x : x > a\}$ and $\{x : x < -a\}$.

1.4.2. Clearly $|0| = 0$. If $x \neq 0$, then, $x > 0$ or $-x > 0$, hence, $|x| = \max(x, -x) > 0$. If $x > 0$ and $y > 0$, then, $|xy| = xy = |x||y|$. If $x > 0$ and

$y < 0$, then, xy is negative, so, $|xy| = -(xy) = x(-y) = |x| |y|$. Similarly, for the other two cases.

1.4.3. If $n = 1$, the inequality is true. Assume it is true for n . Then,

$$\begin{aligned} |a_1 + \cdots + a_n + a_{n+1}| &\leq |a_1 + \cdots + a_n| + |a_{n+1}| \\ &\leq |a_1| + \cdots + |a_n| + |a_{n+1}| \end{aligned}$$

by the triangle inequality and the inductive hypothesis. Hence, it is true for $n + 1$. By induction, it is true for all naturals n .

1.4.4. Assume first that $a > 1$. Let $S = \{x : x \geq 1 \text{ and } x^2 < a\}$. Since $1 \in S$, S is nonempty. Also $x \in S$ implies $x = x \cdot 1 \leq x^2 < a$, so, S is bounded above. Let $s = \sup S$. We claim that $s^2 = a$. Indeed, if $s^2 < a$, note that

$$\begin{aligned} \left(s + \frac{1}{n}\right)^2 &= s^2 + \frac{2s}{n} + \frac{1}{n^2} \\ &\leq s^2 + \frac{2s}{n} + \frac{1}{n} = s^2 + \frac{2s+1}{n} < a \end{aligned}$$

if $(2s+1)/n < a - s^2$, i.e., if $n > (2s+1)/(a - s^2)$. Since $s^2 < a$, $b = (2s+1)/(a - s^2)$ is a perfectly well defined, positive real. Since $\sup \mathbb{N} = \infty$, such a natural $n > b$ can always be found. This rules out $s^2 < a$. If $s^2 > a$, then, $b = (s^2 - a)/2s$ is positive. Hence, there is a natural n satisfying $1/n < b$ which implies $s^2 - 2s/n > a$. Hence,

$$\left(s - \frac{1}{n}\right)^2 = s^2 - \frac{2s}{n} + \frac{1}{n^2} > a,$$

so (by Exercise 6 of §1.2), $s - 1/n$ is an upper bound for S . This shows that s is not the *least* upper bound, contradicting the definition of s . Thus, we are forced to conclude that $s^2 = a$. Now, if $a < 1$, then, $1/a > 1$, and $1/\sqrt{1/a}$ is a positive square root of a . The square root is unique by Exercise 6 of §1.2.

1.4.5. By completing the square, x solves $ax^2 + bx + c = 0$ iff x solves $(x + b/2a)^2 = (b^2 - 4ac)/4a^2$. If $b^2 - 4ac < 0$, this shows that there are no solutions. If $b^2 - 4ac = 0$, this shows that $x = -b/2a$ is the only solution. If $b^2 - 4ac > 0$, take the square root of both sides to obtain $x + b/2a = \pm(\sqrt{b^2 - 4ac})/2a$.

1.4.6. The inequality $(1+a)^n \leq 1 + (2^n - 1)a$ is clearly true for $n = 1$. So, assume it is true for n . Then, $(1+a)^{n+1} = (1+a)^n(1+a) \leq (1 + (2^n - 1)a)(1+a) = 1 + 2^n a + (2^n - 1)a^2 \leq 1 + (2^{n+1} - 1)a$ since $0 \leq a^2 \leq a$. Hence, it is true for all n . The inequality $(1+b)^n \geq 1 + nb$ is true for $n = 1$. So, suppose that it is true for n . Since $1+b \geq 0$, then, $(1+b)^{n+1} = (1+b)^n(1+b) \geq (1+nb)(1+b) = 1 + (n+1)b + nb^2 \geq 1 + (n+1)b$. Hence, the inequality is true for all n .

1.4.7. If $0 \leq a < b$, then, $a^n < b^n$ is true for $n = 1$. If it is true for n , then, $a^{n+1} = a^n a < b^n a < b^n b = b^{n+1}$. Hence, by induction, it is true for all n . Hence, $0 \leq a \leq b$ implies $a^n \leq b^n$. If $a^n \leq b^n$ and $a > b$, then, by applying the previous, we obtain $a^n > b^n$, a contradiction. Hence, for $a, b \geq 0$, $a \leq b$ iff $a^n \leq b^n$. For

the second part, assume that $a > 1$, and let $S = \{x : x \geq 1 \text{ and } x^n \leq a\}$. Since $x \geq 1$, $x^{n-1} \geq 1$. Hence, $x = x1 \leq xx^{n-1} = x^n < a$. Thus, $s = \sup S$ exists. We claim that $s^n = a$. If $s^n < a$, then, $b = s^{n-1}(2^n - 1)/(a - s^n)$ is a well defined, positive real. Choose a natural $k > b$. Then, $s^{n-1}(2^n - 1)/k < a - s^n$. Hence, by Exercise 6,

$$\begin{aligned} \left(s + \frac{1}{k}\right)^n &= s^n \left(1 + \frac{1}{sk}\right)^n \\ &\leq s^n \left(1 + \frac{2^n - 1}{sk}\right) \\ &= s^n + \frac{s^{n-1}(2^n - 1)}{k} < a. \end{aligned}$$

Hence, $s + 1/k \in S$. Hence, s is not an upper bound for S , a contradiction. If $s^n > a$, $b = ns^{n-1}/(s^n - a)$ is a well defined, positive real, so, choose $k > b$. By Exercise 6,

$$\begin{aligned} \left(s - \frac{1}{k}\right)^n &= s^n \left(1 - \frac{1}{sk}\right)^n \\ &\geq s^n \left(1 - \frac{n}{sk}\right) \\ &= s^n - \frac{s^{n-1}n}{k} > a. \end{aligned}$$

Hence, by the first part of this Exercise, $s - 1/k$ is an upper bound for S . This shows that s is not the least upper bound, a contradiction. We conclude that $s^n = a$. Uniqueness follows from the first part of this Exercise.

1.4.8. If $t = k/(n\sqrt{2})$ is a rational p/q , then, $\sqrt{2} = (kq)/(np)$ is rational, a contradiction.

1.4.9. Let (b) denote the fractional part of b . If $a = 0$, the result is clear. If $a \neq 0$, the fractional parts of $a, 2a, 3a, \dots$, form a sequence in $[0, 1]$. Now, divide $[0, 1]$ into finitely many subintervals, each of length, at most, ϵ . Then, the fractional parts of at least two terms pa and qa , $p \neq q$, $p, q \in \mathbb{N}$, must lie in the same subinterval. Hence, $|(pa) - (qa)| < \epsilon$. Since $pa - (pa) \in \mathbb{Z}$, $qa - (qa) \in \mathbb{Z}$, we obtain $|(p - q)a - m| < \epsilon$ for some integer m . Choosing $n = p - q$, we obtain $|na - m| < \epsilon$.

1.4.10. The key issue here is that $|2n^2 - m^2| = n^2|f(m/n)|$ is a nonzero integer, since all the roots of $f(x) = 0$ are irrational. Hence, $|2n^2 - m^2| \geq 1$. But $2n^2 - m^2 = (n\sqrt{2} - m)(n\sqrt{2} + m)$, so, $|n\sqrt{2} - m| \geq 1/(n\sqrt{2} + m)$. Dividing by n , we obtain

$$(1.1) \quad \left|\sqrt{2} - \frac{m}{n}\right| \geq \frac{1}{(\sqrt{2} + m/n)n^2}.$$

Now, if $|\sqrt{2} - m/n| \geq 1$, the result we are trying to show is clear. So, let us assume that $|\sqrt{2} - m/n| \leq 1$. In this case, $\sqrt{2} + m/n = 2\sqrt{2} + (m/n - \sqrt{2}) \leq 2\sqrt{2} + 1$. Inserting this in the denominator of the right side of (1.1), the result follows.

1.4.11. By Exercise 5, the (real) roots of $f(x) = 0$ are $\pm a$. The key issue here is that $|m^4 - 2m^2n^2 - n^4| = n^4|f(m/n)|$ is a nonzero integer, since all the roots of $f(x) = 0$ are irrational. Hence, $n^4|f(m/n)| \geq 1$. But, by factoring, $f(x) = (x - a)g(x)$ with $g(x) = (x + a)(x^2 + \sqrt{2} - 1)$, so,

$$(1.2) \quad \left| a - \frac{m}{n} \right| \geq \frac{1}{n^4 g(m/n)}.$$

Now, there are two cases: If $|a - m/n| \geq 1$, we obtain $|a - m/n| \geq 1/n^4$ since $1 \geq 1/n^4$. If $|a - m/n| \leq 1$, then, $0 < m/n < 3$. So, from the formula for g , we obtain $0 < g(m/n) \leq 51$. Inserting this in the denominator of the right side of (1.2), the result follows with $c = 1/51$.

1.4.12. First, we verify the absolute value properties (1), (2), and (3) for $x, y \in \mathbb{Z}$. (1) is clear. For $x, y \in \mathbb{Z}$, let $x = 2^k p$ and $y = 2^j q$ with p, q odd. Then, $xy = 2^{j+k} pq$ with pq odd, establishing (2) for $x, y \in \mathbb{Z}$. For (3), let $i = \min(j, k)$ and note $x + y = 0$ or $x + y = 2^i r$ with r odd. In the first case, $|x + y|_2 = 0$, whereas, in the second, $|x + y|_2 = 2^{-i}$. Hence, $|x + y|_2 \leq 2^{-i} = \max(2^{-j}, 2^{-k}) = \max(|x|_2, |y|_2) \leq |x|_2 + |y|_2$. Now, using (2) for $x, y, z \in \mathbb{Z}$, $|zx|_2 = |z|_2|x|_2$ and $|zy|_2 = |z|_2|y|_2$. Hence, $|zx/zy|_2 = |zx|_2/|zy|_2 = |z|_2|x|_2/|z|_2|y|_2 = |x|_2/|y|_2 = |x/y|_2$. Hence, $|\cdot|_2$ is well defined on \mathbb{Q} . Now, using (1), (2), and (3) for $x, y \in \mathbb{Z}$, one checks (1), (2), and (3) for $x, y \in \mathbb{Q}$.

Exercises 1.5

1.5.1. If (a_n) is increasing, $\{a_n : n \geq 1\}$ and $\{a_n : n \geq N\}$ have the same sups. Similarly, for decreasing. If $a_n \rightarrow L$, then, $a_n^* \searrow L$. Hence, $a_{n+N}^* \searrow L$ and $a_{n*} \nearrow L$. Hence, $a_{(n+N)*} \nearrow L$. We conclude that $a_{n+N} \rightarrow L$.

1.5.2. If $a_n \nearrow L$, then, $L = \sup\{a_n : n \geq 1\}$, so, $-L = \inf\{-a_n : n \geq 1\}$, so, $-a_n \searrow -L$. Similarly, if $a_n \searrow L$, then, $-a_n \nearrow -L$. If $a_n \rightarrow L$, then, $a_n^* \searrow L$, so, $(-a_n)^* = -a_n^* \nearrow -L$, and $a_{n*} \nearrow L$, so, $(-a_n)^* = -a_{n*} \searrow -L$. Hence, $-a_n \rightarrow -L$.

1.5.3. First, if $A \subset \mathbb{R}^+$, let $1/A = \{1/x : x \in A\}$. Then, $\inf(1/A) = 0$ implies that, for all $c > 0$, there exists $x \in A$ with $1/x < 1/c$ or $x > c$. Hence, $\sup A = \infty$. Conversely, $\sup A = \infty$ implies $\inf(1/A) = 0$. If $\inf(1/A) > 0$, then, $c > 0$ is a lower bound for $1/A$ iff $1/c$ is an upper bound for A . Hence, $\sup A < \infty$ and $\inf(1/A) = 1/\sup A$. If $1/\infty$ is interpreted as 0, we obtain $\inf(1/A) = 1/\sup A$ in all cases. Applying this to $A = \{a_k : k \geq n\}$ yields $b_{n*} = 1/a_n^*, n \geq 1$. Moreover, A is bounded above iff $\inf(1/A) > 0$. Applying this to $A = \{b_{n*} : n \geq 1\}$ yields $\sup\{b_{n*} : n \geq 1\} = \infty$ since $a_n^* \searrow 0$. Hence, $b_n \rightarrow \infty$. For the converse, $b_n \rightarrow \infty$ implies $\sup\{b_{n*} : n \geq 1\} = \infty$. Hence, $\inf\{a_n^* : n \geq 1\} = 0$. Hence, $a_n \rightarrow 0$.

1.5.4. Since $k_n \geq n$, $a_{n*} \leq a_{k_n} \leq a_n^*$. Since $a_n \rightarrow L$ means $a_n^* \rightarrow L$ and $a_{n*} \rightarrow L$, the ordering property implies $a_{k_n} \rightarrow L$. Now assume (a_n) is increasing. Then, (a_{k_n}) is increasing. Suppose that $a_n \rightarrow L$ and $a_{k_n} \rightarrow M$. Since (a_n) is increasing and $k_n \geq n$, $a_{k_n} \geq a_n$, $n \geq 1$. Hence, by ordering, $M \geq L$. On the other hand, $\{a_{k_n} : n \geq 1\} \subset \{a_n : n \geq 1\}$. Since M and L are the sups of these sets, $M \leq L$. Hence, $M = L$.

1.5.5. From the text, we know that all but finitely many a_n lie in $(L - \epsilon, L + \epsilon)$, for any $\epsilon > 0$. Choosing $\epsilon = L$ shows that all but finitely many terms are positive.

1.5.6. The sequence $a_n = \sqrt{n+1} - \sqrt{n} = 1/(\sqrt{n+1} + \sqrt{n})$ is decreasing. Since $a_{n^2} \leq 1/n$ has limit zero, so does (a_n) . Hence, $a_n^* = a_n$ and $a_{n*} = 0 = a_* = a^*$.

1.5.7. We do only the case a^* finite. Since $a_n^* \rightarrow a^*$, a_n^* is finite for each $n \geq N$ beyond some $N \geq 1$. Now, $a_n^* = \sup\{a_k : k \geq n\}$, so, for each $n \geq 1$, we can choose $k_n \geq n$, such that $a_n^* \geq a_{k_n} > a_n^* - 1/n$. Then, the sequence (a_{k_n}) lies between (a_n^*) and $(a_n^* - 1/n)$, hence, converges to a^* . But (a_{k_n}) may not be a subsequence of (a_n) because the sequence (k_n) may not be strictly increasing. To take care of this, note that, since $k_n \geq n$, we can choose a subsequence (k_{j_n}) of (k_n) which is strictly increasing. Then, $(a_p : p = k_{j_n})$ is a subsequence of (a_n) converging to a^* . Similarly (or multiply by minuses), for a_* .

1.5.8. If $a_n \not\rightarrow L$, then, by definition, either $a^* \neq L$ or $a_* \neq L$. For definiteness, suppose that $a_* \neq L$. Then, from Exercise 7, there is a subsequence (a_{k_n}) converging to a_* . From §1.5, if $2\epsilon = |L - a_*| > 0$, all but finitely many of the terms a_{k_n} lie in the interval $(a_* - \epsilon, a_* + \epsilon)$. Since ϵ is chosen to be half the distance between a_* and L , this implies that these same terms lie outside the interval $(L - \epsilon, L + \epsilon)$. Hence, these terms form a subsequence as requested.

1.5.9. From Exercise 7, we know that x_* and x^* are limit points. If (x_{k_n}) is a subsequence converging to a limit point L , then, since $k_n \geq n$, $x_{n*} \leq x_{k_n} \leq x_n^*$ for all $n \geq 1$. By the ordering property for sequences, taking the limit yields $x_* \leq L \leq x^*$.

1.5.10. If $x_n \rightarrow L$, then, $x_* = x^* = L$. Since x_* and x^* are the smallest and the largest limit points, L must be the only one. Conversely, if there is only one limit point, then, $x_* = x^*$ since x_* and x^* are always limit points.

1.5.11. If $M < \infty$, then, for each $n \geq 1$, the number $M - 1/n$ is not an upper bound for the displayed set. Hence, there is an $x_n \in (a, b)$ with $f(x_n) > M - 1/n$. Since $f(x_n) \leq M$, we see that $f(x_n) \rightarrow M$, as $n \nearrow \infty$. If $M = \infty$, for each $n \geq 1$, the number n is not an upper bound for the displayed set. Hence, there is an $x_n \in (a, b)$ with $f(x_n) > n$. Then, $f(x_n) \rightarrow \infty = M$.

1.5.12. Note that

$$(1.3) \quad \frac{1}{2} \left(a + \frac{2}{a} \right) - \sqrt{2} = \frac{1}{2a} (a - \sqrt{2})^2 \geq 0, \quad a > 0.$$

Since $e_1 = 2 - \sqrt{2}$, $e_1 \geq 0$. By (1.3), $e_{n+1} \geq 0$ as soon as $e_n \geq 0$. Hence, $e_n \geq 0$ for all $n \geq 1$ by induction. Similarly, (1.3) with $a = d_n$ plugged in and $d_n \geq \sqrt{2}$, $n \geq 1$, yield $e_{n+1} \leq e_n^2/2\sqrt{2}$, $n \geq 1$.

1.5.13. If $f(a) = 1/(q + a)$, then, $|f(a) - f(b)| \leq f(a)f(b)|a - b|$. This implies (1). Now, (1) implies $|x - x_n| \leq x_n|x' - x'_n| \leq x_n x'_n |x'' - x''_n| \leq \dots$, where $x_n^{(k)}$ denotes x_n with k layers “peeled off.” Hence,

$$(1.4) \quad |x - x_n| \leq x_n x'_n x''_n \dots x_n^{(n-1)}, \quad n \geq 1.$$

Since $x_n^{(n-1)} = 1/q_n$, (1.4) implies (2). For (3), note that, since $q_n \geq 1$, $x \leq 1/[1 + 1/(q_2 + 1)] = (q_2 + 1)/(q_2 + 2)$. Let $a = (c + 1)/(c + 2)$. Now, if one of the q_k 's is bounded by c , (1.4) and (3) imply $|x - x_n| \leq a$, as soon as n is large enough, since all the factors in (1.4) are bounded by 1. Similarly, if two of the q_k 's are bounded by c , $|x - x_n| \leq a^2$, as soon as n is large enough. Continuing in this manner, we obtain (4). If $q_n \rightarrow \infty$, we are done, by (2). If not, then, there is a c , such that $q_k \leq c$ for infinitely many n . By (4), we conclude that the upper and lower limits of $(|x - x_n|)$ lie between 0 and a^N , for all $N \geq 1$. Since $a^N \rightarrow 0$, the upper and lower limits are 0. Hence, $|x - x_n| \rightarrow 0$.

Exercises 1.6

1.6.1. First, suppose that the decimal expansion of x is as advertised. Then, the fractional part of $10^{n+m}x$ is identical to the fractional part of $10^m x$. Hence, $x(10^{n+m} - 10^m) \in \mathbb{Z}$, or $x \in \mathbb{Q}$. Conversely, if $x = m/n \in \mathbb{Q}$, perform long division to obtain the digits: From Exercise 8 of §1.3, $10m = nd_1 + r_1$, obtaining the quotient d_1 and remainder r_1 . Similarly, $10r_1 = nd_2 + r_2$, obtaining the quotient d_2 and remainder r_2 . Similarly, $10r_2 = nd_3 + r_3$, obtaining d_3 and r_3 , and so on. Here, d_1, d_2, \dots are digits (since $0 < x < 1$), and r_1, r_2, \dots are zero or naturals less than n . At some point the remainders must start repeating, and, therefore, the digits also.

1.6.2. We assume $d_N > e_N$, and let $x = .d_1 d_2 \dots = .e_1 e_2 \dots$. If

$$y = .d_1 d_2 \dots d_{N-1} (e_N + 1) 00 \dots,$$

then, $x \geq y$. If

$$z = .e_1 e_2 \dots e_N 99 \dots,$$

then, $z \geq x$. Since $.99 \dots = 1$, $z = y$. Hence, $x = y$ and $x = z$. Since $x = y$, $d_N = e_N + 1$ and $d_j = 0$ for $j > N$. Since $x = z$, $e_j = 9$ for $j > N$. Clearly, this happens iff $10^N x \in \mathbb{Z}$.

1.6.3. Since $2^{1-1} \leq 1!$, the statement is true for $n = 1$. Assume it is true for n . Then, $2^{(n+1)-1} = 22^{n-1} \leq 2n! \leq (n+1)n! = (n+1)!$.

1.6.4. From Exercise 1.4.6, $(1 + b)^n \geq 1 + nb$ for $b \geq -1$. In this inequality, replace n by N and b by $-1/N(n+1)$ to obtain $[1 - 1/N(n+1)]^N \geq 1 - 1/(n+1) = n/(n+1)$. By Exercise 7 of §1.4, we may take N th roots of both sides, yielding

(1). (2) follows by multiplying (1) by $(n+1)^{1/N}$ and rearranging. If $a_n = 1/n^{1/N}$, then, by (2),

$$\begin{aligned} a_n - a_{n+1} &= \frac{(n+1)^{1/N} - n^{1/N}}{n^{1/N}(n+1)^{1/N}} \\ &\geq \frac{1}{N(n+1)^{(N-1)/N}n^{1/N}(n+1)^{1/N}} \\ &\geq \frac{1}{N(n+1)^{1+1/N}}. \end{aligned}$$

Summing over $n \geq 1$ yields (3).

1.6.5. Since $e_1 = 2 - \sqrt{2}$ and $e_{n+1} \leq e_n^2/2\sqrt{2}$, $e_2 \leq e_1^2/2\sqrt{2} = (3 - 2\sqrt{2})/\sqrt{2}$. Similarly,

$$\begin{aligned} e_3 &\leq \frac{e_2^2}{2\sqrt{2}} \\ &\leq \frac{(3 - 2\sqrt{2})^2}{4\sqrt{2}} \\ &= \frac{17 - 12\sqrt{2}}{4\sqrt{2}} \\ &= \frac{1}{4(17\sqrt{2} + 24)} \leq \frac{1}{100}. \end{aligned}$$

Now, assume the inductive hypothesis $e_{n+2} \leq 10^{-2^n}$. Then,

$$e_{(n+1)+2} = e_{n+3} \leq \frac{e_{n+2}^2}{2\sqrt{2}} \leq e_{n+2}^2 = (10^{-2^n})^2 = 10^{-2^{n+1}}.$$

Thus, the inequality is true by induction.

1.6.6. Since $[0, 2] = 2[0, 1]$, given $z \in [0, 1]$, we have to find $x \in C$ and $y \in C$ satisfying $x + y = 2z$. Let $z = .d_1d_2d_3\dots$. Then, for all $n \geq 1$, $2d_n$ is an even integer satisfying $0 \leq 2d_n \leq 18$. Thus, there are digits, zero or odd (i.e., 0, 1, 3, 5, 7, 9), $d'_n, d''_n, n \geq 1$, satisfying $d'_n + d''_n = 2d_n$. Now, set $x = .d'_1d'_2d'_3\dots$ and $y = .d''_1d''_2d''_3\dots$

Exercises 1.7

1.7.1. Since B is countable, there is a bijection $f : B \rightarrow \mathbb{N}$. Then, f restricted to A is a bijection between A and $f(A) \subset \mathbb{N}$. Thus, it is enough to show that $C = f(A)$ is countable or finite. If C is finite, we are done, so, assume C is infinite. Since $C \subset \mathbb{N}$, let $c_1 = \min C$, $c_2 = \min C \setminus \{c_1\}$, $c_3 = \min C \setminus \{c_1, c_2\}$, etc. Then, $c_1 < c_2 < c_3 < \dots$. Since C is infinite, $C_n = C \setminus \{c_1, \dots, c_n\}$ is not empty, allowing us to set $c_{n+1} = \min C_n$, for all $n \geq 1$. Since (c_n) is strictly increasing, we must have $c_n \geq n$ for $n \geq 1$. If $m \in C \setminus \{c_n : n \geq 1\}$, then, by

construction, $m \geq c_n$ for all $n \geq 1$, which is impossible. Thus, $C = \{c_n : n \geq 1\}$ and $g : \mathbb{N} \rightarrow C$ given by $g(n) = c_n, n \geq 1$, is a bijection.

1.7.2. With each rational r , associate the pair $f(r) = (m, n)$, where $r = m/n$ in lowest terms. Then, $f : \mathbb{Q} \rightarrow \mathbb{N}^2$ is an injection. Since \mathbb{N}^2 is countable and an infinite subset of a countable set is countable, so is \mathbb{Q} .

1.7.3. If $x \in \bigcup_{n=1}^{\infty} A_n$, then, $x \in A_n$ for some n . Let n be the least such natural. Now, since A_n is countable, there is a natural $m \geq 1$ such that x is the m th element of A_n . Define $f : \bigcup_{n=1}^{\infty} A_n \rightarrow \mathbb{N} \times \mathbb{N}$ by $f(x) = (m, n)$. Then, f is an injection. Since $\mathbb{N} \times \mathbb{N}$ is countable, we conclude that $\bigcup_{n=1}^{\infty} A_n$ is countable. To show that $\mathbb{Q} \times \mathbb{Q}$ is countable, let $\mathbb{Q} = (r_1, r_2, r_3, \dots)$, and, for each $n \geq 1$, set $A_n = \{r_n\} \times \mathbb{Q}$. Then, A_n is countable for each $n \geq 1$. Hence, so is $\mathbb{Q} \times \mathbb{Q}$, since it equals $\bigcup_{n=1}^{\infty} A_n$.

1.7.4. Suppose that $[0, 1]$ is countable, and list the elements as

$$\begin{aligned} a_1 &= .d_{11}d_{12}\dots \\ a_2 &= .d_{21}d_{22}\dots \\ a_3 &= .d_{31}d_{32}\dots \\ &\dots \end{aligned}$$

Let $a = .d_1d_2\dots$, where d_n is any digit chosen, such that $d_n \neq d_{nn}, n \geq 1$ (the “diagonal” in the above listing). Then, a is not in the list, so, the list is not complete, a contradiction. Hence, $[0, 1]$ is not countable.

1.7.5. Note that $i + j = n + 1$ implies $i^3 + j^3 \geq (n + 1)^3/8$ since at least one of i or j is $\geq (n + 1)/2$. Sum (7.6) in the order of \mathbb{N}^2 given in §1.7:

$$\begin{aligned} \sum_{(m,n) \in \mathbb{N}^2} \frac{1}{m^3 + n^3} &= \sum_{n=1}^{\infty} \left(\sum_{i+j=n+1} \frac{1}{i^3 + j^3} \right) \\ &\leq \sum_{n=1}^{\infty} \left[\sum_{i+j=n+1} \frac{8}{(n+1)^3} \right] \\ &= \sum_{n=1}^{\infty} \frac{8n}{(n+1)^3} \leq 8 \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty. \end{aligned}$$

1.7.6. Since $n^{-s} < 1$, the geometric series implies

$$\frac{1}{n^s - 1} = \frac{n^{-s}}{1 - n^{-s}} = \sum_{m=1}^{\infty} (n^{-s})^m.$$

Summing over $n \geq 2$,

$$\begin{aligned}\sum_{n=2}^{\infty} \frac{1}{n^s - 1} &= \sum_{n=2}^{\infty} \left(\sum_{m=1}^{\infty} n^{-sm} \right) \\ &= \sum_{m=1}^{\infty} \left(\sum_{n=2}^{\infty} n^{-ms} \right) \\ &= \sum_{m=1}^{\infty} Z(ms).\end{aligned}$$

1.7.7. Since $\sum |a_n|$ and $\sum |b_n|$ converge, from the text, we know that their Cauchy product converges. Thus, with $c_n = \sum_{i+j=n+1} a_i b_j$ and by the triangle inequality,

$$\sum_{n=1}^{\infty} |c_n| \leq \sum_{n=1}^{\infty} \left(\sum_{i+j=n+1} |a_i| |b_j| \right) < \infty.$$

Hence, $\sum_{n=1}^{\infty} c_n$ converges absolutely. Now, in the difference

$$D_N = \sum_{n=1}^N \left(\sum_{i+j=n+1} a_i b_j \right) - \left(\sum_{i=1}^N a_i \right) \left(\sum_{j=1}^N b_j \right)$$

there will be cancellation, with the absolute value of each of the remaining terms occurring as summands in

$$\sum_{n=N+1}^{\infty} \left(\sum_{i+j=n+1} |a_i| |b_j| \right).$$

Thus, $D_N \rightarrow 0$, as $N \nearrow \infty$. On the other hand, $D_N \rightarrow \sum_{n=1}^{\infty} c_n - (\sum a_i)(\sum b_j)$, hence, the result.

1.7.8. Let $\bar{a}_n = (-1)^{n+1} a_n$ and $\bar{b}_n = (-1)^{n+1} b_n$. If $\sum \bar{c}_n$ is the Cauchy product of $\sum \bar{a}_n$ and $\sum \bar{b}_n$, then,

$$\bar{c}_n = \sum_{i+j=n+1} \bar{a}_i \bar{b}_j = \sum_{i+j=n+1} (-1)^{i+1} (-1)^{j+1} a_i b_j = (-1)^{n+1} c_n,$$

where $\sum c_n$ is the Cauchy product of $\sum a_n$ and $\sum b_n$.

1.7.9. As in Exercise 13 of §1.5, $|x_n - x_m| \leq x_n x'_n x''_n \dots x_n^{(n-1)}$, for $m \geq n \geq 1$. Since $x_n^{(n-1)} = 1/q_n$, this yields $|x_n - x_m| \leq 1/q_n$, for $m \geq n \geq 1$. Hence, if $q_n \rightarrow \infty$, $(1/q_n)$ is an error sequence for (x_n) . Now, suppose that $q_n \not\rightarrow \infty$. Then, there a c with $q_n \leq c$ for infinitely many n . Hence, if N_n is the number of q_k 's, $k \leq n+2$, bounded by c , $\lim_{n \nearrow \infty} N_n = \infty$. Since $x_n^{(k)} \leq (q_{k+2} + 1)/(q_{k+2} + 2)$, the first inequality above implies that

$$|x_n - x_m| \leq \left(\frac{c+1}{c+2} \right)^{N_n}, \quad m \geq n \geq 1.$$

Set $a = (c + 1)/(c + 2)$. Then, in this case, (a^{N_n}) is an error sequence for (x_n) . For the golden mean x , note that $x = 1 + 1/x$, hence, $x = (1 \pm \sqrt{5})/2$. Since $x > 0$, we must take the $+$.

A.2 Chapter 2

Exercises 2.1

2.1.1. Let Q_n be the rectangle $(0, 1) \times (1/n, 1)$, $n \geq 1$. Then, (Q_n) covers $R = (0, 1) \times (0, 1)$, but no finite subcollection covers R .

2.1.2. Let $Q_n = (-\infty, \infty) \times (1/n, \infty)$, $Q'_n = (1/n, \infty) \times (-\infty, \infty)$, and $Q'' = [0, 0] \times [0, 0] = \{(0, 0)\}$. Arrange these rectangles into a single sequence $Q'', Q_1, Q'_1, Q_2, Q'_2, \dots$. This sequence covers $R = [0, 1] \times [0, 1]$, but no finite subcollection covers R .

2.1.3. For simplicity, we assume $a = 0$, $b = 1$. Let the limiting point be $L = .d_1d_2\dots$. By the construction of L , it is a limit point. Since x_* is the smallest limit point (Exercise 9 of §1.5), $x_* \leq L$. Now, note that if $t \in [0, 1]$ satisfies $t \leq x_n$ for all $n \geq 1$, then, $t \leq$ any limit point of (x_n) . Hence, $t \leq x_*$. Since changing finitely many terms of a sequence does not change x_* , we conclude that $x_* \geq t$ for all t satisfying $t \leq x_n$ for all but finitely n . Now, by construction, there are, at most, finitely many terms $x_n \leq .d_1$. Hence, $.d_1 \leq x_*$. Similarly, there are finitely many terms $x_n \leq .d_1d_2$. Hence, $.d_1d_2 \leq x_*$. Continuing in this manner, we conclude that $.d_1d_2\dots d_N \leq x_*$. Letting $N \nearrow \infty$, we obtain $L \leq x_*$. Hence, $x_* = L$.

Exercises 2.2

2.2.1. If $\lim_{x \rightarrow c} f(x) \neq 0$, there is at least one sequence $x_n \rightarrow c$ with $x_n \neq c$, $n \geq 1$, and $f(x_n) \not\rightarrow 0$. From Exercise 8 of §1.5, this means there is a subsequence (x_{k_n}) and an $N \geq 1$, such that $|f(x_{k_n})| \geq 1/N$, $n \geq 1$. But this means that, for all $n \geq 1$, the reals x_{k_n} are rationals with denominators bounded in absolute value by N . Hence, $N!x_{k_n}$ are integers converging to $N!c$. But this cannot happen unless $N!x_{k_n} = N!c$ from some point on, i.e., $x_{k_n} = c$ from some point on, contradicting $x_n \neq c$ for all $n \geq 1$. Hence, the result.

2.2.2. Let $L = \inf\{f(x) : a < x < b\}$. We have to show that $f(x_n) \rightarrow L$ whenever $x_n \rightarrow a+$. So, suppose that $x_n \rightarrow a+$, and assume, first, $x_n \searrow a$, i.e., (x_n) is decreasing. Then, $(f(x_n))$ is decreasing. Hence, $f(x_n)$ decreases to some limit M . Since $f(x_n) \geq L$ for all $n \geq 1$, $M \geq L$. If $d > 0$, then, there is an $x \in (a, b)$ with $f(x) < L + d$. Since $x_n \searrow a$, there is an $n \geq 1$ with $x_n < x$. Hence, $M \leq f(x_n) \leq f(x) < L + d$. Since $d > 0$ is arbitrary, we conclude that $M = L$ or $f(x_n) \searrow L$. In general, if $x_n \rightarrow a+$, $x_n^* \searrow a$. Hence, $f(x_n^*) \searrow L$. But $x_n \leq x_n^*$, hence, $L \leq f(x_n) \leq f(x_n^*)$. So, by the ordering property, $f(x_n) \rightarrow L$.

This establishes the result for the inf. For the sup, repeat the reasoning, or apply the first case to $g(x) = -f(-x)$.

2.2.3. Assume f is increasing. If $a < c < b$, then, apply the previous exercise to f on (c, b) , concluding that $f(c+)$ exists. Since $f(x) \geq f(c)$ for $x > c$, we obtain $f(c+) \geq f(c)$. Apply the previous exercise to f on (a, c) to conclude that $f(c-)$ exists and $f(c-) \leq f(c)$. Hence, $f(c)$ is between $f(c-)$ and $f(c+)$. Now, if $a < A < B < b$ and there are N points in $[A, B]$ where f jumps by at least $\delta > 0$, then, we must have $f(B) - f(A) \geq N\delta$. Hence, given δ and A, B , there are, at most, finitely many such points. Choosing $A = a + 1/n$ and $B = b - 1/n$ and taking the union of all these points over all the cases $n \geq 1$, we see that there are, at most, countably many points in (a, b) at which the jump is at least δ . Now, choosing δ equal $1, 1/2, 1/3, \dots$, and taking the union of all the cases, we conclude that there are, at most, countably many points $c \in (a, b)$ at which $f(c+) > f(c-)$. The decreasing case is obtained by multiplying by minus.

2.2.4. Choose any $c \in (a, b)$, and consider the partition with just two points $a = x_0 < x_1 < x_2 < x_3 = b$, where $x_1 = c$ and $x_2 = x$. Then, the variation corresponding to this partition is $|f(x) - f(c)|$. Hence, $|f(x) - f(c)| \leq I$ for $x > c$. Similarly, $|f(x) - f(c)| \leq I$ for $x < c$. Hence, $f(x)$ is bounded by $I + |f(c)|$.

2.2.5. If f is increasing, then, there are no absolute values in the variation (2.1) which, therefore, collapses to $f(x_n) - f(x_1)$. Since f is also bounded, this last quantity is bounded by some number I . Hence, f is of bounded variation. Now, note f and g of bounded variation implies $-f$ and $f + g$ are also of bounded variation, since the minus does not alter the absolute values, and the variation of the sum $f + g$, by the triangle inequality, is less or equal to the sum of the variation of f and the variation of g . This implies the second statement.

2.2.6. If $a < x < y < b$, every partition $a = x_0 < x_1 < x_2 < \dots < x_n = x < x_{n+1} = b$ with $x_n = x$ yields, by adding the point $\{y\}$, a partition $a = x_0 < x_1 < x_2 < \dots < x_n = x < x_{n+1} = y < x_{n+2} = b$ with $x_{n+1} = y$. Since $v(x)$ and $v(y)$ are the sup of the variations over all such partitions, respectively, this and (2.1) yield $v(x) + |f(y) - f(x)| \leq v(y)$. Hence, v is increasing and, throwing away the absolute value, $v(x) - f(x) \leq v(y) - f(y)$, i.e., $v - f$ is increasing. Thus, $f = v - (v - f)$ is the difference of two increasing functions. Finally, since $0 \leq v \leq I$, both v and $v - f$ are bounded.

2.2.7. Look at the partition $x_1 = 1, x'_1, x_2 = 1/2, x'_2, \dots, x_n = 1/n, x'_n$, where x'_i is an irrational between x_i and x_{i+1} , $1 \leq i \leq n$ (for this to make sense, take $x_{n+1} = 0$). Then, the variation corresponding to this partition is $2s_n - 1$, where s_n is the n th partial sum of the harmonic series. But $s_n \nearrow \infty$.

Exercises 2.3

2.3.1. Let f be a polynomial with odd degree n and highest order coefficient a_0 . Since $x^k/x^n \rightarrow 0$, as $x \rightarrow \pm\infty$, for $n > k$, it follows that $f(x)/x^n \rightarrow a_0$. Since $x^n \rightarrow \pm\infty$, as $x \rightarrow \pm\infty$, it follows that $f(\pm\infty) = \pm\infty$, at least when $a_0 > 0$. When $a_0 < 0$, the same reasoning leads to $f(\pm\infty) = \mp\infty$. Thus, there are reals a, b with $f(a) > 0$ and $f(b) < 0$. Hence, by the intermediate value property, there is a c with $f(c) = 0$.

2.3.2. By definition of μ_c , $|f(x) - f(c)| \leq \mu_c(\delta)$ when $|x - c| < \delta$. Since $|x - c| < 2|x - c|$ for $x \neq c$, choosing $\delta = 2|x - c|$ yields $|f(x) - f(c)| \leq \mu_c(2|x - c|)$, for $x \neq c$. First, assume $\mu_c(0+) = 0$, and suppose that $x_n \rightarrow c$ with $x_n \neq c$ for all $n \geq 1$. Then, $|x_n - c| \rightarrow 0$. Hence, $|f(x_n) - f(c)| \leq \mu_c(2|x_n - c|) \rightarrow 0$. Hence, f is continuous at c . Conversely, if $\mu_c(0+) \neq 0$, setting $\epsilon = \mu_c(0+)/2$, $\mu_c(1/n) \geq 2\epsilon$. By the definition of the sup in the definition of μ_c , this implies that, for each $n \geq 1$, there is an $x_n \in (a, b)$ with $|x_n - c| \leq 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$. Since the sequence (x_n) converges to c and $f(x_n) \not\rightarrow f(c)$, it follows that f is not continuous at c .

2.3.3. Let $A = f((a, b))$. To show that A is an interval, it is enough to show that $(\inf A, \sup A) \subset A$. By definition of \inf and \sup , there are sequences $m_n \rightarrow \inf A$ and $M_n \rightarrow \sup A$ with $m_n \in A$, $M_n \in A$. Hence, there are reals c_n, d_n with $f(c_n) = m_n$, $f(d_n) = M_n$. Since $f([c_n, d_n])$ is a compact interval, it follows that $[m_n, M_n] \subset A$ for all $n \geq 1$. Hence, $(\inf A, \sup A) \subset A$. For the second part, if $f((a, b))$ is not an open interval, then, there is a $c \in (a, b)$ with $f(c)$ a max or a min. But this cannot happen: Since f is strictly monotone, we can always find an x and y to the right and to the left of c such that $f(x)$ and $f(y)$ are larger and smaller than $f(c)$. Thus, $f((a, b))$ is an open interval.

2.3.4. Let $a = \sup A$. Since $a \geq x$ for $x \in A$ and f is increasing, $f(a) \geq f(x)$ for $x \in A$, or $f(a)$ is an upper bound for $f(A)$. Since $a = \sup A$, there is a sequence $(x_n) \subset A$ with $x_n \rightarrow a$. By continuity, $f(x_n) \rightarrow f(a)$. Now, let M be any upper bound for $f(A)$. Then, $M \geq f(x_n)$, $n \geq 1$. Hence, $M \geq f(a)$. Thus, $f(a)$ is the least upper bound for $f(A)$. Similarly, for \inf .

2.3.5. Let $y_n = f(x_n)$. From Exercise 4, $f(x_n^*) = \sup\{f(x_k) : k \geq n\} = y_n^*$, $n \geq 1$. Since $x_n^* \rightarrow x^*$ and f is continuous, $y_n^* = f(x_n^*) \rightarrow f(x^*)$. Thus, $y^* = f(x^*)$. Similarly, for lower stars.

2.3.6. Remember x^r is defined as $(x^m)^{1/n}$ when $r = m/n$. Since

$$[(x^{1/n})^m]^n = (x^{1/n})^{mn} = [(x^{1/n})^n]^m = x^m,$$

$x^r = (x^{1/n})^m$ also. With $r = m/n \in \mathbb{Q}$ and p in \mathbb{Z} , $(x^r)^p = [(x^{1/n})^m]^p = (x^{1/n})^{mp} = x^{mp/n} = x^{rp}$. Now, let $r = m/n$ and $s = p/q$ with m, n, p, q integers and $nq \neq 0$. Then, $[(x^r)^s]^{nq} = (x^r)^{snq} = x^{rsnq} = x^{mp}$. By uniqueness of roots, $(x^r)^s = (x^{mp})^{1/nq} = x^{rs}$. Similarly, $(x^r x^s)^{nq} = x^{rnq} x^{snq} = x^{rnq+snq} = x^{(r+s)nq} = (x^{r+s})^{nq}$. By uniqueness of roots, $x^r x^s = x^{r+s}$.

2.3.7. We are given $a^b = \sup\{a^r : 0 < r < b, r \in \mathbb{Q}\}$, and we need to show that $a^b = c$, where $c = \inf\{a^s : s > b, s \in \mathbb{Q}\}$. If r, s are rationals with $r < b < s$, then, $a^r < a^s$. Taking the sup over all $r < b$ yields $a^b \leq a^s$. Taking the inf over all $s > b$ implies $a^b \leq c$. On the other hand, choose $r < b < s$ rational with $s - r < 1/n$. Then, $c \leq a^s < a^r a^{1/n} \leq a^b a^{1/n}$. Taking the limit as $n \nearrow \infty$, we obtain $c \leq a^b$.

2.3.8. In this solution, r, s , and t denote rationals. Given b , let (r_n) be a sequence of rationals with $r_n \rightarrow b^-$. If $t < bc$, then, $t < r_n c$ for n large. Pick one such r_n and call it r . Then, $s = t/r < c$ and $t = rs$. Thus, $t < bc$ iff t is of the form rs with $r < b$ and $s < c$. By Exercise 6, $(a^b)^c = \sup\{(a^b)^s : 0 < s < c\} = \sup\{(\sup\{a^r : 0 < r < b\})^s : 0 < s < c\} = \sup\{a^{rs} : 0 < r < b, 0 < s < c\} = \sup\{a^t : 0 < t < bc\} = a^{bc}$.

2.3.9. Since $f(x + x') = f(x)f(x')$, by induction, we obtain $f(nx) = f(x)^n$. Hence, $f(n) = f(1)^n = a^n$ for n natural. Also $a = f(1) = f(1 + 0) = f(1)f(0) = af(0)$. Hence, $f(0) = 1$. Since $1 = f(n - n) = f(n)f(-n) = a^n f(-n)$, we obtain $f(-n) = a^{-n}$ for n natural. Hence, $f(n) = a^n$ for $n \in \mathbb{Z}$. Now, $(f(m/n))^n = f(n(m/n)) = f(m) = a^m$, so, by uniqueness of roots, $f(m/n) = a^{m/n}$. Hence, $f(r) = a^r$ for $r \in \mathbb{Q}$. If x is real, choose rationals $r_n \rightarrow x$. Then, $a^{r_n} = f(r_n) \rightarrow f(x)$. Since we know that a^x is continuous, $a^{r_n} \rightarrow a^x$. Hence, $f(x) = a^x$.

2.3.10. Given $\epsilon > 0$, we seek $\delta > 0$, such that $|x - 1| < \delta$ implies $|(1/x) - 1| < \epsilon$. By the triangle inequality, $|x| = |(x - 1) + 1| \geq 1 - |x - 1|$. So, $\delta < 1$ and $|x - 1| < \delta$ implies $|x| > 1 - \delta$ and

$$\left| \frac{1}{x} - 1 \right| = \frac{|x - 1|}{|x|} < \frac{\delta}{1 - \delta}.$$

Solving $\delta/(1 - \delta) = \epsilon$, we have found a δ , $\delta = \epsilon/(1 + \epsilon)$, satisfying $0 < \delta < 1$ and the ϵ - δ criterion.

2.3.11. Let A_n be the set of all real roots of all polynomials with degree d and rational coefficients a_0, a_1, \dots, a_d , with denominators bounded by n and satisfying

$$|a_0| + |a_1| + \dots + |a_d| + d \leq n.$$

Since each polynomial has finitely many roots and there are finitely many polynomials involved here, for each n , the set A_n is finite. But the set of algebraic numbers is $\bigcup_{n=1}^{\infty} A_n$, hence, is countable.

2.3.12. Let b be a root of f , $b \neq a$, and let $f(x) = (x - b)g(x)$. If b is rational, then, the coefficients of g are necessarily rational (this follows from the construction of g in the text) and $0 = f(a) = (a - b)g(a)$. Hence, $g(a) = 0$. But the degree of g is less than the degree of f . This contradiction shows that b is irrational.

2.3.13. Write $f(x) = (x - a)g(x)$. By the previous exercise, $f(m/n)$ is never zero. Since $n^d f(m/n)$ is an integer,

$$n^d |f(m/n)| = n^d |m/n - a| |g(m/n)| \geq 1,$$

or

$$(2.1) \quad \left| a - \frac{m}{n} \right| \geq \frac{1}{n^d |g(m/n)|}.$$

Since g is continuous at a , choose $\delta > 0$ such that $\mu_a(\delta) < 1$, i.e., such that $|x - a| < \delta$ implies $|g(x) - g(a)| \leq \mu_a(\delta) < 1$. Then, $|x - a| < \delta$ implies $|g(x)| < |g(a)| + 1$. Now, we have two cases: Either $|a - m/n| \geq \delta$ or $|a - m/n| < \delta$. In the first case, we obtain the required inequality with $c = \delta$. In the second case, we obtain $|g(m/n)| < |g(a)| + 1$. Inserting this in the denominator of the right side of (2.1) yields

$$\left| a - \frac{m}{n} \right| \geq \frac{1}{n^d (|g(a)| + 1)},$$

which is the required inequality with $c = 1/[|g(a)| + 1]$. Now, let

$$c = \min \left(\delta, \frac{1}{|g(a)| + 1} \right).$$

Then, in either case, the required inequality holds with this choice of c .

2.3.14. Let a be the displayed real. Then, a is irrational since its decimal expansion is not repeating. Let s_n denote the n th partial sum. Then, for $k \geq n$, $(k + 1)! \geq n!k$ and $10^{n!} \geq 2$, so,

$$\begin{aligned} |a - s_n| &= \sum_{k=n+1}^{\infty} \frac{1}{10^{(k+1)!}} \\ &\leq \sum_{k=n+1}^{\infty} \frac{1}{10^{n!k}} \\ &= \frac{1}{(10^{n!})^{n+1}} \sum_{k=0}^{\infty} \frac{1}{(10^{n!})^k} \\ &\leq \frac{1}{(10^{n!})^{n+1}} \sum_{k=0}^{\infty} \frac{1}{2^k} = \frac{1}{(10^{n!})^n} \cdot \frac{2}{10^{n!}}. \end{aligned}$$

Now, write $s_n = M/N$ with $M \in \mathbb{N}$ and $N = 10^{n!}$. Then, for all $\epsilon > 0$, choose $n \geq 1$ satisfying $2/10^{n!} < \epsilon$. This yields

$$\left| a - \frac{M}{N} \right| < \frac{\epsilon}{N^n}.$$

By Exercise 13, this shows that the algebraic order of a is more than n . Since $n \geq 1$ may be chosen arbitrarily large and $\epsilon > 0$ may be chosen arbitrarily small, a is transcendental.

2.3.15. From §1.6, we know that $\sum n^{-r}$ converges when $r = 1 + 1/N$. Given $s > 1$ real, we can always choose N with $1 + 1/N < s$. The result follows by comparison.

2.3.16. To show that $b^{\log_a c} = c^{\log_a b}$, apply \log_a obtaining $\log_a b \log_a c$ from either side. For the second part, $\sum 1/5^{\log_3 n} = \sum 1/n^{\log_3 5}$ which converges since $\log_3 5 > 1$.

2.3.17. Such an example cannot be continuous by the results of §2.3. Let $f(x) = x + 1/2$, $0 \leq x < 1/2$, and $f(x) = x - 1/2$, $1/2 \leq x < 1$, $f(1) = 1$. Then, f is a bijection, hence, invertible.

2.3.18. First, assume f is increasing. Then, by Exercise 3 of §2.2, $f(c-) = f(c) = f(c+)$ for all except, at most, countably many points $c \in (a, b)$, where there are, at worst, jumps. Hence, f is continuous for all but at most countably many points at which there are at worst jumps. If f is of bounded variation, then, $f = g - h$ with g and h bounded increasing. But, then, f is continuous wherever both g and h are continuous. Thus, the set of discontinuities of f is, at most, countable with the discontinuities at worst jumps.

2.3.19. Let $M = \sup\{f(x) : x \in \mathbb{R}\}$. Then, $M > -\infty$. If $M < \infty$, for each $n \geq 1$, choose x_n with $f(x_n) > M - 1/n$. Then, $f(x_n) \rightarrow M$. If $M = \infty$, for each $n \geq 1$, choose x_n with $f(x_n) > n$. Then, $f(x_n) \rightarrow M$. Now, from §2.1, (x_n) subconverges to some x where x may equal $\pm\infty$. If (x_n) subconverges to $\pm\infty$, then, $f(x_n)$ subconverges to $-\infty$ since $f(\infty) = f(-\infty) = -\infty$. But this cannot happen since $M > -\infty$. Hence, (x_n) must subconverge to some real x . Since f is continuous, $(f(x_n))$ must subconverge to $f(x)$. Hence, $f(x) = M$. This shows that M is finite and M is a max.

2.3.20. If $x_n \rightarrow \infty$, then, $x_n y - f(x_n) = x_n(y - f(x_n)/x_n) \rightarrow \infty(y - \infty) = -\infty$ by superlinearity. If $x_n \rightarrow -\infty$, by superlinearity, $x_n y - f(x_n) \rightarrow -\infty$. Hence, for each fixed y , the function $h(x) = xy - f(x)$ satisfies $h(\pm\infty) = -\infty$. Hence, $g(y)$ is well defined by the previous problem. Now, for $x > 0$ fixed and $y_n \rightarrow \infty$, $g(y_n) \geq xy_n - f(x)$. Hence,

$$\frac{g(y_n)}{y_n} \geq x - \frac{f(x)}{y_n}.$$

It follows that the lower limit of $(g(y_n)/y_n)$ is $\geq x$. Since x is arbitrary, it follows that the lower limit is ∞ . Hence, $g(y_n)/y_n \rightarrow \infty$. Since (y_n) was any sequence converging to ∞ , we conclude that $\lim_{y \rightarrow \infty} g(y)/|y| = \infty$. Similarly, $\lim_{y \rightarrow -\infty} g(y)/|y| = \infty$. Thus, g is superlinear.

2.3.21. Suppose that $y_n \rightarrow y$. We want to show that $g(y_n) \rightarrow g(y)$. Let $L^* \geq L_*$ be the upper and lower limits of $(g(y_n))$. For all z , $g(y_n) \geq zy_n - f(z)$. Hence, $L_* \geq zy - f(z)$. Since z is arbitrary, taking the sup over all z , we obtain $L_* \geq g(y)$. For the reverse inequality, let (y'_n) be a subsequence of (y_n) satisfying $g(y'_n) \rightarrow L^*$. Pick, for each $n \geq 1$, x'_n with $g(y'_n) = x'_n y'_n - f(x'_n)$. From §2.1, (x'_n) subconverges to some x , possibly infinite. If $x = \pm\infty$, then, superlinearity

(see previous solution) implies the subconvergence of $(g(y'_n))$ to $-\infty$. But $L^* \geq L_* \geq g(y) > -\infty$, so, this cannot happen. Thus, (x'_n) subconverges to a finite x . Hence, by continuity, $g(y'_n) = x'_n y'_n - f(x'_n)$ subconverges to $xy - f(x)$ which is $\leq g(y)$. Since by construction, $g(y'_n) \rightarrow L^*$, this shows that $L^* \leq g(y)$. Hence, $g(y) \leq L_* \leq L^* \leq g(y)$ or $g(y_n) \rightarrow g(y)$.

2.3.22. Note that $0 \leq f(x) \leq 1$ and $f(x) = 1$ iff $x \in \mathbb{Z}$. We are supposed to take the limit in m first, then, n . If $x \in \mathbb{Q}$, then, there is an $N \in \mathbb{N}$, such that $n!x \in \mathbb{Z}$ for $n \geq N$. Hence, $f(n!x) = 1$ for $n \geq N$. For such an x , $\lim_{m \rightarrow \infty} [f(n!x)]^m = 1$, for every $n \geq N$. Hence, the double limit is 1 for $x \in \mathbb{Q}$. If $x \notin \mathbb{Q}$, then, $n!x \notin \mathbb{Q}$, so, $f(n!x) < 1$, so, $[f(n!x)]^m \rightarrow 0$, as $m \nearrow \infty$, for every $n \geq 1$. Hence, the double limit is 0 for $x \notin \mathbb{Q}$.

2.3.23. In the definition of $\mu_c(\delta)$, we are to maximize $|(1/x) - (1/c)|$ over all $x \in (0, 1)$ satisfying $|x - c| < \delta$ or $c - \delta < x < c + \delta$. In the first case, if $\delta \geq c$, then, $c - \delta \leq 0$. Hence, all points x near and to the right of 0 satisfy $|x - c| < \delta$. Since $\lim_{x \rightarrow 0+} (1/x) = \infty$, in this case, $\mu_c(\delta) = \infty$. In the second case, if $0 < \delta < c$, then, x varies between $c - \delta > 0$ and $c + \delta$. Hence,

$$\left| \frac{1}{x} - \frac{1}{c} \right| = \frac{|x - c|}{xc}$$

is largest when the numerator is largest ($|x - c| = \delta$) and the denominator is smallest ($x = c - \delta$). Thus,

$$\mu_c(\delta) = \begin{cases} \delta/(c^2 - c\delta), & 0 < \delta < c, \\ \infty, & \delta \geq c. \end{cases}$$

Now, $\mu(\delta)$ equals the sup of $\mu_c(\delta)$ for all $c \in (0, 1)$. But, for δ fixed and $c \rightarrow 0+$, $\delta \geq c$ eventually. Hence, $\mu(\delta) = \infty$ for all $\delta > 0$. Hence, $\mu(0+) = \infty$, or f is not uniformly continuous on $(0, 1)$.

2.3.24. Follow the proof of the uniform continuity theorem. If $\mu(0+) > 0$, set $\epsilon = \mu(0+)/2$. Then, since μ is increasing, $\mu(1/n) \geq 2\epsilon$ for all $n \geq 1$. Hence, for each $n \geq 1$, by the definition of the sup in the definition of $\mu(1/n)$, there is a $c_n \in \mathbb{R}$ with $\mu_{c_n}(1/n) > \epsilon$. Now, by the definition of the sup in $\mu_{c_n}(1/n)$, for each $n \geq 1$, there is an $x_n \in \mathbb{R}$ with $|x_n - c_n| < 1/n$ and $|f(x_n) - f(c_n)| > \epsilon$. Then, by compactness (§2.1), (x_n) subconverges to some real x or to $x = \pm\infty$. It follows that (c_n) subconverges to the same x . Hence, $\epsilon < |f(x_n) - f(c_n)|$ subconverges to $|f(x) - f(x)| = 0$, a contradiction.

2.3.25. If $\sqrt{2}^{\sqrt{2}}$ is rational, we are done. Otherwise $a = \sqrt{2}^{\sqrt{2}}$ is irrational. In this case, let $b = \sqrt{2}$. Then, $a^b = (\sqrt{2}^{\sqrt{2}})^{\sqrt{2}} = \sqrt{2}^2 = 2$ is rational.

A.3 Chapter 3

Exercises 3.1

3.1.1. Since $a > 0$, $f(0) = 0$. If $a = 1$, we already know that f is not differentiable at 0. If $a < 1$, then, $g(x) = (f(x) - f(0))/(x - 0) = |x|^a/x$ satisfies $|g(x)| \rightarrow \infty$ as $x \rightarrow 0$, so, f is not differentiable at 0. If $a > 1$, then, $g(x) \rightarrow 0$ as $x \rightarrow 0$. Hence, $f'(0) = 0$.

3.1.2. Since $\sqrt{2}$ is irrational, $f(\sqrt{2}) = 0$. Hence, $(f(x) - f(\sqrt{2}))/(x - \sqrt{2}) = f(x)/(x - \sqrt{2})$. Now, if x is irrational, this expression vanishes whereas, if x is rational with denominator d ,

$$q(x) \equiv \frac{f(x) - f(\sqrt{2})}{x - \sqrt{2}} = \frac{1}{d^3(x - \sqrt{2})}.$$

By Exercise 10 of §1.4 it seems that the limit will be zero, as $x \rightarrow \sqrt{2}$. To prove this, suppose that the limit of $q(x)$ is not zero, as $x \rightarrow \sqrt{2}$. Then, there exists at least one sequence $x_n \rightarrow \sqrt{2}$ with $q(x_n) \not\rightarrow 0$. It follows that there is a $\delta > 0$ and a sequence $x_n \rightarrow \sqrt{2}$ with $|q(x_n)| \geq \delta$. But this implies all the reals x_n are rational. If d_n is the denominator of x_n , $n \geq 1$, we obtain, from Exercise 10 of §1.4,

$$|q(x_n)| \leq \frac{d_n^2}{d_n^3 c} = \frac{1}{d_n c}.$$

Since $d_n \rightarrow \infty$ (Exercise 1 of §2.2), we conclude that $q(x_n) \rightarrow 0$, contradicting our assumption. Hence, our assumption must be false, i.e., $\lim_{x \rightarrow \sqrt{2}} q(x) = 0$ or $f'(\sqrt{2}) = 0$.

3.1.3. Since f is superlinear (Exercise 20 of §2.3), $g(y)$ is finite, and the max is attained at some critical point x . Differentiating $xy - ax^2/2$ with respect to x yields $0 = y - ax$, or $x = y/a$ for the critical point, which, as previously said, must be the global max. Hence,

$$g(y) = (y/a)y - a(y/a)^2/2 = y^2/2a.$$

Since $f'(x) = ax$ and $g'(y) = y/a$, it is clear they are inverses.

3.1.4. Suppose that $g'(\mathbb{R})$ is bounded above. Then, $g'(x) \leq c$ for all x . Hence, $g(x) - g(0) = g'(z)(x - 0) \leq cx$ for $x > 0$, which implies $g(x)/x \leq c$ for $x > 0$, which contradicts superlinearity. Hence, $g'(\mathbb{R})$ is not bounded above. Similarly, $g'(\mathbb{R})$ is not bounded below.

3.1.5. To show that $f'(c)$ exists, let $x_n \rightarrow c$ with $x_n \neq c$ for all $n \geq 1$. Then, for each $n \geq 1$ there is a y_n strictly between c and x_n , such that $f(x_n) - f(c) = f'(y_n)(x_n - c)$. Since $x_n \rightarrow c$, $y_n \rightarrow c$, and $y_n \neq c$ for all $n \geq 1$. Since $\lim_{x \rightarrow c} f'(x) = L$, it follows that $f'(y_n) \rightarrow L$. Hence, $(f(x_n) - f(c))/(x_n - c) \rightarrow L$. Since (x_n) was arbitrary, we conclude that $f'(c) = L$.

3.1.6. To show that $f'(c) = f'(c+)$, let $x_n \rightarrow c+$. Then, for each $n \geq 1$, there is a y_n between c and x_n , such that $f(x_n) - f(c) = f'(y_n)(x_n - c)$. Since $x_n \rightarrow c+$, $y_n \rightarrow c+$. It follows that $f'(y_n) \rightarrow f'(c+)$. Hence, $(f(x_n) - f(c))/(x_n - c) \rightarrow f'(c+)$, i.e., $f'(c) = f'(c+)$. Similarly, for $f'(c-)$.

3.1.7. If $a = x_0 < x_1 < x_2 < \cdots < x_{n+1} = b$ is a partition, the mean value theorem says $f(x_k) - f(x_{k-1}) = f'(z_k)(x_k - x_{k-1})$ for some z_k between x_{k-1} and x_k , $1 \leq k \leq n+1$. Since $|f'(x)| \leq I$, we obtain $|f(x_k) - f(x_{k-1})| \leq I(x_k - x_{k-1})$. Summing over $1 \leq k \leq n+1$, we see that the variation corresponding to this partition is $\leq I(b - a)$. Since the partition was arbitrary, the result follows.

3.1.8. Let $c \in \mathbb{Q}$. We have to show that, for some $n \geq 1$, $f(c) \geq f(x)$ for all x in $(c - 1/n, c + 1/n)$. If this were not the case, for each $n \geq 1$, we can find a real x_n satisfying $|x_n - c| < 1/n$ and $f(x_n) > f(c)$. But, by Exercise 1 of §2.2, we know that $f(x_n) \rightarrow 0$ since $x_n \rightarrow c$ and $x_n \neq c$, contradicting $f(c) > 0$. Hence, c must be a local maximum.

3.1.9. If f is even, then, $f(-x) = f(x)$. Differentiating yields $-f'(-x) = f'(x)$, or f' is odd. Similarly, if f is odd. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is even,

$$\begin{aligned} g(-y) &= \max_{-\infty < x < \infty} (x(-y) - f(x)) \\ &= \max_{-\infty < -x < \infty} (x(-y) - f(x)) \\ &= \max_{-\infty < x < \infty} ((-x)(-y) - f(-x)) \\ &= \max_{-\infty < x < \infty} (xy - f(x)) = g(y). \end{aligned}$$

Hence, g is even.

Exercises 3.2

3.2.1. Let $f(x) = e^x$. Since $f(x) - f(0) = f'(c)x$ for some $0 < c < x$, we have $e^x - 1 = e^c x$. Since $c > 0$, $e^c > 1$. Hence, $e^x - 1 \geq x$ for $x \geq 0$.

3.2.2. Since g is even (Exercise 9 of §3.1), it is enough to compute $g(y)$ for $y \geq 0$. In this case, since $f(x) \geq 0$, the maximum is attained for $x \geq 0$,

$$g(y) = \max_{x \geq 0} (xy - f(x)),$$

so, we look only at $x \geq 0$. Also $(xy - f(x))' = 0$ iff $y = f'(x)$, i.e., $y = e^x$. Thus, $x > 0$ is a critical point if $y > 1$ and $x = \log y$, which gives $xy - f(x) = y \log y - y + 1$. If $0 \leq y \leq 1$, the function $x \mapsto xy - f(x)$ has no critical points in $(0, \infty)$. Hence, it is maximized at $x = 0$, i.e., $g(y) = 0$ when $0 \leq y \leq 1$. If $y > 1$, we obtain the critical point $x = \log y$, the corresponding critical value $y \log y - y + 1$, and the endpoint values 0 and $-\infty$. To see which of these three values is largest, note that $(y \log y - y + 1)' = \log y > 0$ for $y > 1$ and, thus, $y \log y - y + 1 \geq 0$ for $y \geq 1$. Hence, $g(y) = y \log y - y + 1$ for $y \geq 1$ and $g(y) = 0$ for $0 \leq y \leq 1$.

3.2.3. If $f(x) = \log(1+x)$, then, by l'Hopital's rule $\lim_{x \rightarrow 0} \log(1+x)/x = \lim_{x \rightarrow 0} 1/(1+x) = 1$. This proves the first limit. If $a \neq 0$, set $x_n = a/n$. Then, $x_n \rightarrow 0$ with $x_n \neq 0$ for all $n \geq 1$. Hence,

$$\lim_{n \nearrow \infty} n \log(1 + a/n) = a \lim_{n \nearrow \infty} \frac{\log(1 + x_n)}{x_n} = a.$$

By taking exponentials, we obtain $\lim_{n \nearrow \infty} (1 + a/n)^n = e^a$ when $a \neq 0$. If $a = 0$, this is immediate, so, the second limit is true for all a . Now, if $a_n \rightarrow a$, then, for some $N \geq 1$, $a - \epsilon \leq a_n \leq a + \epsilon$ for all $n \geq N$. Hence,

$$\left(1 + \frac{a - \epsilon}{n}\right)^n \leq \left(1 + \frac{a_n}{n}\right)^n \leq \left(1 + \frac{a + \epsilon}{n}\right)^n$$

for $n \geq N$. Thus, the upper and lower limits of the sequence in the middle lie between $e^{a-\epsilon}$ and $e^{a+\epsilon}$. Since $\epsilon > 0$ is arbitrary, the upper and lower limits must both equal e^a . Hence, we obtain the third limit.

3.2.4. Let $b = x/(n+1)$, $v = (n+1)/n$, and $e_n = (1 + x/n)^n$. Then, $|b| < 1$, so, by (2.2), $(1+b)^{nv} \geq (1+vb)^n$. Hence, $e_{n+1} \geq e_n$.

3.2.5. Since $p > 1$, f is superlinear, so, the max exists. First, note that g is even by Exercise 9 of §3.1. Therefore, we need compute $g(y)$ only for $y \geq 0$. If $y = 0$, we obtain $g(0) = 0$, whereas, if $y > 0$, we need consider only $x > 0$, since $f(x) \geq 0$, in computing the sup for $g(y)$. To find the critical points, solve $0 = (xy - f(x))' = y - f'(x)$ for $x > 0$ obtaining $x^{p-1} = y$ or $x = y^{1/(p-1)}$. Plugging this into $xy - f(x)$ yields the required g . Finally, f' and g' are odd, and, for $x \geq 0$, $f'(x) = x^{p-1}$ and $g'(y) = y^{q-1}$ are inverses since $(p-1)(q-1) = 1$.

3.2.6. If $f(x) = (1+x^2)^{-1/2}$, then, $f(x) - f(0) = f'(c)(x-0)$ with $0 < c < x$. Since $f'(c) = -c(1+c^2)^{-3/2} > -c > -x$ for $x > 0$,

$$\frac{1}{\sqrt{1+x^2}} - 1 = f'(c)x \geq -x^2,$$

which implies the result.

3.2.7. With $f(t) = -t^{-x}$ and $f'(t) = xt^{-x-1}$, the left side of the displayed inequality equals $f(2j) - f(2j-1)$, which equals $f'(c)$ with $2j-1 < c < 2j$. Hence, the left side is $xc^{-x-1} \leq x(2j-1)^{-x-1}$.

3.2.8. First, suppose that $L = 0$. If there is no such x , then, f' is never zero. Hence, $f' > 0$ on (a, b) or $f' < 0$ on (a, b) , contradicting $f'(c) < 0 < f'(d)$. Hence, there is an x satisfying $f'(x) = 0$. In general, let $g(x) = f(x) - Lx$. Then, $f'(c) < L < f'(d)$ implies $g'(c) < 0 < g'(d)$, so, the general case follows from the case $L = 0$.

3.2.9. From Exercise 4 of §3.1, $g'(\mathbb{R})$ is not bounded above nor below. But g' satisfies the intermediate value property. Hence, the range of g' is an interval. Hence, $g'(\mathbb{R}) = \mathbb{R}$.

Exercises 3.3

3.3.1. Since $f'(x) = (1/2) - (1/x^2)$, the only positive critical point (Figure A.1) is $x = \sqrt{2}$. Moreover, $f(\infty) = f(0+) = \infty$, so, $\sqrt{2}$ is a global minimum over $(0, \infty)$, and $f(\sqrt{2}) = \sqrt{2}$. Also, $f''(x) = 2/x^3 > 0$, so, f is convex.




FIGURE A.1. The graph of $(x + 2/x)/2$.

3.3.2. If $a < b < c$ and $t = (b - a)/(c - a)$, then, $b = (1 - t)a + tc$. Hence, by convexity,

$$f(b) \leq (1 - t)f(a) + tf(c).$$

Subtracting $f(a)$ from both sides and, then, dividing by $b - a$ yields $s[a, b] \leq s[a, c]$. Instead, if we subtract both sides from $f(c)$ and, then, divide by $c - b$, we obtain $s[a, c] \leq s[b, c]$.

3.3.3. Applying the previous to $x < c < t$ yields $s[x, c] \leq s[c, t]$. Since $s[c, t] \rightarrow f'(c)$ as $t \rightarrow c+$, we obtain $s[x, c] \leq f'(c)$. Similarly, for $t < c < y$, $s[t, c] \leq s[c, y]$. Since $s[t, c] \rightarrow f'(c)$, as $t \rightarrow c-$, we obtain $f'(c) \leq s[c, y]$. Hence, $s[x, c] \leq f'(c) \leq s[c, y]$. Now, clear denominators in $s[x, c] \leq f'(c)$ to obtain $f(x) \geq f(c) + f'(c)(x - c)$ for $x < c$. Repeating this with $f'(c) \leq s[c, x]$, $x > c$, we obtain the same inequality for $x > c$.

3.3.4. Exercise 2 says $x \mapsto s[c, x]$ is an increasing function of x . Hence,

$$f'_+(c) \equiv \lim_{x \rightarrow c+} s[c, x] = \inf\{s[c, x] : x > c\}$$

exists. Similarly $f'_-(c) = \lim_{y \rightarrow c-} s[y, c]$ exists. Since, for $y < c < x$, $s[y, c] \leq s[c, x]$, we conclude that $f'_-(c) \leq f'_+(c)$.

3.3.5. If c is a maximum of f , then, for $a < y < c < x < b$, $s[y, c] \geq 0$ and $s[c, x] \leq 0$. But, from convexity, $s[y, c] \leq s[c, x]$. Hence, $s[c, x] = s[y, c] = 0$ for $a < y < c < x < b$, which implies $f(x) = f(c) = f(y)$ or f is constant.

3.3.6. Let $s^f[c, x]$, $s^g[c, x]$, and $s^{f-g}[c, x]$ denote the difference quotients corresponding to f , g , and $f - g$, respectively. Since c is a local maximum of $f - g$, we have $s^{f-g}[c, x] \leq 0$ for $x > c$. Hence,

$$\begin{aligned} f'_+(c) &= \lim_{x \rightarrow c+} s^f[c, x] \\ &= \lim_{x \rightarrow c+} s^{f-g}[c, x] + s^g[c, x] \end{aligned}$$

$$\leq \lim_{x \rightarrow c+} s^g[c, x] = g'(c).$$

Similarly, $g'(c) \leq f'_-(c)$. But $f'_-(a) \leq f'_+(a)$. Hence, $f'(c)$ exists and equals $g'(c)$.

3.3.7. g is convex because it is the sup of lines. To see this, fix $a < b$ and $0 \leq t \leq 1$. Then,

$$x[(1-t)a+tb]-f(x) = (1-t)[xa-f(x)]+t[xb-f(x)] \leq (1-t)g(a)+tg(b).$$

Since this is true for every x , taking the sup of the left side over x yields the convexity of g . Now, fix y , and suppose that $g(y) = xy - f(x)$, i.e., x attains the max in the definition of $g(y)$. Then, for any z , $g(z) \geq xz - f(x)$. Hence, for $z > y$,

$$\frac{g(z) - g(y)}{z - y} \geq \frac{xz - f(x) - xy + f(x)}{z - y} = x.$$

Letting $z \rightarrow y+$ yields $g'_+(y) \geq x$. Considering $z < y$ instead, we obtain $g'_-(y) \leq x$.

3.3.8. Fix y , suppose that $g(y) = xy - f(x)$ and $g(y) = x'y - f(x')$, and let $x'' = (1-t)x + tx'$ with $0 < t < 1$. Then, by strict convexity,

$$x''y - f(x'') > (1-t)(xy - f(x)) + t(x'y - f(x')) = (1-t)g(y) + tg(y) = g(y),$$

contradicting the definition of $g(y)$. Thus, for each y , the max is attained at a unique x , call it $x^\#(y)$. To show that $g'(y) = x^\#(y)$, we show that $g'_\pm(y) = x^\#(y)$, since g is differentiable at y iff $g'_+(y) = g'_-(y)$. From the previous exercise, we know $g'_+(y) \geq x$. To obtain the reverse inequality, let $y_n \rightarrow y+$ and let $x_n = x^\#(y_n)$, i.e., $g(y_n) = x_n y_n - f(x_n)$. Then, as in Exercise 19 of §2.3, (x_n) must be bounded. If z is any limit point of (x_n) , then, since f is continuous, $x_n y_n - f(x_n)$ subconverges to $zy - f(z)$. Since g is continuous, $g(y_n)$ converges to $g(y)$. Hence, $g(y) = xz - f(z)$ or $z = x^\#(y)$. This shows that (x_n) converges to $x^\#(y)$ (Exercise 10 of §1.5). Now, since $g(y) \geq x_n y - f(x_n)$,

$$\frac{g(y_n) - g(y)}{y_n - y} \leq \frac{x_n y_n - f(x_n) - x_n y + f(x_n)}{y_n - y} = x_n \rightarrow x^\#(y),$$

hence, $g'_+(y) \leq x^\#(y)$. This establishes $g'_+(y) = x^\#(y)$. Similarly, $g'_-(y) = x^\#(y)$. Hence, $g'(y) = x^\#(y)$.

3.3.9. The only part we have not derived in the previous exercises is the formula for $f(x)$. To see this, note, by the formula for $g(y)$, that $f(x) + g(y) \geq xy$ for all x, y , which implies $f(x) \geq \max_y [xy - g(y)]$. To obtain equality, the previous exercise shows that, for each y , $f(x) + g(y) = xy$ iff $x = g'(y)$. But g' is surjective by Exercise 9 of §3.2. Hence, for each x , we can find a y satisfying $x = g'(y)$. For this y , $f(x) + g(y) = xy$. Hence, $f(x) = \max_y (xy - g(y))$.

3.3.10. If the max is attained at x , x is a critical point. Hence, $(xy - f(x))' = 0$ or $y = f'(x)$. Since y is arbitrary, f' is surjective.

3.3.11. From above, we know that g is superlinear, differentiable and convex. Moreover, $f(x) + g(y) = xy$ iff $x = g'(y)$ iff $y = f'(x)$. Hence, f' and g' are inverses. Since f is strictly convex, f' is strictly increasing. Hence, its inverse g' is strictly increasing. Hence, g is strictly convex.

3.3.12. Here, f' does not exist at 0. However, the previous Exercise says that g' is trying to be the inverse of f' which suggests that $f'(0)$ should be defined to be (Figure A.2) the line segment $[-1, 1]$ on the vertical axis. Of course, with such a definition, f' is no longer a function, but something more general (a set-valued mapping).




FIGURE A.2. The graphs of f, g, f', g' (3.3.12).

3.3.13. Since $(e^x)'' > 0$, e^x is convex. Hence, $e^{t \log a + (1-t) \log b} \leq t e^{\log a} + (1-t) e^{\log b}$. But this simplifies to $a^t b^{1-t} \leq ta + (1-t)b$.

3.3.14. If f and g are as in Exercise 11, then, $g'(f'(x)) = x$. Differentiating yields $g''[f'(x)]f''(x) = 1$, so, $f''(x) \neq 0$. Since convexity implies $f''(x) \geq 0$, we obtain $f''(x) > 0$ for all x . Conversely, if $f''(x) > 0$, by the inverse function theorem g is twice differentiable with $g''(x) = (g')'(x) = 1/(f')'[g'(x)] = 1/f''[g'(x)]$. Hence,

$$g''(x) = \frac{1}{f''[g'(x)]}.$$

Since g is twice differentiable, the right side of this last equation is differentiable. Hence, g'' is differentiable or g is thrice differentiable. If g is n times differentiable, then, the smoothness of f and the right side imply g is $(n+1)$ times differentiable. Hence, g is smooth.

Exercises 3.4

3.4.1. By Taylor's Theorem, $f(c+t) = f(c) + f'(c)t + f''(\eta)t^2/2$ with η between c and $c+t$. Since $f(c+t) \geq 0$ and $f''(\eta) \leq 1/2$, we obtain

$$0 \leq f(c) + f'(c)t + t^2/4, \quad -\infty < t < \infty.$$

Hence, the quadratic $Q(t) = f(c) + f'(c)t + t^2/4$ has at most one solution. But this implies (Exercise 5 of §1.4) $f'(c)^2 - 4(1/4)f(c) \leq 0$, which gives the result.

3.4.2. Since $n! \geq 2^{n-1}$ (Exercise 3 of §1.6),

$$\sum_{k \geq n+1} \frac{1}{k!} \leq \sum_{k \geq n+1} 2^{1-k} = 2^{1-n}.$$

Now choose $n = 15$. Then, $2^{n-1} > 10^4$, so, adding the terms of the series up to $n = 15$ yields accuracy to four decimals. Adding these terms yields $e \sim 2.718281829$ where \sim means that the error is $< 10^{-4}$.

3.4.3. For $n = 1$, (1) is true since we can choose $R_1(x) = 1$. Also (2) is true for $n = 1$ since $h(0) = 0$. Now, assume that (1) and (2) are true for n . Then,

$$\lim_{x \rightarrow 0+} \frac{h^{(n)}(x) - h^{(n)}(0)}{x} = \lim_{x \rightarrow 0+} \frac{R_n(x)}{x} e^{-1/x} = \lim_{t \rightarrow \infty} t R_n(1/t) e^{-t} = 0$$

since $t^d e^{-t} \rightarrow 0$, as $t \rightarrow \infty$, and R_n is rational. Since $\lim_{x \rightarrow 0+} [h^{(n)}(x) - h^{(n)}(0)]/x = 0$, this establishes (2) for $n+1$. Now, establish (1) for $n+1$ using the product rule and the fact that the derivative of a rational function is rational. Thus, (1) and (2) hold by induction for all $n \geq 1$.

3.4.4. If $n \geq 100$, then,

$$n! \geq 101 \cdot 102 \cdot \dots \cdot n \geq 100 \cdot 100 \cdot \dots \cdot 100 = 100^{n-100}.$$

. Hence, $(n!)^{1/n} \geq 100^{(n-100)/n}$, which clearly approaches 100. Thus, the lower limit of $((n!)^{1/n})$ is ≥ 100 . Since 100 may be replaced by any N , the result follows.

3.4.5. Apply the binomial theorem with $v = -1/2$ to obtain

$$\frac{1}{\sqrt{1+x}} = 1 - \frac{1}{2}x + \frac{1}{2} \cdot \frac{3}{4}x^2 - \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6}x^3 + \dots$$

Now, replace x by $-x^2$.

3.4.6. If $f(x) = \log(1+x)$, then, $f(0) = 0$, $f'(x) = 1/(1+x)$, $f''(x) = -1/(1+x)^2$, and $f^{(n)}(x) = (-1)^{n-1}(n-1)!/(1+x)^n$ for $n \geq 1$. Hence, $f^{(n)}(0)/n! = (-1)^{n-1}/n$ or

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

3.4.7. Inserting $x = 0$ in the series yields $f(0) = a_0$. Now,

$$f'(x) = a_1 + 2a_2x + 3a_3x^2 + \dots,$$

so, inserting $x = 0$ yields $f'(0) = a_1$. Differentiating the series repeatedly yields

$$f^{(n)}(x) = n!a_n + (n+1)!a_{n+1}\frac{x}{1!} + (n+2)!a_{n+2}\frac{x^2}{2!} + \dots$$

Hence, $f^{(n)}(0) = n!a_n$ or $f^{(n)}(0)/n! = a_n$. Thus, the series is the Taylor series centered at zero.

3.4.8. If $1/(1+x) = \sum_{n=0}^{\infty} a_n x^n$ and $\log(1+x) = \sum_{n=1}^{\infty} b_n x^n$, then, $\log(1+x)/(1+x) = \sum_{n=1}^{\infty} c_n x^n$, where

$$c_n = a_0 b_n + a_1 b_{n-1} + \dots + a_{n-1} b_1 + a_n b_0.$$

Since $a_n = (-1)^n$, $b_n = (-1)^{n-1}/n$, we obtain

$$\begin{aligned} c_n &= \sum_{i=0}^n a_{n-i} b_i = \sum_{i=0}^n (-1)^{n-i} (-1)^{i-1}/i \\ &= -(-1)^n \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right). \end{aligned}$$

3.4.9. First, $f(x) = f(0) + f'(0)x + h(x)x^2/2$ with h continuous and $h(0) = f''(0) = q$. So,

$$f\left(\frac{x}{\sqrt{n}}\right) = 1 + \frac{h(x/\sqrt{n})x^2}{2n}.$$

Now, apply Exercise 3 of §3.2 with $a_n = h(x/\sqrt{n})x^2/2 \rightarrow qx^2/2$ to obtain the result.

3.4.10. Since $\sinh(\pm\infty) = \pm\infty$, $\sinh(\mathbb{R}) = \mathbb{R}$. Since $\sinh' = \cosh > 0$, \sinh is bijective, hence, invertible. Note that $\cosh^2 - \sinh^2 = 1$, so,

$$\cosh^2(\operatorname{arcsinh} x) = 1 + x^2.$$

The derivative of $\operatorname{arcsinh} : \mathbb{R} \rightarrow \mathbb{R}$ is (by the IFT)

$$\operatorname{arcsinh}'(x) = \frac{1}{\sinh'(\operatorname{arcsinh} x)} = \frac{1}{\cosh(\operatorname{arcsinh} x)} = \frac{1}{\sqrt{1+x^2}}.$$

Since $1/\sqrt{1+x^2}$ is smooth, so is $\operatorname{arcsinh}$. Now, \cosh is superlinear since $\cosh x \geq e^{|x|}/2$ and strictly convex since $\cosh'' = \cosh > 0$. Hence, the max in

$$g(y) = \max_{-\infty < x < \infty} (xy - \cosh(x))$$

is attained at $x = \operatorname{arcsinh} y$. We obtain $g(y) = y \operatorname{arcsinh} y - \sqrt{1+y^2}$.

3.4.11. With $a_n = (-1)^n/4^n(n!)^2$, use the ratio test,

$$\frac{|a_n|}{|a_{n+1}|} = 4(n+1)^2 \rightarrow \infty.$$

Hence, the radius ρ equals ∞ .

3.4.12. Here, neither the ratio test nor the root test work. If $|x| \geq 1$, by the n th term test, the series diverges, whereas, if $|x| < 1$, the series converges absolutely by comparison with the geometric series. Hence, $R = 1$.

3.4.13. For $n = 0$, this is immediate. If $n \geq 1$,

$$\begin{aligned} \binom{-1/2}{n} &= \frac{\left(-\frac{1}{2}\right)\left(-\frac{1}{2}-1\right)\cdots\left(-\frac{1}{2}-n+1\right)}{1 \cdot 2 \cdot \dots \cdot n} \\ &= \frac{(-1)^n 1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2^n n!} \\ &= \frac{(-1)^n 1 \cdot 2 \cdot 3 \cdot 4 \cdot \dots \cdot (2n-1) \cdot (2n)}{2^n n! \cdot 2 \cdot 4 \cdot 6 \cdot \dots \cdot (2n)} \\ &= \frac{(-1)^n (2n)!}{2^n n! \cdot 2^n \cdot 1 \cdot 2 \cdot 3 \cdot \dots \cdot n} \\ &= \frac{(-1)^n (2n)!}{4^n (n!)^2}. \end{aligned}$$

3.4.14. Inserting $-x$ for x in the series yields $f(-x) = a_0 - a_1x + a_2x^2 - \dots$. Hence,

$$f^e(x) = \frac{f(x) + f(-x)}{2} = a_0 + a_2x^2 + \dots$$

But f is even iff $f = f^e$, so, the result follows. The odd case is similar.

3.4.15. Define $h(t)$ by $e^t = 1 + t + t^2h(t)/2$, $t \neq 0$. Then, $e^t - 1 = t(1 + th(t)/2)$, and, by the exponential series, $\lim_{t \rightarrow 0} h(t) = 1$. Now,

$$\begin{aligned} \frac{1}{e^t - 1} - \frac{1}{t} &= \frac{1}{t[1 + th(t)/2]} - \frac{1}{t} \\ &= \frac{1}{t} \left[\frac{1}{1 + th(t)/2} - 1 \right] \\ &= \frac{1}{t} \cdot \frac{th(t)/2}{1 + th(t)/2} \\ &= \frac{-h(t)/2}{1 + th(t)/2}. \end{aligned}$$

This shows that the limit is $-1/2$.

Exercises 3.5

3.5.1. We have to show that $(xx' + yy')^2 \leq (x^2 + y^2)(x'^2 + y'^2)$. By multiplying, show that

$$(xx' + yy')^2 + (xy' - x'y)^2 = (x^2 + y^2)(x'^2 + y'^2).$$

This implies Cauchy-Schwarz.

3.5.2. Since $\sin^2 + \cos^2 = 1$, $\tan^2 + 1 = 1/\cos^2$, or $\cos^2 = 1/(1 + \tan^2)$. Hence, $\cos^2(\theta/2) = 1/(1 + t^2)$, which gives

$$\cos \theta = 2 \cos(\theta/2)^2 - 1 = \frac{2}{1 + t^2} - 1 = \frac{1 - t^2}{1 + t^2}.$$

Also,

$$\sin \theta = 2 \sin(\theta/2) \cos(\theta/2) = 2t \cos^2(\theta/2) = \frac{2t}{1 + t^2}.$$

Also,

$$\tan \theta = \sin \theta / \cos \theta = \frac{2t}{1 - t^2}.$$

3.5.3. f is differentiable at all nonzero reals, hence, continuous there. Since $|f(x)| \leq |x|$, f is also continuous at $x = 0$. Compute the variation of f corresponding to the partition $x_k = 2/(k\pi)$, $k = 1, \dots, n$. Since $f(x_k) = 0$ for k even and $f(x_k) = \pm 2/k\pi$ for k odd, the variation is larger than $(2/\pi)(1/2 + 1/3 + \dots + 1/n)$. Hence, f is not of bounded variation near 0.

3.5.4. If $x \neq 0$, then, $f'(x) = 2x \sin(1/x) - \cos(1/x)$. If $x = 0$, then, $f'(0) = \lim_{x \rightarrow 0} f(x)/x = \lim_{x \rightarrow 0} x \sin(1/x) = 0$. Hence, $|f'(x)| \leq 1 + 2|x|$ for all x . By Exercise 7 of §3.1, f is of bounded variation on any bounded interval.

3.5.5. If $(x', y') = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$ and $(x'', y'') = (x' \cos \phi - y' \sin \phi, x' \sin \phi + y' \cos \phi)$, then, by the addition formulas,

$$\begin{aligned} x'' &= (x \cos \theta - y \sin \theta) \cos \phi - (x \sin \theta + y \cos \theta) \sin \phi \\ &= x(\cos \theta \cos \phi - \sin \theta \sin \phi) - y(\sin \theta \cos \phi + \cos \theta \sin \phi) \\ &= x \cos(\theta + \phi) - y \sin(\theta + \phi). \end{aligned}$$

Similarly,

$$y'' = x \sin(\theta + \phi) + y \cos(\theta + \phi).$$

Thus, $R_\theta \circ R_\phi = R_{\theta+\phi}$.

3.5.6. Draw the line L through C parallel to the line through A and B , and mark two points A' and B' on L , one on either side of C . Then, by rotation and translation invariance, $\angle CAB = \angle ACA'$, and $\angle ABC = \angle B'CB$. Hence,

$$\angle ABC + \angle BCA + \angle CAB = \angle B'CB + \angle BCA + \angle ACA' = B'CA' = \pi$$

since B' , C , and A' all lie on L .

3.5.7. It is enough to show that $x^n/n! \geq x^{n+1}/(n+1)!$ for $0 \leq x \leq 3$ and $n \geq 3$. But, simplifying, we see that the inequality holds iff $x \leq n+1$, which is true, since $x \leq 3 \leq n+1$.

3.5.8. If $(z, w) = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$ and $(z', w') = (x' \cos \theta - y' \sin \theta, x' \sin \theta + y' \cos \theta)$, then,

$$zz' + ww' = (x \cos \theta - y \sin \theta)(x' \cos \theta - y' \sin \theta)$$

$$\begin{aligned}
& + (x \sin \theta + y \cos \theta)(x' \sin \theta + y' \cos \theta) \\
& = (xx' + yy')(\cos^2 \theta + \sin^2 \theta) = xx' + yy'.
\end{aligned}$$

3.5.9. Since both sides are translation invariant, we may assume that $B = O$, $A = (a, a')$, and $C = (c, c')$. Then,

$$\begin{aligned}
|AC|^2 &= (a - c)^2 + (a' - c')^2 = (a^2 + a'^2) + (c^2 + c'^2) - 2(ac + a'c') \\
&= |AO|^2 + |OC|^2 - 2|AO||OC|\cos(AOC)
\end{aligned}$$

by (5.5).

3.5.10. Let $\theta = \pi/9$. Then, $\sin(8\theta) = \sin(\theta)$ since $8\theta + \theta = 9\theta = \pi$. Hence,

$$\begin{aligned}
\sin \theta &= \sin(8\theta) \\
&= 2 \sin(4\theta) \cos(4\theta) \\
&= 4 \sin(2\theta) \cos(2\theta) \cos(4\theta) \\
&= 8 \sin(\theta) \cos(\theta) \cos(2\theta) \cos(4\theta).
\end{aligned}$$

Now, divide both sides by $8 \sin \theta$.

3.5.11. $\sin(\pi/3) = \sin(\pi - \pi/3)$ by (5.3). Hence,

$$\sin(\pi/3) = \sin(2\pi/3) = 2 \sin(\pi/3) \cos(\pi/3),$$

or $\cos(\pi/3) = 1/2$, which implies $\sin(\pi/3) = \sqrt{3}/2$ and $\tan(\pi/3) = \sqrt{3}$. From (5.2), we obtain $\sin(\pi/2 - x) = \cos x$ and $\cos(\pi/2 - x) = \sin x$. Hence, $\sin(\pi/6) = 1/2$, $\cos(\pi/6) = \sqrt{3}/2$, and $\tan(\pi/6) = 1/\sqrt{3}$. Also, $0 = \cos(\pi/2) = 2 \cos(\pi/4)^2 - 1$, so, $\cos(\pi/4) = 1/\sqrt{2}$, $\sin(\pi/4) = 1/\sqrt{2}$, and $\tan(\pi/4) = 1$.

3.5.12. Let s_n denote the sum on the left. Since $2 \cos a \sin b = \sin(a + b) - \sin(a - b)$ from (5.3), with $a = x$ and $b = x/2$,

$$\begin{aligned}
\sin(x/2)s_1 &= \sin(x/2) + 2 \cos x \sin(x/2) \\
&= \sin(x/2) + \sin(3x/2) - \sin(x/2) = \sin(3x/2).
\end{aligned}$$

Thus, the result is true when $n = 1$. Assuming the result is true for n and repeating the same reasoning with $a = (n + 1)x$ and $b = x/2$,

$$\begin{aligned}
\sin(x/2)s_{n+1} &= \sin(x/2)(s_n + 2 \cos((n + 1)x)) \\
&= \sin((n + 1/2)x) + 2 \sin(x/2) \cos((n + 1)x) \\
&= \sin((n + 1/2)x) + \sin((n + 3/2)x) - \sin((n + 1/2)x) \\
&= \sin((n + 3/2)x).
\end{aligned}$$

This derives the result for $n + 1$. Hence, the result is true for all $n \geq 1$.

3.5.13. Divide $2 \cos(2x) = 2 \cos^2 x - 2 \sin^2 x$ by $\sin(2x) = 2 \sin x \cos x$.

3.5.14. The first identity is established using the double-angle formula. When $n = 2$, the identity (n) says $(x^4 - 1) = (x^2 - 1)(x^2 + 1)$ and is true. Now, assume

the validity of the identity (n). To obtain the identity (2n), replace x by x^2 in (n), and use the first identity. Then,

$$\begin{aligned}
 \frac{x^{4n} - 1}{x^4 - 1} &= \prod_{k=1}^{n-1} [x^4 - 2x^2 \cos(k\pi/n) + 1] \\
 &= \prod_{k=1}^{n-1} [x^2 - 2x \cos(k\pi/2n) + 1] \cdot \prod_{k=1}^{n-1} [x^2 - 2x \cos(\pi - k\pi/2n) + 1] \\
 &= \prod_{k=1}^{n-1} [x^2 - 2x \cos(k\pi/2n) + 1] \cdot \prod_{k=1}^{n-1} [x^2 - 2x \cos((2n - k)\pi/2n) + 1] \\
 &= \prod_{k=1}^{n-1} [x^2 - 2x \cos(k\pi/2n) + 1] \cdot \prod_{k=n+1}^{2n-1} [x^2 - 2x \cos(k\pi/2n) + 1] \\
 &= \prod_{\substack{k \neq n \\ 1 \leq k \leq 2n-1}} [x^2 - 2x \cos(k\pi/2n) + 1] \\
 &= \frac{1}{(x^2 + 1)} \cdot \prod_{k=1}^{2n-1} [x^2 - 2x \cos(k\pi/2n) + 1].
 \end{aligned}$$

Multiplying by $(x^4 - 1) = (x^2 - 1)(x^2 + 1)$, we obtain the result.

3.5.15. Let $a_n = 1/n$ and $c_n = \cos(nx)$, $n \geq 1$. Then, for $x \notin 2\pi\mathbb{Z}$, by Exercise 12, the sequence $b_n = c_1 + \cdots + c_n$, $n \geq 1$, is bounded. Hence, by the Dirichlet test, $\sum \cos(nx)/n$ converges.

Exercises 3.6

3.6.1. With $dv = e^x dx$ and $u = \cos x$, $v = e^x$ and $du = -\sin x dx$. So,

$$I = \int e^x \cos x dx = e^x \cos x + \int e^x \sin x dx.$$

Repeat with $dv = e^x dx$ and $u = \sin x$. We get $v = e^x$ and $du = \cos x dx$. Hence,

$$\int e^x \sin x dx = e^x \sin x - \int e^x \cos x dx.$$

Now, insert the second equation into the first to yield

$$I = e^x \cos x + [e^x \sin x - I].$$

Solving for I yields

$$\int e^x \cos x dx = \frac{1}{2} (e^x \cos x + e^x \sin x).$$

3.6.2. Let $u = \arcsin x$. Then, $x = \sin u$, so, $dx = \cos u du$. So,

$$\int e^{\arcsin x} dx = \int e^u \cos u du = \frac{1}{2} e^u (\sin u + \cos u)$$

by Exercise 1. Since $\cos u = \sqrt{1-x^2}$, we obtain

$$\int e^{\arcsin x} dx = \frac{1}{2} e^{\arcsin x} (x + \sqrt{1-x^2}).$$

3.6.3.

$$\begin{aligned} \int \frac{x+1}{\sqrt{1-x^2}} dx &= \int \frac{x}{\sqrt{1-x^2}} dx + \int \frac{dx}{\sqrt{1-x^2}} \\ &= -\frac{1}{2} \int \frac{d(1-x^2)}{\sqrt{1-x^2}} + \arcsin x \\ &= -(1-x^2)^{1/2} + \arcsin x = \arcsin x - \sqrt{1-x^2}. \end{aligned}$$

3.6.4. If $u = \arctan x$, then,

$$\int \frac{\arctan x}{1+x^2} dx = \int u du = \frac{1}{2} u^2 = \frac{1}{2} (\arctan x)^2.$$

3.6.5. If $u = (\log x)^2$ and $dv = x^2 dx$, then, $v = x^3/3$ and $du = 2(\log x) dx/x$. Hence,

$$\int x^2 (\log x)^2 dx = \frac{1}{3} x^3 (\log x)^2 - \frac{2}{3} \int x^2 \log x dx.$$

If $u = \log x$ and $dv = x^2 dx$, then, $v = x^3/3$ and $du = dx/x$. Hence,

$$\int x^2 \log x dx = \frac{1}{3} x^3 \log x - \frac{1}{3} \int x^2 dx = \frac{1}{3} x^3 \log x - \frac{1}{9} x^3.$$

Now, insert the second integral into the first equation, and rearrange to obtain

$$\int x^2 (\log x)^2 dx = \frac{x^3}{27} (9 \log^2 x - 6 \log x + 2).$$

3.6.6. Take $u = \sqrt{1-e^{-2x}}$. Then, $u^2 = 1 - e^{-2x}$, so, $2u du = 2e^{-2x} dx = 2(1-u^2) dx$. Hence,

$$\int \sqrt{1-e^{-2x}} dx = \int \frac{u^2 du}{1-u^2} = \int \frac{du}{1-u^2} - u = \frac{1}{2} \log \left(\frac{1+u}{1-u} \right) - u$$

which simplifies to

$$\int \sqrt{1-e^{-2x}} dx = \log \left(1 + \sqrt{1-e^{-2x}} \right) + x - \sqrt{1-e^{-2x}}.$$

3.6.7. Since $|\sin| \leq 1$, $F'(0) = \lim_{x \rightarrow 0} F(x)/x = \lim_{x \rightarrow 0} x \sin(1/x) = 0$. Moreover,

$$F'(x) = 2x \sin(1/x) - \cos(1/x), \quad x \neq 0.$$

So, F' is not continuous at zero.

3.6.8. If f is of bounded variation, then, its discontinuities are, at worst, jumps (Exercise 18 of §2.3). But, if $f = F'$, then, f is a derivative and Exercise 6 of §3.1 says f cannot have any jumps. Thus, f must be continuous.

3.6.9. Let $\theta = \arcsin \sqrt{x}$. Then, $x = \sin^2 \theta$ and the left side of (6.5) equals 2θ . Now, $2x - 1 = 2\sin^2 \theta - 1 = -\cos(2\theta)$, so, the right side equals

$$\pi/2 - \arcsin(\cos(2\theta)) = \arccos[\cos(2\theta)] = 2\theta.$$

3.6.10. Since $(\arcsin x)' = 1/\sqrt{1-x^2}$ and the derivative of the exhibited series is the Taylor series (Exercise 5 of §3.4) of $1/\sqrt{1-x^2}$, the result follows from the theorem in this section.

3.6.11. Integration by parts: With $u = f(-x)$ and $dv = e^x dx$, $v = e^x$, and $du = -f'(-x)dx$. So,

$$\int e^x f(-x) dx = e^x f(-x) + \int e^x f'(-x) dx.$$

Repeating this procedure with derivatives of f replacing f , we obtain the result.

3.6.12. Divide the first equation in (5.2) by the second equation in (5.2). You obtain the tangent formula. Now, set $a = \arctan(1/5)$ and $b = \arctan(1/239)$. Then, $\tan a = 1/5$, so, $\tan(2a) = 5/12$, so, $\tan(4a) = 120/119$. Also, $\tan b = 1/239$, so,

$$\begin{aligned} \tan(4a - b) &= \frac{\tan(4a) - \tan b}{1 + \tan(4a) \tan b} = \frac{(120/119) - (1/239)}{1 + (120/119)(1/239)} \\ &= \frac{120 \cdot 239 - 119}{119 \cdot 239 + 120} = 1. \end{aligned}$$

Hence, $4a - b = \pi/4$.

3.6.13. Since

$$\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

is alternating with decreasing terms (as long as $0 < x < 1$), plugging in $x = 1/5$ and adding the first two terms yields $\arctan(1/5) = .1973$ with an error less than the third term which is less than 1×10^{-4} . Now, plugging $x = 1/239$ into the first term yields .00418 with an error less than the second term which is less than 10^{-6} . Since 16 times the first error plus 4 times the second error is less than 10^{-2} , $\pi = 16 \arctan(1/5) - 4 \arctan(1/239) = 3.14$ with an error less than 10^{-2} .

3.6.14. If $\theta = \arcsin(\sin 100)$, then, $\sin(\theta) = \sin 100$ and $|\theta| \leq \pi/2$. But $32\pi = 32(3.14) = 100.48$, and $31.5\pi = 98.9$, with an error less than $32 \times 10^{-2} = .32$. Hence, we are sure that $31.5\pi < 100 < 32\pi$ or $-\pi/2 < 100 - 32\pi < 0$, i.e., $\theta = 100 - 32\pi$.

3.6.15. Let $u = 1 - x^2$. Then, $du = -2x dx$. Hence,

$$\int \frac{-4x}{1-x^2} dx = \int \frac{2du}{u} = 2 \log u = 2 \log(1-x^2).$$

3.6.16. Completing the square, $x^2 - \sqrt{2}x + 1 = (x - 1/\sqrt{2})^2 + 1/2$. So, with $u = \sqrt{2}x - 1$ and $v = (\sqrt{2}x - 1)^2 + 1 = 2x^2 - 2\sqrt{2}x + 2 = 2(x^2 - \sqrt{2}x + 1)$,

$$\begin{aligned} \int \frac{4\sqrt{2} - 4x}{x^2 - \sqrt{2}x + 1} dx &= \int \frac{8\sqrt{2} - 8x}{(\sqrt{2}x - 1)^2 + 1} dx \\ &= \int \frac{4\sqrt{2}}{(\sqrt{2}x - 1)^2 + 1} dx - 2 \int \frac{2\sqrt{2}(\sqrt{2}x - 1)}{(\sqrt{2}x - 1)^2 + 1} dx \\ &= \int \frac{4du}{u^2 + 1} - 2 \int \frac{dv}{v} \\ &= 4 \arctan u - 2 \log v + 2 \log 2 \\ &= 4 \arctan(\sqrt{2}x - 1) - 2 \log(x^2 - \sqrt{2}x + 1). \end{aligned}$$

3.6.17. Let $s_n(x)$ denote the n th partial sum in (6.3). Then, if $0 < x < 1$, by the Leibnitz test,

$$s_{2n}(x) \leq \log(1+x) \leq s_{2n-1}(x), \quad n \geq 1.$$

In this last inequality, the number of terms in the partial sums is finite. Letting $x \nearrow 1$, we obtain

$$s_{2n}(1) \leq \log 2 \leq s_{2n-1}(1), \quad n \geq 1.$$

Now, let $n \nearrow \infty$.

A.4 Chapter 4

Exercises 4.1

4.1.1. The first subrectangle thrown out has area $(1/3) \times 1 = 1/3$, the next two each have area $1/9$, the next four each have area $1/27$, and so on. So, the areas of the removed rectangles sum to $(1/3)(1 + (2/3) + (2/3)^2 + \dots) = 1$. Hence, the area of what is left, C' , is zero. At the n th stage, the width of each of the remaining rectangles in C'_n is 3^{-n} . Since $C' \subset C'_n$, no rectangle in C' can have width greater than 3^{-n} . Since $n \geq 1$ is arbitrary, no open rectangle can lie in C' .

4.1.2. Here, the widths of the removed rectangles sum to $\alpha/3 + 2\alpha/3^2 + 4\alpha/3^3 + \dots = \alpha$, so, the area of what is left, C^α , is $1 - \alpha > 0$. At the n th stage, the width of each of the remaining rectangles in C_n^α is $3^{-n}\alpha$. Since $C^\alpha \subset C_n^\alpha$, no rectangle in C^α can have width greater than $3^{-n}\alpha$. Since $n \geq 1$ is arbitrary, no open rectangle can lie in C^α .

4.1.3. Here, all expansions are ternary. Since $[0, 2] \times [0, 2] = 2C_0$, given $(x, y) \in C_0$, we have to find $(x', y') \in C$ and $(x'', y'') \in C$ satisfying $x' + x'' = 2x$ and $y' + y'' = 2y$. Let $x = .d_1d_2d_3\dots$ and $y = .e_1e_2e_3\dots$. Then, for all $n \geq 1$, $2d_n$ and $2e_n$ are 0, 2, or 4. Thus, there are digits $d'_n, d''_n, e'_n, e''_n, n \geq 1$, equalling 0 or 2 and satisfying $d'_n + d''_n = 2d_n$ and $e'_n + e''_n = 2e_n$. Now, set $x' = .d'_1d'_2d'_3\dots$, $y' = .e'_1e'_2e'_3\dots$, $x'' = .d''_1d''_2d''_3\dots$, $y'' = .e''_1e''_2e''_3\dots$.

Exercises 4.2

4.2.1. If Q is a rectangle, then, so is $-Q$ and $\|Q\| = \|-Q\|$. If (Q_n) is a paving of A , then, $(-Q_n)$ is a paving of $-A$, so,

$$\text{area}(-A) \leq \sum_{n=1}^{\infty} \|-Q_n\| = \sum_{n=1}^{\infty} \|Q_n\|.$$

Since $\text{area}(A)$ is the inf of the sums on the right, we obtain $\text{area}(-A) \leq \text{area}(A)$. Applying this to $-A$, instead of A , yields $\text{area}(A) \leq \text{area}(-A)$ which yields reflection invariance, when combined with the previous inequality. For monotonicity, if $A \subset B$ and (Q_n) is a paving of B , then, (Q_n) is a paving of A . So, $\text{area}(A) \leq \sum_{n=1}^{\infty} \|Q_n\|$. Since the inf of the sums on the right over all pavings of B equals $\text{area}(B)$, $\text{area}(A) \leq \text{area}(B)$.

4.2.2. Let L be a line segment. If L is vertical, we already know that $\text{area}(L) = 0$. Otherwise, by translation and dilation invariance, we may assume that $L = \{(x, y) : 0 \leq x \leq 1, y = mx\}$. If $Q_i = ((i-1)/n, i/n) \times (m(i-1)/n, mi/n)$, $i = 1, \dots, n$, then, (Q_1, \dots, Q_n) covers L , except for the corner points. If $c > 1$, then, (cQ_1, \dots, cQ_n) is a paving of L , and $\sum_{i=1}^n \|cQ_i\| = c^2 m/n$. Since $n \geq 1$ is arbitrary, we conclude that $\text{area}(L) = 0$. Since any line L is a countable union of line segments, by subadditivity, $\text{area}(L) = 0$. Since $Q \subset \overline{Q}$, by monotonicity, $\text{area}(Q) \leq \text{area}(\overline{Q})$. Conversely, $\overline{Q} = Q \cup L_1 \cup L_2 \cup L_3 \cup L_4$ where L_i are bounded line segments, so, by subadditivity, $\text{area}(\overline{Q}) \leq \text{area}(Q) + \sum_i \text{area}(L_i) = \text{area}(Q)$. Hence, $\text{area}(Q) = \text{area}(\overline{Q})$. Given a compact rectangle $Q = [a, b] \times [c, d]$, let $Q' = (a, b) \times (c, d)$. Then, $\text{area}(Q) = \text{area}(Q') \leq \|Q'\| = \|Q\|$.

4.2.3. First, write $P = A \cup B \cup C$, where A and B are triangles with horizontal bases and C is a rectangle, all intersecting only along their edges. Since the sum of the naive areas of A , B , and C is the naive area of P , subadditivity yields

$$\begin{aligned} \text{area}(P) &\leq \text{area}(A) + \text{area}(B) + \text{area}(C) \\ &= \|A\| + \|B\| + \|C\| = \|P\|. \end{aligned}$$

To obtain the reverse inequality, draw two triangles B and C with horizontal bases, such that $P \cup B \cup C$ is a rectangle and P , B , and C intersect only along their edges. Then, the sum of the naive areas of P , B , and C equals the naive area of $P \cup B \cup C$, so, by subadditivity of area,

$$\begin{aligned} \|P\| + \|B\| + \|C\| &= \|P \cup B \cup C\| \\ &= \text{area}(P \cup B \cup C) \\ &\leq \text{area}(P) + \text{area}(B) + \text{area}(C) \\ &\leq \text{area}(P) + \|B\| + \|C\|. \end{aligned}$$

Cancelling $\|B\|$ and $\|C\|$, we obtain the reverse inequality $\|P\| \leq \text{area}(P)$.

4.2.4. If T is the trapezoid, T can be broken up into the union of a rectangle and two triangles. As before, by subadditivity, this yields $\text{area}(T) \leq \|T\|$. Also

two triangles can be added to T to obtain a rectangle. As before, this yields $\|T\| \leq \text{area}(T)$.

4.2.5. By extending the sides of the rectangles, $A \cup B$ can be decomposed into the union of finitely many rectangles intersecting only along their edges (seven rectangles in the general case). Moreover, since $A \cap B$ is one of these rectangles and is counted twice, using subadditivity, we obtain $\text{area}(A \cup B) \leq \text{area}(A) + \text{area}(B) - \text{area}(A \cap B)$. To obtain the reverse inequality, add two rectangles C and D , to fill in the corners, obtaining a rectangle $A \cup B \cup C \cup D$, and proceed as before.

4.2.6. If Q is a rectangle, then, $H(Q)$ is a rectangle and $\|H(Q)\| = |k| \cdot \|Q\|$. If (Q_n) is a paving of A , then, $(H(Q_n))$ is a paving of $H(A)$. So,

$$\text{area}[H(A)] \leq \sum_{n=1}^{\infty} \|H(Q_n)\| = \sum_{n=1}^{\infty} |k| \cdot \|Q_n\| = |k| \sum_{n=1}^{\infty} \|Q_n\|.$$

Taking the inf over all pavings of A yields $\text{area}[H(A)] \leq |k| \cdot \text{area}(A)$. In this last inequality replace A by $H^{-1}(A)$ and k by $1/k$ to obtain $|k| \cdot \text{area}(A) \leq \text{area}(H(A))$. Thus, $\text{area}[H(A)] = |k| \cdot \text{area}(A)$. V is similar.

4.2.7. Suppose that $(X, Y) = (ax + by, cx + dy)$ and (x, y) lies on a line with rise m and run n , $(x, y) = (nt + p, mt + q)$. Then, $(X, Y) = (Nt + P, Mt + Q)$, where $M = cn + dm$ and $N = an + bm$, or (X, Y) lies on a line with rise M and run N . Thus, L sends lines to lines. Since the slope of the new line depends only on the slope of the old line, parallel lines are sent to parallel lines. Hence, parallelograms are sent to parallelograms. Now, L is a bijection iff the pair of equations $(X, Y) = (ax + by, cx + dy)$ can be solved uniquely for (x, y) . But these equations are equivalent to $(dX - bY, -cX + dY) = (ad - bc)(x, y)$ which can be solved iff $\det L = ad - bc \neq 0$. Moreover, the solution is $K(X, Y) = (dX - bY, -cX + aY)/\det L = (AX + BY, CX + DY)$. So, $A = d/\det L$, $B = -b/\det L$, $C = -c/\det L$, and $D = a/\det L$, which shows that K is linear and $\det K = AD - BC = 1/\det L$.

4.2.8. If $Q = [0, 1] \times [0, 1]$, then, $L(Q)$ is a parallelogram with corners $(0, 0)$, (a, c) , (b, d) , and $(a + b, c + d)$. Since we know that the formulas for the areas of rectangles and parallelograms, we obtain (2.9) in this case (draw a picture). By dilation invariance, then, we obtain (2.9) when Q is any rectangle with the bottom left corner at the origin. By translation invariance, (2.9), then, is true for any rectangle Q . Now, use pavings as in Exercise 6 to obtain half of (2.9). Use invertibility to obtain the other half.

4.2.9. If L is not invertible, then, $ad - bc = 0$ or $ad = bc$. With the notation of solution 4.2.7, this implies $cX = aY$ and $dX = bY$. Thus, if not all of a, b, c, d are zero, at least one of the sets $A = \{(X, Y) : cX = aY\}$, $A' = \{(X, Y) : dX = bY\}$ is a line passing through the origin and $L(\mathbb{R}^2) \subset A \cap A'$. This shows that $L(\mathbb{R}^2)$ is contained in a line, when not all of a, b, c, d are zero. When $a = b = c = d = 0$, we have $(X, Y) = (0, 0)$. Hence, $L(\mathbb{R}^2) = \{(0, 0)\}$ is, again, contained in a line.

Thus, in either case, $L(\mathbb{R}^2)$ is contained in a line. Since a line has zero area and $\det(L) = 0$, in this case, both sides of (2.9) are equal.

4.2.10. If (a, b) is in the unit disk, then, $a^2 + b^2 < 1$. Hence, $|a| < 1$. Hence,

$$\sqrt{(\sqrt{2} - a)^2 + (0 - b)^2} \geq |\sqrt{2} - a| \geq \sqrt{2} - |a| \geq \sqrt{2} - 1.$$

Hence, $d(D, \{(\sqrt{2}, 0)\}) > 0$. For the second part, let (a_n) denote a sequence of rationals converging to $\sqrt{2}$. Then, $(a_n, 0) \in \mathbb{Q} \times \mathbb{Q}$, and

$$d((a_n, 0), (\sqrt{2}, 0)) \rightarrow 0, \quad n \nearrow \infty.$$

4.2.11. For $1 > h > 0$, let $D_h^+ = \{(x, y) \in D^+ : y > h\}$. Then, $D^+ \setminus D_h^+$ is contained in a rectangle with area h , and D_h^+ and $D^- = \{(x, y) \in D : y < 0\}$ are well separated. By reflection invariance, $\text{area}(D^+) = \text{area}(D^-)$. So, by subadditivity,

$$\begin{aligned} \text{area}(D) &\geq \text{area}(D_h^+ \cup D^-) = \text{area}(D_h^+) + \text{area}(D^-) \\ &\geq \text{area}(D^+) - \text{area}(D^+ \setminus D_h^+) + \text{area}(D^+) \\ &\geq 2 \cdot \text{area}(D^+) - h. \end{aligned}$$

Since $h > 0$ is arbitrary, we obtain $\text{area}(D) \geq 2 \cdot \text{area}(D^+)$. The reverse inequality follows by subadditivity.

4.2.12. The derivation is similar to the derivation of $\text{area}(C) = 0$ presented at the end of §4.2.

4.2.13. If T is the triangle joining $(0, 0)$, (a, b) , $(a, -b)$, where

$$(a, b) = (\cos(\pi/n), \sin(\pi/n)),$$

then, $\text{area}(T) = \sin(\pi/n) \cos(\pi/n)$. Since D_n is the union of n triangles T_1, \dots, T_n , each having the same area as T (rotation invariance), subadditivity yields $\text{area}(D_n) \leq n \sin(\pi/n) \cos(\pi/n) = n \sin(2\pi/n)/2$. The reverse inequality is obtained by shrinking each triangle T_i towards its center and using well separated additivity.

4.2.14. Let α denote the inf on the right side. Since (T_n) is a cover, we obtain $\text{area}(A) \leq \sum_{n=1}^{\infty} \text{area}(T_n) = \sum_{n=1}^{\infty} \|T_n\|$. Hence, $\text{area}(A) \leq \alpha$. On the other hand, if (Q_n) is any paving of A , write each $Q_n = T_n \cup T'_n$ as the union of two triangles to obtain a triangular paving $(T_n) \cup (T'_n)$. Hence,

$$\alpha \leq \sum_{n=1}^{\infty} \|T_n\| + \|T'_n\| = \sum_{n=1}^{\infty} \|Q_n\|.$$

Taking the inf over all pavings (Q_n) , we obtain $\alpha \leq \text{area}(A)$.

4.2.15. Assume not. Then, for every rectangle Q , $\text{area}(Q \cap A) \leq \alpha \cdot \text{area}(Q)$. If (Q_n) is any paving of A , then, $(Q_n \cap A)$ is a cover of A . So

$$\text{area}(A) \leq \sum_{n=1}^{\infty} \text{area}(Q_n \cap A) \leq \alpha \sum_{n=1}^{\infty} \text{area}(Q_n).$$

Taking the inf of the right side over all (Q_n) , $\text{area}(A) \leq \alpha \cdot \text{area}(A)$. Since $\alpha < 1$, this yields $\text{area}(A) = 0$.

Exercises 4.3

4.3.1. Apply dilation invariance to $g(x) = f(x)/x$. Then,

$$\begin{aligned} \int_0^{\infty} f(kx)x^{-1} dx &= k \int_0^{\infty} f(kx)(kx)^{-1} dx = k \int_0^{\infty} g(kx) dx \\ &= k \cdot \frac{1}{k} \int_0^{\infty} g(x) dx = \int_0^{\infty} f(x)x^{-1} dx. \end{aligned}$$

4.3.2. Let G be the subgraph of f over (a, b) , and let $H(x, y) = (-x, y)$. Then, $H(G)$ equals $\{(-x, y) : a < x < b, 0 < y < f(x)\}$. But this equals $\{(x, y) : -b < x < -a, 0 < y < f(-x)\}$, which is the subgraph of $f(-x)$ over $(-b, -a)$. Thus, by Exercise 6 of §4.2,

$$\int_a^b f(x) dx = \text{area}(G) = \text{area}(H(G)) = \int_{-b}^{-a} f(-x) dx.$$

4.3.3. Since f is uniformly continuous on $[a, b]$, there is a $\delta > 0$, such that $\mu(\delta) < \epsilon/(b-a)$. Here, μ is the uniform modulus of continuity of f over $[a, b]$. In §2.3, we showed that, with this choice of δ , for any partition $a = x_0 < x_1 < \dots < x_{n+1} = b$ of mesh $< \delta$ and choice of intermediate points $x_1^{\#}, \dots, x_{n+1}^{\#}$, the piecewise constant function $g(x) = f(x_i^{\#})$, $x_{i-1} < x < x_i$, $1 \leq i \leq n+1$, satisfies $|f(x) - g(x)| < \epsilon/(b-a)$ over (a, b) . Now, by additivity,

$$\int_a^b [g(x) \pm \epsilon/(b-a)] dx = \sum_{i=1}^{n+1} f(x_i^{\#})(x_i - x_{i-1}) \pm \epsilon,$$

and

$$g(x) - \epsilon/(b-a) < f(x) < g(x) + \epsilon/(b-a), \quad a < x < b.$$

Hence, by monotonicity,

$$\sum_{i=1}^{n+1} f(x_i^{\#})(x_i - x_{i-1}) - \epsilon \leq \int_a^b f(x) dx \leq \sum_{i=1}^{n+1} f(x_i^{\#})(x_i - x_{i-1}) + \epsilon,$$

which is to say that

$$\left| I - \sum_{i=1}^{n+1} f(x_i^{\#})(x_i - x_{i-1}) \right| \leq \epsilon.$$

4.3.4. Since $f(x) \leq x = g(x)$ on $(0, 1)$ and the subgraph of g is a triangle, by monotonicity, $\int_0^1 f(x) dx \leq \int_0^1 x dx = 1/2$. On the other hand, the subgraph of g equals the union of the subgraphs of f and $g - f$. So, by subadditivity,

$$1/2 = \int_0^1 g(x) dx \leq \int_0^1 f(x) dx + \int_0^1 [g(x) - f(x)] dx.$$

But $g(x) - f(x) > 0$ iff $x \in \mathbb{Q}$. So, the subgraph of $g - f$ is a countable (§1.7) union of line segments. By subadditivity, again, the area $\int_0^1 [g(x) - f(x)] dx$ of the subgraph of $g - f$ equals zero. Hence, $\int_0^1 f(x) dx = 1/2$.

4.3.5. Suppose that g is a constant $c \geq 0$, and let G be the subgraph of f . Then, the subgraph of $f + c$ is the union of the rectangle $Q = (a, b) \times (0, c]$ and the vertical translate $G + (0, c)$. Thus, by subadditivity and translation invariance,

$$\begin{aligned} \int_a^b [f(x) + g(x)] dx &= \text{area}[Q \cup (G + (0, c))] \\ &\leq \text{area}(G) + \text{area}(Q) \\ &= \int_a^b f(x) dx + \int_a^b g(x) dx. \end{aligned}$$

Let αQ denote the centered dilate of Q , $0 < \alpha < 1$. Then, αQ and $G + (0, c)$ are well separated. So,

$$\begin{aligned} \int_a^b [f(x) + g(x)] dx &\geq \text{area}[\alpha Q \cup (G + (0, c))] \\ &= \text{area}(G) + \alpha^2 \text{area}(Q) \\ &= \int_a^b f(x) dx + \alpha^2 \int_a^b g(x) dx. \end{aligned}$$

Let $\alpha \rightarrow 1$ to get the reverse inequality. Thus, the result is true when g is constant. If g is piecewise constant over a partition $a = x_0 < x_1 < \cdots < x_{n+1} = b$, then, apply the constant case to the intervals (x_{i-1}, x_i) , $i = 1, \dots, n+1$, and sum.

4.3.6. By subadditivity, $\int_0^\infty f(x) dx \leq \sum_{n=1}^\infty \int_{n-1}^n f(x) dx = \sum_{n=1}^\infty c_n$. For the reverse inequality,

$$\int_0^\infty f(x) dx \geq \int_0^N f(x) dx = \sum_{n=1}^N \int_{n-1}^n f(x) dx = \sum_{n=1}^N c_n.$$

Now, let $N \nearrow \infty$. This establishes the nonnegative case. Now, apply the nonnegative case to $|f|$. Then, f is integrable iff $\sum |c_n| < \infty$. Now, apply the nonnegative case to f^+ and f^- , and subtract to get the result for the integrable case.

4.3.7. To be Riemann integrable, the Riemann sum must be close to a specific real I for any partition with small enough mesh and any choice of intermediate points $x_1^\#, \dots, x_{n+1}^\#$. But, for any partition $a = x_0 < x_1 < \cdots < x_{n+1} =$

b , with any mesh size, we can choose intermediate points that are irrational, leading to a Riemann sum of 1. We can also choose intermediate points that are rational, leading to a Riemann sum of 0. Since no real I can be close to 0 and 1, simultaneously, f is not Riemann integrable.

4.3.8. Apply the integral test to $f(x) = g(x\delta)$ to get

$$\int_1^\infty g(x\delta) dx \leq \sum_{n=1}^\infty g(n\delta) \leq \int_1^\infty g(x\delta) dx + g(\delta).$$

By dilation invariance,

$$\int_\delta^\infty g(x) dx \leq \delta \sum_{n=1}^\infty g(n\delta) \leq \int_\delta^\infty g(x) dx + \delta g(\delta).$$

Since g is bounded, $\delta g(\delta) \rightarrow 0$, as $\delta \rightarrow 0+$. Now, let $\delta \rightarrow 0+$, and use continuity at the endpoints.

4.3.9. If f is even and nonnegative or integrable, by Exercise 2,

$$\begin{aligned} \int_{-b}^b f(x) dx &= \int_{-b}^0 f(x) dx + \int_0^b f(x) dx \\ &= \int_0^b f(-x) dx + \int_0^b f(x) dx \\ &= 2 \int_0^b f(x) dx. \end{aligned}$$

If f is odd and integrable,

$$\begin{aligned} \int_{-b}^b f(x) dx &= \int_{-b}^0 f(x) dx + \int_0^b f(x) dx \\ &= \int_0^b f(-x) dx + \int_0^b f(x) dx = 0. \end{aligned}$$

4.3.10. By the previous Exercise, $\int_{-\infty}^\infty e^{-a|x|} dx = 2 \int_0^\infty e^{-ax} dx$. By the integral test, $\int_1^\infty e^{-ax} dx \leq \sum_{n=1}^\infty e^{-an} = 1/(1 - e^{-a}) < \infty$. Hence,

$$\int_0^\infty e^{-ax} dx = \int_0^1 e^{-ax} dx + \int_1^\infty e^{-ax} dx \leq 1 + \frac{1}{1 - e^{-a}}.$$

4.3.11. Since f is superlinear, for any $M > 0$, there is a $b > 0$, such that $f(x)/x > M$ for $x > b$. Hence, $\int_b^\infty e^{sx} e^{-f(x)} dx \leq \int_b^\infty e^{(s-M)x} dx = \int_b^\infty e^{-(M-s)|x|} dx$. Similarly, there is an $a < 0$ such that $f(x)/(-x) > M$ for $x < a$. Hence, $\int_{-\infty}^a e^{sx} e^{-f(x)} dx \leq \int_{-\infty}^a e^{(s+M)x} dx = \int_{-\infty}^a e^{-(M+s)|x|} dx$. Since f is continuous, f is bounded on (a, b) , hence, integrable over (a, b) . Thus, with $g(x) = e^{sx} e^{-f(x)}$,

$$\int_{-\infty}^\infty e^{sx} e^{-f(x)} dx = \int_{-\infty}^a g(x) dx + \int_a^b g(x) dx + \int_b^\infty g(x) dx$$

$$\leq \int_{-\infty}^a e^{-(M+s)|x|} dx + \int_a^b e^{sx} e^{-f(x)} dx + \int_b^{\infty} e^{-(M-s)|x|} dx,$$

which is finite, as soon as M is chosen $> |s|$.

4.3.12. Suppose that there were such a δ . First, choose $f(x) = 1$, for all $x \in \mathbb{R}$, in (3.12) to get $\int_{-\infty}^{\infty} \delta(x) dx = 1$. Thus, δ is integrable over \mathbb{R} . Now, let f equal 1 at all points except at zero, where we set $f(0) = 0$. Then, (3.12) fails because the integral is still 1, but the right side vanishes. However, this is of no use, since we are assuming (3.12) for f continuous only. Because of this, we let $f_n(x) = 1$ for $|x| \geq 1/n$ and $f_n(x) = n|x|$ for $|x| \leq 1/n$. Then, f_n is continuous and nonnegative. Hence, by monotonicity,

$$0 \leq \int_{|x| \geq 1/n} \delta(x) dx \leq \int_{-\infty}^{\infty} \delta(x) f_n(x) dx = f_n(0) = 0,$$

for all $n \geq 1$. This shows that $\int_{|x| \geq 1/n} \delta(x) dx = 0$ for all $n \geq 1$. But, by continuity at the endpoints,

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \delta(x) dx = \int_{-\infty}^0 \delta(x) dx + \int_0^{\infty} \delta(x) dx \\ &= \lim_{n \nearrow \infty} \left(\int_{-\infty}^{-1/n} \delta(x) dx + \int_{1/n}^{\infty} \delta(x) dx \right) \\ &= \lim_{n \nearrow \infty} (0 + 0) = 0, \end{aligned}$$

a contradiction.

Exercises 4.4

4.4.1. $F(x) = e^{-sx}/(-s)$ is a primitive of $f(x) = e^{-sx}$, and e^{-sx} is positive. So,

$$\int_0^{\infty} e^{-sx} dx = \frac{1}{-s} e^{-sx} \Big|_0^{\infty} = \frac{1}{s}, \quad s > 0.$$

4.4.2. x^r/r is a primitive of x^{r-1} for $r \neq 0$, and $\log x$ is a primitive, when $r = 0$. Thus, $\int_0^1 dx/x = \log x|_0^1 = 0 - (-\infty) = \infty$ and $\int_1^{\infty} dx/x = \log x|_1^{\infty} = \infty - 0 = \infty$. Hence, all three integrals are equal to ∞ , when $r = 0$. Now,

$$\int_0^1 x^{r-1} dx = \frac{1}{r} - \frac{1}{r} \lim_{x \rightarrow 0^+} x^r = \begin{cases} \frac{1}{r}, & r > 0, \\ \infty, & r < 0. \end{cases}$$

Also,

$$\int_1^{\infty} x^{r-1} dx = \frac{1}{r} \lim_{x \rightarrow \infty} x^r - \frac{1}{r} = \begin{cases} -\frac{1}{r}, & r < 0, \\ \infty, & r > 0. \end{cases}$$

Since $\int_0^{\infty} = \int_0^1 + \int_1^{\infty}$, $\int_0^{\infty} x^{r-1} dx = \infty$ in all cases.

4.4.3. Pick c in (a, b) . Since any primitive F differs from F_c by a constant, it is enough to verify the result for F_c . But f bounded and (a, b) bounded implies f integrable. So, $F_c(a+)$ and $F_c(b-)$ exist and are finite by continuity at the endpoints.

4.4.4. Take $u = 1/x$ and $dv = \sin x \, dx$. Then, $du = -dx/x^2$ and $v = -\cos x$. So,

$$\int_1^b \frac{\sin x}{x} \, dx = \left. \frac{-\cos x}{x} \right|_1^b + \int_1^b \frac{\cos x}{x^2} \, dx.$$

But, by (3.1), $\cos x/x^2$ is integrable over $(1, \infty)$. So, by continuity at the endpoints,

$$\lim_{b \rightarrow \infty} \int_1^b \frac{\sin x}{x} \, dx = \cos 1 + \int_1^{\infty} \frac{\cos x}{x^2} \, dx.$$

Since $F(b) - \int_1^b \sin x(dx/x)$ does not depend on b , $F(\infty)$ exists and is finite.

4.4.5. The function $g(t) = e^{-t}$ is strictly monotone with $g((0, \infty)) = (0, 1)$. Now, apply substitution.

4.4.6. Let $u = x^n$ and $dv = e^{-sx} \, dx$. Then, $du = nx^{n-1} \, dx$, and $v = e^{-sx}/(-s)$. Hence,

$$\int_0^{\infty} e^{-sx} x^n \, dx = \left. \frac{e^{-sx} x^n}{-s} \right|_0^{\infty} + \frac{n}{s} \int_0^{\infty} e^{-sx} x^{n-1} \, dx.$$

If we call the integral on the left I_n , this says that $I_n = (n/s)I_{n-1}$. Iterating this down to $n = 0$ yields $I_n = n!/s^n$ since $I_0 = 1/s$ from Exercise 1.

4.4.7. Call the integrals I_s and I_c . Let $u = \sin(sx)$ and $dv = e^{-nx} \, dx$. Then, $du = s \cos(sx) \, dx$ and $v = e^{-nx}/(-n)$. So,

$$I_s = \left. \frac{e^{-nx} \sin(sx)}{-n} \right|_0^{\infty} + \frac{s}{n} \int_0^{\infty} e^{-nx} \cos(sx) \, dx = \frac{s}{n} I_c.$$

Now, let $u = \cos(sx)$ and $dv = e^{-nx} \, dx$. Then, $du = -s \sin(sx) \, dx$ and $v = e^{-nx}/(-n)$. So

$$I_c = \left. \frac{e^{-nx} \cos(sx)}{-n} \right|_0^{\infty} - \frac{s}{n} \int_0^{\infty} e^{-nx} \sin(sx) \, dx = \frac{1}{n} - \frac{s}{n} I_s.$$

Thus, $nI_s = sI_c$, and $nI_c = 1 - sI_s$. Solving, we obtain $I_s = s/(n^2 + s^2)$ and $I_c = n/(n^2 + s^2)$.

4.4.8. Let $u = t^{x-1}$ and $dv = e^{-t^2/2} t \, dt$. Then, $du = (x-1)t^{x-2} \, dt$, and $v = -e^{-t^2/2}$. So,

$$\begin{aligned} \int_0^{\infty} e^{-t^2/2} t^x \, dx &= \left. -e^{-t^2/2} t^{x-1} \right|_0^{\infty} + (x-1) \int_0^{\infty} e^{-t^2/2} t^{x-2} \, dt \\ &= (x-1) \int_0^{\infty} e^{-t^2/2} t^{x-2} \, dt. \end{aligned}$$

If $I_n = \int_0^\infty e^{-t^2/2} t^n dt$, then,

$$\begin{aligned} I_{2n+1} &= 2n \cdot I_{2n-1} = 2n \cdot (2n-2) I_{2n-3} \\ &= \cdots = 2n \cdot (2n-2) \cdots 4 \cdot 2 \cdot I_1 = 2^n n! I_1. \end{aligned}$$

But, substituting $u = t^2/2$, $du = t dt$ yields

$$I_1 = \int_0^\infty e^{-t^2/2} t dt = \int_0^\infty e^{-u} du = 1.$$

4.4.9. Let $u = (1-t)^n$ and $dv = t^{x-1} dt$. Then, $du = -n(1-t)^{n-1} dt$, and $v = t^x/x$. So,

$$\begin{aligned} \int_0^1 (1-t)^n t^{x-1} dt &= \left. \frac{(1-t)^n t^x}{x} \right|_0^1 + \frac{n}{x} \int_0^1 (1-t)^{n-1} t^{x+1} dt \\ &= \frac{n}{x} \int_0^1 (1-t)^{n-1} t^{x+1} dt. \end{aligned}$$

Thus, integrating by parts increases the x by 1 and decreases the n by 1. Iterating this n times,

$$\int_0^1 (1-t)^n t^{x-1} dt = \frac{n \cdot (n-1) \cdots 1}{x \cdot (x+1) \cdots (x+n-1)} \cdot \int_0^1 t^{x+n-1} dt.$$

But $\int_0^1 t^{x+n-1} dt = 1/(x+n)$. So,

$$\int_0^1 (1-t)^n t^{x-1} dt = \frac{n!}{x \cdot (x+1) \cdots (x+n)}.$$

4.4.10. Let $t = -\log x$, $x = e^{-t}$. Then, from Exercises 5 and 6,

$$\int_0^1 (-\log x)^n dx = \int_0^\infty e^{-t} t^n dt = n!.$$

4.4.11. Let $I_n = \int_{-1}^1 (x^2 - 1)^n dx$ and $u = (x^2 - 1)^n$, $dv = dx$. Then,

$$\begin{aligned} I_n &= x(x^2 - 1)^n \Big|_{-1}^1 - 2n \int_{-1}^1 (x^2 - 1)^{n-1} x^2 dx \\ &= -2n I_n - 2n I_{n-1}. \end{aligned}$$

Solving for I_n , we obtain

$$I_n = -\frac{2n}{2n+1} I_{n-1} = \cdots = (-1)^n \frac{2n \cdot (2n-2) \cdots 2}{(2n+1) \cdot (2n-1) \cdots 3} \cdot 2$$

since $I_0 = 2$.

4.4.12. Let $f(x) = (x^2 - 1)^n$. Then, $P_n(x) = f^{(n)}(x)/2^n n!$. Note that $f(\pm 1) = 0$, $f'(\pm 1) = 0$, ..., and $f^{(n-1)}(\pm 1) = 0$, since all these derivatives have at least one factor $(x^2 - 1)$ by the product rule. Hence, integrating by parts,

$$\int_{-1}^1 f^{(n)}(x) f^{(n)}(x) dx = - \int_{-1}^1 f^{(n-1)}(x) f^{(n+1)}(x) dx$$

increases one index and decreases the other. Iterating, we get

$$\int_{-1}^1 P_n(x)^2 dx = \frac{1}{(2^n n!)^2} \int_{-1}^1 [f^{(n)}(x)]^2 dx = \frac{(-1)^n}{(2^n n!)^2} \int_{-1}^1 f(x) f^{(2n)}(x) dx.$$

But f is a polynomial of degree $2n$ with highest order coefficient 1. Hence, $f^{(2n)}(x) = (2n)!$. So,

$$\int_{-1}^1 P_n(x)^2 dx = \frac{(-1)^n (2n)!}{(2^n n!)^2} \int_{-1}^1 (x^2 - 1)^n dx.$$

Now, inserting the result of the previous Exercise and simplifying leads to $2/(2n+1)$.

4.4.13. By the integral test, $\zeta(s)$ differs from $\int_1^\infty x^{-s} dx$ by, at most, 1. But $\int_1^\infty x^{-s} dx$ converges for $s > 1$ by Exercise 2.

4.4.14. Here, $f(x) = 1/x$ and $\int_1^{n+1} f(x) dx = \log(n+1)$. Since $1/(n+1) \rightarrow 0$, by the integral test,

$$\begin{aligned} \gamma &= \lim_{n \nearrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n \right) \\ &= \lim_{n \nearrow \infty} \left[1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} + \frac{1}{n+1} - \log(n+1) \right] \\ &= \lim_{n \nearrow \infty} \left[f(1) + f(2) + \cdots + f(n) - \int_1^{n+1} f(x) dx \right] \end{aligned}$$

exists and satisfies $0 < \gamma < 1$.

4.4.15. Call the integrals I_n^c and I_n^s . Then, $I_0^s = 0$, and $I_n^c = 0$, $n \geq 0$, since the integrand is odd. Also, $I_n^s = 2 \int_0^\pi x \sin(nx) dx$ since the integrand is even. Now, for $n \geq 1$,

$$\begin{aligned} \int_0^\pi x \sin(nx) dx &= -\frac{x \cos(nx)}{n} \Big|_0^\pi + \frac{1}{n} \int_0^\pi \cos(nx) dx \\ &= -\frac{\pi \cos(n\pi)}{n} + \frac{1}{n} \cdot \frac{\sin(nx)}{n} \Big|_0^\pi \\ &= \frac{(-1)^{n-1} \pi}{n}. \end{aligned}$$

Thus, $I_n^s = 2\pi(-1)^{n-1}/n$, $n \geq 1$.

4.4.16. By oddness, $\int_{-\pi}^\pi \sin(nx) \cos(mx) dx = 0$ for all $m, n \geq 0$. For $m \neq n$, using (5.3) in §3.5,

$$\begin{aligned} \int_{-\pi}^\pi \sin(nx) \sin(mx) dx &= \frac{1}{2} \int_{-\pi}^\pi [\cos((n-m)x) - \cos((n+m)x)] dx \\ &= \frac{1}{2} \left(\frac{\sin((n-m)x)}{n-m} - \frac{\sin((n+m)x)}{n+m} \right) \Big|_{-\pi}^\pi \end{aligned}$$

$$= 0.$$

For $m = n$,

$$\begin{aligned}\int_{-\pi}^{\pi} \sin(nx) \sin(mx) dx &= \frac{1}{2} \int_{-\pi}^{\pi} [1 - \cos(2nx)] dx \\ &= \frac{1}{2} [x - \sin(2nx)/2n] \Big|_{-\pi}^{\pi} = \pi.\end{aligned}$$

Similarly, for $\int_{-\pi}^{\pi} \cos(nx) \cos(mx) dx$. Hence,

$$\int_{-\pi}^{\pi} \cos(nx) \cos(mx) dx = \int_{-\pi}^{\pi} \sin(nx) \sin(mx) dx = \begin{cases} 0, & n \neq m, \\ \pi, & n = m. \end{cases}$$

4.4.17. By linearity, $q(t) = at^2 + 2bt + c$ where $a = \int_a^b g(t)^2 dt$, $b = \int_a^b f(t)g(t) dt$, and $c = \int_a^b f(t)^2 dt$. Since q is nonnegative, q has, at most, one root. Hence, $b^2 - ac \leq 0$, which is Cauchy–Schwarz.

4.4.18. By substituting $t = n(1 - s)$, $dt = nds$, and equation (3.2) in §2.3,

$$\begin{aligned}\int_0^n \frac{1 - (1 - t)^n}{t} dt &= \int_0^1 \frac{1 - s^n}{1 - s} ds \\ &= \int_0^1 [1 + s + \cdots + s^{n-1}] ds \\ &= 1 + \frac{1}{2} + \cdots + \frac{1}{n}.\end{aligned}$$

4.4.19. Continuity of F was established in §4.3. Now, f is continuous on the subinterval (x_{i-1}, x_i) . Hence, $F(x) = F(x_{i-1}) + F_{x_{i-1}}(x)$ is differentiable by the (first version of) the fundamental theorem.

4.4.20. For any f , let $I(f) = \int_a^b f(x) dx$. If $g : [a, b] \rightarrow \mathbb{R}$ is nonnegative and continuous, we can (§2.3) find a piecewise constant $g_\epsilon \geq 0$, such that $g_\epsilon(x) \leq g(x) + \epsilon \leq g_\epsilon(x) + 2\epsilon$ on $a \leq x \leq b$. By monotonicity,

$$I(g_\epsilon) \leq I(g + \epsilon) \leq I(g_\epsilon + 2\epsilon).$$

But, by Exercise 5 of §4.3, $I(g + \epsilon) = I(g) + \epsilon(b - a)$ and $I(g_\epsilon + 2\epsilon) = I(g_\epsilon) + 2\epsilon(b - a)$. Hence,

$$I(g_\epsilon) \leq I(g) + \epsilon(b - a) \leq I(g_\epsilon) + 2\epsilon(b - a),$$

or

$$|I(g) - I(g_\epsilon)| \leq \epsilon(b - a).$$

Similarly, since $f(x) + g_\epsilon(x) \leq f(x) + g(x) + \epsilon \leq f(x) + g_\epsilon(x) + 2\epsilon$,

$$|I(f + g) - I(f) - I(g_\epsilon)| \leq \epsilon(b - a),$$

where we have used Exercise 5 of §4.3, again. Thus, $|I(f + g) - I(f) - I(g)| \leq 2\epsilon(b - a)$. Since ϵ is arbitrary, we conclude that $I(f + g) = I(f) + I(g)$.

4.4.21. Let $m_i = g(t_i)$, $i = 0, 1, \dots, n+1$. For each $i = 1, \dots, n+1$, define $\#_i : (m, M) \rightarrow \{0, 1\}$ by setting $\#_i(x) = 1$ if x is between m_{i-1} and m_i and $\#_i(x) = 0$, otherwise. Since the m_i 's may not be increasing, for a given x , more than one $\#_i(x)$, $i = 1, \dots, n+1$, may equal one. In fact, for any x not equal to the m_i 's,

$$\#(x) = \#_1(x) + \dots + \#_{n+1}(x).$$

Since G is strictly monotone on (t_{i-1}, t_i) ,

$$\int_{t_{i-1}}^{t_i} f(g(t))|g'(t)| dt = \int_{m_{i-1}}^{m_i} f(x) dx = \int_m^M f(x)\#_i(x) dx.$$

Now, add these equations over $1 \leq i \leq n+1$ to get

$$\begin{aligned} \int_a^b f(g(t))|g'(t)| dt &= \sum_{i=1}^{n+1} \int_m^M f(x)\#_i(x) dx \\ &= \int_m^M \sum_{i=1}^{n+1} f(x)\#_i(x) dx = \int_m^M f(x)\#(x) dx. \end{aligned}$$

Here, the last equality follows from the fact that $\#(x)$ and $\sum_{i=1}^{n+1} \#_i(x)$ differ only on finitely many points in (m, M) .

4.4.22. If $a = x_0 < x_1 < \dots < x_{n+1} = b$ is a partition, then,

$$|f(x_i) - f(x_{i-1})| = \left| \int_{x_{i-1}}^{x_i} f'(x) dx \right| \leq \int_{x_{i-1}}^{x_i} |f'(x)| dx$$

by the fundamental theorem. Summing this over $1 \leq i \leq n+1$ yields the first part. Also, since this was any partition, taking the sup over all partitions shows that the total variation is $\leq \int_a^b |f'(x)| dx$. Call this last integral I . To show that the total variation equals I , given ϵ , we will exhibit a partition whose variation is within ϵ of I . Now, since $|f'|$ is continuous over $[a, b]$, $|f'|$ is Riemann integrable. Hence (Exercise 3 of §4.3), given $\epsilon > 0$, there is a partition $a = x_0 < x_1 < \dots < x_{n+1} = b$ whose corresponding Riemann sum $\sum_{i=1}^{n+1} |f'(x_i^\#)|(x_i - x_{i-1})$ is within ϵ of I , for any choice of intermediate points $x_i^\#$, $i = 1, \dots, n+1$. But, by the mean value theorem,

$$\sum_{i=1}^{n+1} |f(x_i) - f(x_{i-1})| = \sum_{i=1}^{n+1} |f'(x_i^\#)|(x_i - x_{i-1})$$

for some intermediate points $x_i^\#$, $i = 1, \dots, n+1$. Thus, the variation of this partition is within ϵ of I .

Exercises 4.5

4.5.1. Let $Q = (a, b) \times (c, d)$. If $(x, y) \in Q$ and $(x', y') \notin Q$, then, the distance from (x, y) to (x', y') is no smaller than the distance from (x, y) to the boundary of Q , which, in turn, is no smaller than the minimum of $|x - a|$ and $|x - b|$.

4.5.2. Let $Q_n = (-1/n, 1/n) \times (-1/n, 1/n)$, $n \geq 1$. Then, Q_n is an open set for each $n \geq 1$, and $\bigcap_{n=1}^{\infty} Q_n$ is a single point $\{(0, 0)\}$, which is not open.

4.5.3. If Q is compact, then, Q^c is a union of four open rectangles. So, Q^c is open. So, Q is closed. If C_n , $n \geq 1$, is closed, then, C_n^c is open. So,

$$\left(\bigcap_{n=1}^{\infty} C_n \right)^c = \bigcup_{n=1}^{\infty} C_n^c$$

is open. Hence, $\bigcap_{n=1}^{\infty} C_n$ is closed. Let $Q_n = [0, 1] \times [1/n, 1]$, $n \geq 1$. Then, Q_n is closed, but

$$\bigcup_{n=1}^{\infty} Q_n = [0, 1] \times (0, 1]$$

is not.

4.5.4. It is enough to show that C^c is open. But C^c is the union of the four sets (draw a picture) $(-\infty, a) \times \mathbb{R}$, $(b, \infty) \times \mathbb{R}$, $(a, b) \times (-\infty, 0)$, and $\{(x, y) : a < x < b, y > f(x)\}$. The first three sets are clearly open, whereas the fourth is shown to be open using the continuity of f , exactly as in the text. Thus, C is closed. Since C contains the subgraph of f , $\text{area}(C) \geq \int_a^b f(x) dx$. On the other hand, C is contained in the union of the subgraph of $f + \epsilon/(1 + x^2)$ with L_a and L_b . Thus,

$$\begin{aligned} \text{area}(C) &\leq \int_a^b [f(x) + \epsilon/(1 + x^2)] dx + \text{area}(L_a) + \text{area}(L_b) \\ &= \int_a^b f(x) dx + \epsilon \int_a^b \frac{dx}{1 + x^2} \\ &\leq \int_a^b f(x) dx + \epsilon \int_{-\infty}^{\infty} \frac{dx}{1 + x^2} \\ &= \int_a^b f(x) dx + \epsilon\pi. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, the result follows.

4.5.5. Distance is always nonnegative, so (\iff means iff),

$$\begin{aligned} C \text{ is closed} &\iff C^c \text{ is open} \\ &\iff d((x, y), (C^c)^c) > 0 \text{ iff } (x, y) \in C^c \\ &\iff d((x, y), C) > 0 \text{ iff } (x, y) \in C^c \\ &\iff d((x, y), C) = 0 \text{ iff } (x, y) \in C. \end{aligned}$$

This is the first part. For the second, let $(x, y) \in G_n$. If $\alpha = 1/n - d((x, y), C) > 0$ and $|x - x'| < \epsilon$, $|y - y'| < \epsilon$, then, $d((x', y'), C) \leq d((x, y), C) + 2\epsilon = 1/n + 2\epsilon - \alpha$, by the triangle inequality. Thus, for $\epsilon < \alpha/2$, $(x', y') \in G_n$. This shows that $Q_\epsilon \subset G_n$, where Q_ϵ is the open rectangle centered at (x, y) with sides

of length 2ϵ . Thus, G_n is open, and $\bigcap_{n=1}^{\infty} G_n = \{(x, y) : d((x, y), C) = 0\}$, which equals C .

4.5.6. Given $\epsilon > 0$, we have to find an open superset G of A satisfying $\text{area}(G) \leq \text{area}(A) + \epsilon$. If $\text{area}(A) = \infty$, $G = \mathbb{R}^2$ will do. If $\text{area}(A) < \infty$, choose a paving (Q_n) , such that $\sum_{n=1}^{\infty} \text{area}(Q_n) < \text{area}(A) + \epsilon$. Then, $G = \bigcup_{n=1}^{\infty} Q_n$ is open, G contains A , and

$$\text{area}(G) = \text{area}\left(\bigcup_{n=1}^{\infty} Q_n\right) \leq \sum_{n=1}^{\infty} \text{area}(Q_n) < \text{area}(A) + \epsilon.$$

For the second part, let $\alpha = \inf\{\text{area}(G) : A \subset G, G \text{ open}\}$. Choosing G as above, $\alpha \leq \text{area}(G) \leq \text{area}(A) + \epsilon$ for all ϵ . Hence, $\alpha \leq \text{area}(A)$. Conversely, monotonicity implies that $\text{area}(A) \leq \text{area}(G)$ for any superset G . Hence, $\text{area}(A) \leq \alpha$.

4.5.7. For each $\epsilon > 0$, by Exercise 6, choose G_ϵ open such that $A \subset G_\epsilon$ and $\text{area}(G_\epsilon) \leq \text{area}(A) + \epsilon$. Let $I = \bigcap_{n=1}^{\infty} G_{1/n}$. Then, I is interopen, and $\text{area}(I) \leq \inf_{n \geq 1} \text{area}(G_{1/n}) \leq \inf_{n \geq 1} (\text{area}(A) + 1/n) = \text{area}(A)$. But $I \supset A$. So, $\text{area}(I) \geq \text{area}(A)$.

4.5.8. We already know that the intersection of a sequence of measurable sets is measurable. By De Morgan's law (§1.1), M_n measurable implies the complement M_n^c is measurable. So,

$$\left(\bigcup_{n=1}^{\infty} M_n\right)^c = \bigcap_{n=1}^{\infty} M_n^c$$

is measurable. So, the complement $\bigcup_{n=1}^{\infty} M_n$ is measurable.

4.5.9. Let C_n and C be as in §4.1. Since C_n is a finite union of compact rectangles, C_n is closed. Since $C = \bigcap_{n=1}^{\infty} C_n$, C is closed.

4.5.10. Let P_k , $k = 0, \dots, n$, denote the vertices of D'_n . It is enough to show that the closest approach to O of the line joining P_k and P_{k+1} is at the midpoint $M = (P_k + P_{k+1})/2$, where the distance to O equals 1. Let $\theta_k = k\pi/n$. Then, the distance squared from the midpoint to O is given by

$$\begin{aligned} & \frac{[\cos(2\theta_k) + \cos(2\theta_{k+1})]^2 + [\sin(2\theta_k) + \sin(2\theta_{k+1})]^2}{4\cos(\theta_1)^2} \\ &= \frac{2 + 2[\cos(2\theta_k)\cos(2\theta_{k+1}) + \sin(2\theta_k)\sin(2\theta_{k+1})]}{4\cos^2(\theta_1)} \\ &= \frac{2 + 2\cos(2\theta_1)}{4\cos^2(\theta_1)} = 1. \end{aligned}$$

Thus, the distance to the midpoint is 1. To show that this is the minimum, check that the line segments OM and P_kP_{k+1} are perpendicular.

4.5.11. Here, $a_n = n \sin(\pi/n) \cos(\pi/n)$, and $a'_n = n \tan(\pi/n)$. So, $a_n a'_n = n^2 \sin^2(\pi/n)$. But $a_{2n} = 2n \sin(\pi/2n) \cos(\pi/2n) = n \sin(\pi/n)$. So, $a_{2n} =$

$\sqrt{a_n a'_n}$. Also,

$$\begin{aligned} \frac{1}{a_{2n}} + \frac{1}{a'_n} &= \frac{1}{n \sin(\pi/n)} + \frac{1}{n \tan(\pi/n)} \\ &= \frac{\cos(\pi/n) + 1}{n \sin(\pi/n)} \\ &= \frac{2 \cos^2(\pi/2n)}{2n \sin(\pi/2n) \cos(\pi/2n)} \\ &= \frac{2}{2n \tan(\pi/2n)} = \frac{2}{a'_{2n}}. \end{aligned}$$

4.5.12. In the definition of measurable, replace M and A by $A \cup B$ and A , respectively. Then, $A \cap M$ is replaced by A , and $A \cap M^c$ is replaced by B .

4.5.13. From the previous exercise and induction,

$$\text{area} \left(\bigcup_{n=1}^{\infty} A_n \right) \geq \text{area} \left(\bigcup_{n=1}^N A_n \right) = \sum_{n=1}^N \text{area}(A_n).$$

Let $N \nearrow \infty$ to get

$$\text{area} \left(\bigcup_{n=1}^{\infty} A_n \right) \geq \sum_{n=1}^{\infty} \text{area}(A_n).$$

Since the reverse inequality follows from subadditivity, we are done.

4.5.14. Note that A and $B \setminus A$ are disjoint and their union is $A \cup B$. But $B \setminus A$ and $A \cap B$ are disjoint and their union is B . So,

$$\begin{aligned} \text{area}(A \cup B) &= \text{area}(A) + \text{area}(B \setminus A) \\ &= \text{area}(A) + \text{area}(B) - \text{area}(A \cap B). \end{aligned}$$

4.5.15.

$$\begin{aligned} \text{area}(A \cup B \cup C) &= \text{area}(A) + \text{area}(B) + \text{area}(C) \\ &\quad - \text{area}(A \cap B) - \text{area}(B \cap C) - \text{area}(A \cap C) \\ &\quad + \text{area}(A \cap B \cap C). \end{aligned}$$

The general formula, the *inclusion-exclusion principle*, is that the area of a union equals the sum of the areas of the sets minus the sum of the areas of their double intersections plus the sum of the areas of their triple intersections minus the sum of the areas of their quadruple intersections, etc.

4.5.16. By Exercise 6, given $\epsilon > 0$, there is an open superset G of M satisfying $\text{area}(G) \leq \text{area}(M) + \epsilon$. If M is measurable and $\text{area}(M) < \infty$, replace A in (5.4) by G to get $\text{area}(G) = \text{area}(M) + \text{area}(G \setminus M)$. Hence, $\text{area}(G \setminus M) \leq \epsilon$. If $\text{area}(M) = \infty$, write $M = \bigcup_{n=1}^{\infty} M_n$ with $\text{area}(M_n) < \infty$ for all $n \geq 1$. For each $n \geq 1$, choose an open superset G_n of M_n satisfying $\text{area}(G_n \setminus M_n) \leq$

$\epsilon 2^{-n}$. Then, $G = \bigcup_{n=1}^{\infty} G_n$ is an open superset of M , and $\text{area}(G \setminus M) \leq \sum_{n=1}^{\infty} \text{area}(G_n \setminus M_n) \leq \epsilon$. This completes the first part. Conversely, suppose, for all $\epsilon > 0$, there is an open superset G of M satisfying $\text{area}(G \setminus M) \leq \epsilon$, and let A be arbitrary. Since G is measurable,

$$\begin{aligned} \text{area}(A \cap M) + \text{area}(A \cap M^c) & \\ & \leq \text{area}(A \cap G) + \text{area}(A \cap G^c) + \text{area}(A \cap (G \setminus M)) \\ & \leq \text{area}(A) + \epsilon. \end{aligned}$$

Thus, $\text{area}(A \cap M) + \text{area}(A \cap M^c) \leq \text{area}(A)$. Since the reverse inequality follows by subadditivity, M is measurable.

4.5.17. When A is a rectangle, the result is obvious (draw a picture). In fact, for a rectangle Q and $Q' = Q + (a, b)$, $\text{area}(Q \cap Q') \geq (1 - \epsilon)^2 \text{area}(Q)$. To deal with general A , let Q be as in Exercise 15 of §4.2 with α to be determined below, and let $A' = A + (a, b)$. Then, by subadditivity and translation invariance,

$$\begin{aligned} \text{area}(Q \cap Q') & \leq \text{area}((Q \cap A) \cap (Q' \cap A')) \\ & \quad + \text{area}(Q \setminus (Q \cap A)) + \text{area}(Q' \setminus (Q' \cap A')) \\ & = \text{area}((Q \cap A) \cap (Q' \cap A')) + 2 \cdot \text{area}(Q \setminus (Q \cap A)) \\ & \leq \text{area}(A \cap A') + 2 \cdot \text{area}(Q \setminus (Q \cap A)). \end{aligned}$$

But, from Exercise 15 of §4.2 and the measurability of A , $\text{area}[Q \setminus (Q \cap A)] < (1 - \alpha) \text{area}(Q)$. Hence,

$$\begin{aligned} \text{area}(A \cap A') & \geq \text{area}(Q \cap Q') - 2(1 - \alpha) \text{area}(Q) \\ & \geq (1 - \epsilon)^2 \text{area}(Q) - 2(1 - \alpha) \text{area}(Q). \end{aligned}$$

Thus, the result follows as soon as one chooses $2(1 - \alpha) < (1 - \epsilon)^2$.

4.5.18. If $\text{area}[A \cap (A + (a, b))] > 0$, then, $A \cap (A + (a, b))$ is nonempty. If $(x, y) \in A \cap (A + (a, b))$, then, $(x, y) = (x', y') + (a, b)$ with $(x', y') \in A$. Hence, $(a, b) \in A - A$. Since Exercise 17 says that $\text{area}[A \cap (A + (a, b))] > 0$ for all $(a, b) \in Q_\epsilon$, the result follows.

A.5 Chapter 5

Exercises 5.1

5.1.1. Let $f_n(x) = 1/n$ for all $x \in \mathbb{R}$. Then, $\int_{-\infty}^{\infty} f_n(x) dx = \infty$ for all $n \geq 1$, and $f_n(x) \searrow f(x) = 0$.

5.1.2. Let $I_n = \int_a^b f_n(x) dx$ and $I = \int_a^b f(x) dx$. We have to show that $I_* \geq I$. The lower sequence is

$$g_n(x) = \inf\{f_k(x) : k \geq n\}, \quad n \geq 1.$$

Then, $(g_n(x))$ is nonnegative and increasing to $f(x)$, $a < x < b$. So, the monotone convergence theorem applies. So, $J_n = \int_a^b g_n(x) dx \rightarrow \int_a^b f(x) dx = I$. Since $f_n(x) \geq g_n(x)$, $a < x < b$, $I_n \geq J_n$. Hence, $I_* \geq J_* = I$.

5.1.3. Given x fixed, $|x - n| \geq 1$ for n large enough. Hence, $f_0(x - n) = 0$. Hence, $f_n(x) = 0$ for n large enough. Thus, $f(x) = \lim_{n \nearrow \infty} f_n(x) = 0$. But, by translation invariance,

$$\int_{-\infty}^{\infty} f_n(x) dx = \int_{-\infty}^{\infty} h(x) dx = \int_{-1}^1 [1 - x^2] dx = \frac{4}{3} > 0.$$

Since $\int_{-\infty}^{\infty} f(x) dx = 0$, here, the inequality in Fatou's lemma is strict.

5.1.4. By Exercise 4 of §3.2, $(1 - t/n)^n \nearrow e^{-t}$ as $n \nearrow \infty$. To take care of the upper limit of integration that changes with n , let

$$f_n(t) = \begin{cases} \left(1 - \frac{t}{n}\right)^n t^{x-1}, & 0 < t < n, \\ 0, & t \geq n. \end{cases}$$

Then, by the monotone convergence theorem,

$$\begin{aligned} \Gamma(x) &= \int_0^{\infty} e^{-t} t^{x-1} dt \\ &= \int_0^{\infty} \lim_{n \nearrow \infty} f_n(t) dt \\ &= \lim_{n \nearrow \infty} \int_0^{\infty} f_n(t) dt \\ &= \lim_{n \nearrow \infty} \int_0^n \left(1 - \frac{t}{n}\right)^n t^{x-1} dt. \end{aligned}$$

5.1.5. By Exercise 9 of §4.4,

$$\int_0^n \left(1 - \frac{t}{n}\right)^n t^{x-1} dt = n^x \int_0^1 (1-s)^n s^{x-1} ds = \frac{n^x n!}{x \cdot (x+1) \cdot \dots \cdot (x+n)}.$$

5.1.6. Convexity of f_n means that $f_n((1-t)x + ty) \leq (1-t)f_n(x) + tf_n(y)$ for all $a < x < y < b$ and $0 \leq t \leq 1$. Letting $n \nearrow \infty$, we obtain $f((1-t)x + ty) \leq (1-t)f(x) + tf(y)$ for all $a < x < y < b$ and $0 \leq t \leq 1$, which says that f is convex. Now, let

$$f_n(x) = \log \left(\frac{n^x n!}{x \cdot (x+1) \cdot \dots \cdot (x+n)} \right).$$

Then, $\log \Gamma(x) = \lim_{n \nearrow \infty} f_n(x)$ by (1.3) and

$$\frac{d^2}{dx^2} f_n(x) = \frac{d^2}{dx^2} \left(x \log n + \log(n!) - \sum_{k=0}^n \log(x+k) \right) = \sum_{k=0}^n \frac{1}{(x+k)^2},$$

which is positive. Thus, f_n is convex. So, $\log \Gamma$ is convex.

5.1.7. Since $\log \Gamma(x)$ is convex,

$$\log \Gamma((1-t)x + ty) \leq (1-t) \log \Gamma(x) + t \log \Gamma(y)$$

for $0 < x < y < \infty$, $0 \leq t \leq 1$. Since e^x is convex and increasing,

$$\begin{aligned} \Gamma((1-t)x + ty) &= \exp(\log \Gamma((1-t)x + ty)) \\ &\leq \exp((1-t) \log \Gamma(x) + t \log \Gamma(y)) \\ &\leq (1-t) \exp(\log \Gamma(x)) + t \exp(\log \Gamma(y)) \\ &= (1-t)\Gamma(x) + t\Gamma(y) \end{aligned}$$

for $0 < x < y < \infty$ and $0 \leq t \leq 1$.

5.1.8. Use summation under the integral sign, with $f_n(t) = t^{x-1}e^{-nt}$, $n \geq 1$. Then, substituting $s = nt$, $ds = n dt$,

$$\begin{aligned} \int_0^\infty \frac{t^{x-1}}{e^t - 1} dt &= \int_0^\infty \sum_{n=1}^\infty t^{x-1} e^{-nt} dt \\ &= \sum_{n=1}^\infty \int_0^\infty t^{x-1} e^{-nt} dt \\ &= \sum_{n=1}^\infty n^{-x} \int_0^\infty s^{x-1} e^{-s} ds \\ &= \zeta(x)\Gamma(x). \end{aligned}$$

5.1.9. Use summation under the integral sign, with $f_n(t) = t^{x-1}e^{-n^2\pi t}$, $n \geq 1$. Then, substituting $s = n^2\pi t$, $ds = n^2\pi dt$,

$$\begin{aligned} \int_0^\infty \psi(t) t^{x/2-1} dt &= \int_0^\infty \sum_{n=1}^\infty e^{-n^2\pi t} t^{x/2-1} dt \\ &= \sum_{n=1}^\infty \int_0^\infty e^{-n^2\pi t} t^{x/2-1} dt \\ &= \sum_{n=1}^\infty \pi^{-x/2} n^{-x} \int_0^\infty e^{-s} s^{x/2-1} ds \\ &= \pi^{-x/2} \zeta(x)\Gamma(x/2). \end{aligned}$$

5.1.10. By Exercise 5 of §4.4,

$$\int_0^1 t^{x-1} (-\log t)^{n-1} dt = \int_0^\infty e^{-xs} s^{n-1} ds = \frac{\Gamma(n)}{x^n}.$$

Here, we used the substitutions $t = e^{-s}$, then, $xs = r$.

5.1.11. Recall that $\log t < 0$ on $(0, 1)$ and $0 < \log t < t$ on $(1, \infty)$. From the previous Exercise, with $f(t) = e^{-t} t^{x-1} |\log t|^{n-1}$,

$$\begin{aligned} \int_0^\infty e^{-t} t^{x-1} |\log t|^{n-1} dt &= \int_0^1 f(t) dt + \int_1^\infty f(t) dt \\ &\leq \int_0^1 t^{x-1} |\log t|^{n-1} dt + \int_1^\infty e^{-t} t^{x-1} t^{n-1} dt \\ &= \frac{\Gamma(n)}{x^n} + \Gamma(x+n-1). \end{aligned}$$

5.1.12. If $x_n \searrow 1$, then, $k^{-x_n} \nearrow k^{-1}$ for $k \geq 1$. So, by the monotone convergence theorem for series, $\zeta(x_n) = \sum_{k=1}^\infty k^{-x_n} \rightarrow \sum_{k=1}^\infty k^{-1} = \zeta(1) = \infty$. If $x_n \rightarrow 1+$, then, $x_n^* \searrow 1$ (§1.5) and $\zeta(1) \geq \zeta(x_n) \geq \zeta(x_n^*)$. So, $\zeta(x_n) \rightarrow \zeta(1) = \infty$. Thus, $\zeta(1+) = \infty$. Similarly, $\psi(0+) = \infty$.

5.1.13. Since $\tau(t) = \sum_{n=0}^\infty t e^{-nt}$, use summation under the integral sign:

$$\int_0^\infty e^{-xt} \tau(t) dt = \sum_{n=0}^\infty \int_0^\infty e^{-xt} t e^{-nt} dt = \sum_{n=0}^\infty \frac{1}{(x+n)^2}.$$

Here, we used the substitution $s = (x+n)t$, and $\Gamma(2) = 1$.

5.1.14. The problem, here, is that the limits of integration depend on n . So, the monotone convergence theorem is not directly applicable. To remedy this, let $f_n(x) = f(x)$ if $a_n < x < b_n$, and let $f_n(x) = 0$ if $a < x \leq a_n$ or $b_n \leq x < b$. Then, $f_n(x) \nearrow f(x)$ (draw a picture). Hence, by the monotone convergence theorem,

$$\int_{a_n}^{b_n} f(x) dx = \int_a^b f_n(x) dx \rightarrow \int_a^b f(x) dx.$$

Exercises 5.2

5.2.1. First, for $x > 0$,

$$e^{-sx} \left(1 + \frac{x^2}{3!} + \frac{x^4}{5!} + \dots \right) \leq e^{-sx} \sum_{n=0}^\infty \frac{x^{2n}}{(2n)!} \leq e^{-sx} e^x.$$

So, with $g(x) = e^{-(s-1)x}$, we may use summation under the integral sign to get

$$\begin{aligned} \int_0^\infty e^{-sx} \frac{\sin x}{x} dx &= \int_0^\infty \sum_{n=0}^\infty (-1)^n \frac{x^{2n}}{(2n+1)!} e^{-sx} dx \\ &= \sum_{n=0}^\infty \frac{(-1)^n}{(2n+1)!} \int_0^\infty e^{-sx} x^{2n} dx \\ &= \sum_{n=0}^\infty \frac{(-1)^n}{(2n+1)!} \cdot \frac{\Gamma(2n+1)}{s^{2n+1}} \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{(-1)^n (1/s)^{2n+1}}{(2n+1)} \\
&= \arctan\left(\frac{1}{s}\right).
\end{aligned}$$

Here, we used (6.4) in §3.6.

5.2.2. Since $|f_n(x)| \leq g(x)$ for all $n \geq 1$, taking the limit yields $|f(x)| \leq g(x)$. Since g is integrable, so, is f .

5.2.3. $J_0(x)$ is a power series, hence, may be differentiated term by term. The calculation is made simpler by noting that $x[xJ'_0(x)]' = x^2 J''_0(x) + xJ'_0(x)$. Then,

$$xJ'_0(x) = \sum_{n=1}^{\infty} (-1)^n \frac{2n x^{2n}}{4^n (n!)^2},$$

and

$$\begin{aligned}
x^2 J''_0(x) + xJ'_0(x) &= x(xJ'_0(x))' \\
&= \sum_{n=1}^{\infty} (-1)^n \frac{4n^2 x^{2n}}{4^n (n!)^2} \\
&= \sum_{n=0}^{\infty} (-1)^{n+1} \frac{4(n+1)^2 x^{2n+2}}{4^{n+1} ((n+1)!)^2} \\
&= -x^2 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{4^n (n!)^2} \\
&= -x^2 J_0(x).
\end{aligned}$$

5.2.4. With $u = \sin^{n-1} x$ and $dv = \sin x dx$,

$$I_n = \int \sin^n x dx = -\cos x \sin^{n-1} x + (n-1) \int \sin^{n-2} x \cos^2 x dx.$$

Inserting $\cos^2 x = 1 - \sin^2 x$,

$$I_n = -\cos x \sin^{n-1} x + (n-1)(I_{n-2} - I_n).$$

Solving for I_n ,

$$I_n = -\frac{1}{n} \cos x \sin^{n-1} x + \frac{n-1}{n} I_{n-2}.$$

5.2.5. Using the double-angle formula,

$$\begin{aligned}
\sin x &= 2 \cos(x/2) \sin(x/2) \\
&= 4 \cos(x/2) \cos(x/4) \sin(x/4) \\
&= \dots = 2^n \cos(x/2) \cos(x/4) \dots \cos(x/2^n) \sin(x/2^n).
\end{aligned}$$

Now, let $n \nearrow \infty$, and use $2^n \sin(x/2^n) \rightarrow x$.

5.2.6. The integral on the left equals $n!$. So, the left side is $\sum_{n=0}^{\infty} (-1)^n$, which has no sum. The series on the right equals e^{-x} . So, the right side equals $\int_0^{\infty} e^{-2x} dx = 1/2$.

5.2.7. Now, for $x > 0$,

$$\frac{\sin(sx)}{e^x - 1} = \sum_{n=1}^{\infty} e^{-nx} \sin(sx),$$

and

$$\sum_{n=1}^{\infty} e^{-nx} |\sin(sx)| \leq \sum_{n=1}^{\infty} e^{-nx} |s|x = |s|x/(e^x - 1) = g(x),$$

which is integrable ($\int_0^{\infty} g(x) dx = |s|\Gamma(2)\zeta(2)$ by Exercise 8 of §5.1). Hence, we may use summation under the integral sign to obtain

$$\begin{aligned} \int_0^{\infty} \frac{\sin(sx)}{e^x - 1} dx &= \sum_{n=1}^{\infty} \int_0^{\infty} e^{-nx} \sin(sx) dx \\ &= \sum_{n=1}^{\infty} \frac{s}{n^2 + s^2}. \end{aligned}$$

Here, we used Exercise 7 of §4.4.

5.2.8. Writing $\sinh(sx) = (e^{sx} - e^{-sx})/2$ and breaking the integral into two pieces leads to infinities. So, we proceed, as in the previous exercise. For $x > 0$, use the mean value theorem to check

$$\left| \frac{\sinh x}{x} \right| \leq \cosh x \leq e^x.$$

So,

$$\frac{\sinh(sx)}{e^x - 1} = \sum_{n=1}^{\infty} e^{-nx} \sinh(sx),$$

and

$$\sum_{n=1}^{\infty} e^{-nx} |\sinh(sx)| \leq \sum_{n=1}^{\infty} |s|x e^{-nx} e^{|s|x} = g(x),$$

which is integrable when $|s| < 1$ ($\int_0^{\infty} g(x) dx = |s| \sum_{n=1}^{\infty} \Gamma(2)/(n - |s|)^2$). Using summation under the integral sign, we obtain

$$\begin{aligned} \int_0^{\infty} \frac{\sinh(sx)}{e^x - 1} dx &= \sum_{n=1}^{\infty} \int_0^{\infty} e^{-nx} \sinh(sx) dx \\ &= \sum_{n=1}^{\infty} \frac{1}{2} \int_0^{\infty} e^{-nx} (e^{sx} - e^{-sx}) dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{1}{n-s} - \frac{1}{n+s} \right) \\
&= \sum_{n=1}^{\infty} \frac{s}{n^2 + s^2}.
\end{aligned}$$

5.2.9. Because (p_n) is nonzero only when $n-1, n-4, n-5$, or $n-6$ are divisible by 8,

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{p_n}{16^{[n/8]} n} &= \sum_{k=1}^8 \sum_{n=0}^{\infty} \frac{p_{8n+k}}{16^n (8n+k)} \\
&= \sum_{n=0}^{\infty} \frac{4}{(8n+1)16^n} + \sum_{n=0}^{\infty} \frac{-2}{(8n+4)16^n} \\
&\quad + \sum_{n=0}^{\infty} \frac{-1}{(8n+5)16^n} + \sum_{n=0}^{\infty} \frac{-1}{(8n+6)16^n} \\
&= \sum_{n=0}^{\infty} \frac{1}{16^n} \left(\frac{4}{8n+1} - \frac{2}{8n+4} - \frac{1}{8n+5} - \frac{1}{8n+6} \right).
\end{aligned}$$

5.2.10. We have to show that $x_n \rightarrow x$ implies $J_v(x_n) \rightarrow J_v(x)$. But $g(t) = 1$ is integrable over $(0, \pi)$ and dominates the integrands below. So, we can apply the dominated convergence theorem,

$$J_v(x_n) = \frac{1}{\pi} \int_0^{\pi} \cos(vt - x_n \sin t) dt \rightarrow \frac{1}{\pi} \int_0^{\pi} \cos(vt - x \sin t) dt = J_v(x).$$

5.2.11. It is enough to show that ψ is continuous on (a, ∞) for all $a > 0$, for, then, ψ is continuous on $(0, \infty)$. We have to show that $x_n \rightarrow x > a$ implies $\psi(x_n) \rightarrow \psi(x)$. But $x_n > a, n \geq 1$, implies $e^{-k^2 \pi x_n} \leq e^{-k^2 \pi a}, k \geq 1, n \geq 1$, and $\sum g_k = \sum e^{-k^2 \pi a} < \infty$. So, the dominated convergence theorem for series applies, and

$$\psi(x_n) = \sum_{k=1}^{\infty} e^{-k^2 \pi x_n} \rightarrow \sum_{k=1}^{\infty} e^{-k^2 \pi x} = \psi(x).$$

5.2.12. Set $\tilde{f}_n(x) = f_n(x)$ if $a_n < x < b_n$, and $\tilde{f}_n(x) = 0$ if $a < x < a_n \leq a_n$ or $b_n \leq x < b$. Then, $|\tilde{f}_n(x)| \leq g(x)$ on (a, b) , and $\tilde{f}_n(x) \rightarrow f(x)$ for any x in (a, b) . Hence, by the dominated convergence theorem,

$$\int_{a_n}^{b_n} f_n(x) dx = \int_a^b \tilde{f}_n(x) dx \rightarrow \int_a^b f(x) dx.$$

5.2.13. Use the Taylor series for \cos :

$$J_0(x) = \frac{1}{\pi} \int_0^{\pi} \cos(x \sin t) dt = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)! \pi} \int_0^{\pi} \sin^{2n} t dt.$$

But

$$\begin{aligned}\frac{1}{\pi} \int_0^\pi \sin^{2n} t \, dt &= \frac{2}{\pi} \int_0^{\pi/2} \sin^{2n} t \, dt = \frac{2}{\pi} I_{2n} \\ &= \frac{(2n-1) \cdot (2n-3) \cdot \dots \cdot 1}{2n \cdot (2n-2) \cdot \dots \cdot 2} = \frac{(2n)!}{2^{2n}(n!)^2}.\end{aligned}$$

Inserting this in the previous expression, one obtains the series for $J_0(x)$. Here, for x fixed, we used summation under the integral sign with $g(t) = e^x$. Since $\int_0^\pi g(t) \, dt = e^x \pi$, this applies.

5.2.14. By Exercise 18 of §4.4,

$$\begin{aligned}\lim_{n \nearrow \infty} \left[\int_0^1 \frac{1 - (1 - t/n)^n}{t} \, dt - \int_1^n \frac{(1 - t/n)^n}{t} \, dt \right] \\ &= \lim_{n \nearrow \infty} \left[\int_0^n \frac{1 - (1 - t/n)^n}{t} \, dt - \int_1^n \frac{1}{t} \, dt \right] \\ &= \lim_{n \nearrow \infty} \left(1 + \frac{1}{2} + \dots + \frac{1}{n} - \log n \right) \\ &= \gamma.\end{aligned}$$

For the second part, since $(1 - t/n)^n \rightarrow e^{-t}$, we obtain the stated formula by switching the limits and the integrals. To justify the switching, by the mean value theorem with $f(t) = (1 - t/n)^n$,

$$0 \leq \frac{1 - (1 - t/n)^n}{t} = \frac{f(0) - f(t)}{t} = -f'(c) = (1 - c/n)^{n-1} \leq 1.$$

So, we may choose $g(t) = 1$ for the first integral. Since $(1 - t/n)^n \leq e^{-t}$, we may choose $g(t) = e^{-t}/t$ for the second integral.

Exercises 5.3

5.3.1. If $a_n \rightarrow a$ and $b_n \rightarrow b$ with $a > b > 0$, then, there is a $c > 0$ with $a_n > c$ and $b_n > c$ for all $n \geq 1$. Hence,

$$f_n(\theta) = \frac{1}{\sqrt{a_n^2 \cos^2 \theta + b_n^2 \sin^2 \theta}} \leq \frac{1}{\sqrt{c^2 \cos^2 \theta + c^2 \sin^2 \theta}} = \frac{1}{c}.$$

Hence, we may apply the dominated convergence theorem with $g(\theta) = 2/c\pi$.

5.3.2. Since the arithmetic and geometric means of $(1+x, 1-x)$ are $(1, \sqrt{1-x^2})$, $M(1+x, 1-x) = M(1, \sqrt{1-x^2})$. So, the result follows from

$$\cos^2 \theta + (1 - x^2) \sin^2 \theta = 1 - x^2 \sin^2 \theta.$$

5.3.3. By the binomial theorem,

$$\frac{1}{\sqrt{1 - x^2 \sin^2 \theta}} = \sum_{n=0}^{\infty} (-1)^n \binom{-1/2}{n} x^{2n} \sin^{2n} \theta.$$

By Exercise 13 of §3.4, $\binom{-1/2}{n} = (-1)^n 4^{-n} \binom{2n}{n}$. So, this series is positive. Hence, we may apply summation under the integral sign. From Exercise 13 of §5.2, $I_{2n} = (2/\pi) \int_0^{\pi/2} \sin^{2n} \theta d\theta = 4^{-n} \binom{2n}{n}$. Integrating the series term by term, we get the result.

5.3.4. With $t = x/s$, $dt = -xds/s^2$, and $f(t) = 1/\sqrt{(1+t^2)(x^2+t^2)}$, $f(t) dt = -f(s) ds$. So,

$$\begin{aligned} \frac{1}{M(1, x)} &= \frac{2}{\pi} \int_0^\infty \frac{dt}{\sqrt{(1+t^2)(x^2+t^2)}} \\ &= \frac{2}{\pi} \int_0^{\sqrt{x}} f(t) dt + \frac{2}{\pi} \int_{\sqrt{x}}^\infty f(t) dt \\ &= \frac{2}{\pi} \int_0^{\sqrt{x}} f(t) dt + \frac{2}{\pi} \int_0^{\sqrt{x}} f(s) ds \\ &= \frac{4}{\pi} \int_0^{\sqrt{x}} \frac{dt}{\sqrt{(1+t^2)(x^2+t^2)}} \\ &= \frac{4}{\pi} \int_0^{1/\sqrt{x}} \frac{dr}{\sqrt{(1+(xr)^2)(1+r^2)}} \end{aligned}$$

For the last integral, we used $t = xr$, $dt = xdr$.

5.3.5. The AGM iteration yields $(1+x, 1-x) \mapsto (1, x') \mapsto ((1+x')/2, \sqrt{x'})$.

5.3.6. Now, $x < M(1, x) < 1$, and $x' < M(1, x') < 1$. So,

$$\begin{aligned} \left| \frac{1}{M(1, x)} - \frac{1}{Q(x)} \right| &= \frac{1 - M(1, x')}{M(1, x)} \\ &\leq \frac{1 - x'}{x} = \frac{1 - x'^2}{x(1 + x')} = \frac{x}{1 + x'} \leq x. \end{aligned}$$

5.3.7. We already know that

$$Q\left(\frac{1-x'}{1+x'}\right) = \frac{1}{2}Q(x).$$

Substitute $x = 2\sqrt{y}/(1+y)$. Then, $x' = (1-y)/(1+y)$. So, solving for y yields $y = (1-x')/(1+x')$.

5.3.8. From the integral formula, $M(1, x)$ is strictly increasing and continuous. So $M(1, x')$ is strictly decreasing and continuous. So, $Q(x)$ is strictly increasing and continuous. Moreover, $x \rightarrow 0$ implies $x' \rightarrow 1$ implies $M(1, x) \rightarrow 0$ and $M(1, x') \rightarrow 1$, which implies $Q(x) \rightarrow 0$. Thus, $Q(0+) = 0$. If $x \rightarrow 1-$, then, $x' \rightarrow 0+$. Hence, $M(1, x) \rightarrow 1$ and $M(1, x') \rightarrow 0+$. So, $Q(x) \rightarrow \infty$. Thus, $Q(1-) = \infty$, hence, $M(1, \cdot) : (0, 1) \rightarrow (0, 1)$ and $Q : (0, 1) \rightarrow (0, \infty)$ are strictly increasing bijections.

5.3.9. $M(a, b) = 1$ is equivalent to $M(1, b/a) = 1/a$ which is uniquely solvable for b/a , hence, for b by the previous exercise.

5.3.10. Let $x = b/a = f(a)/a$. Then, the stated asymptotic equality is equivalent to

$$\lim_{a \rightarrow \infty} \left[\log(x/4) + \frac{\pi a}{2} \right] = 0.$$

Since $0 < b < 1$, $a \rightarrow \infty$ implies $x \rightarrow 0$ and $M(1, x) = M(1, b/a) = 1/a$ by homogeneity, this follows from (3.8).

Exercises 5.4

5.4.1. If $x = \sqrt{2t}$, $dx = dt/\sqrt{2t}$, and $t = x^2/2$. So

$$\sqrt{\frac{\pi}{2}} = \int_0^\infty e^{-x^2/2} dx = \int_0^\infty e^{-t} \frac{dt}{\sqrt{2t}} = \frac{1}{\sqrt{2}} \Gamma(1/2).$$

Hence, $(1/2)! = \Gamma(3/2) = (1/2)\Gamma(1/2) = \sqrt{\pi}/2$.

5.4.2. Since $(x - s)^2 = x^2 - 2xs + s^2$,

$$e^{-s^2/2} L(s) = \int_{-\infty}^\infty e^{-(x-s)^2/2} dx = \int_{-\infty}^\infty e^{-x^2/2} dx = \sqrt{2\pi}$$

by translation invariance.

5.4.3. By differentiation under the integral sign,

$$L^{(n)}(s) = \int_{-\infty}^\infty e^{sx} x^n e^{-x^2/2} dx.$$

So,

$$L^{(2n)}(0) = \int_{-\infty}^\infty x^{2n} e^{-x^2/2} dx.$$

To justify this, note that for $|s| < b$ and $f(s, x) = e^{sx - x^2/2}$,

$$\begin{aligned} \sum_{k=0}^n \left| \frac{\partial^k}{\partial s^k} f(s, x) \right| &= e^{sx - x^2/2} \sum_{k=0}^n |x|^k \\ &\leq n! e^{sx - x^2/2} \sum_{k=0}^n \frac{|x|^k}{k!} \\ &\leq n! e^{(b+1)|x| - x^2/2} = g(x) \end{aligned}$$

and g is even and integrable ($\int_{-\infty}^\infty g(x) dx \leq 2n! L(b+1)$). Since the integrand is odd for n odd, $L^{(n)}(0) = 0$ for n odd. On the other hand, the exponential series yields

$$L(s) = \sqrt{2\pi} e^{s^2/2} = \sqrt{2\pi} \sum_{n=0}^\infty \frac{s^{2n}}{2^n n!} = \sum_{n=0}^\infty L^{(2n)}(0) \frac{s^{2n}}{(2n)!}.$$

Solving for $L^{(2n)}(0)$, we obtain the result.

5.4.4. With $f(s, x) = e^{-x^2/2} \cos(sx)$,

$$\begin{aligned} |f(s, x)| + \left| \frac{\partial}{\partial s} f(s, x) \right| &= e^{-x^2/2} (|\cos(sx)| + |x| |\sin(sx)|) \\ &\leq e^{-x^2/2} (1 + |x|) = g(x), \end{aligned}$$

which is integrable since $\int_{-\infty}^{\infty} g(x) dx = \sqrt{2\pi} + 2$. Thus, with $u = \sin(sx)$ and $dv = -xe^{-x^2/2} dx$, $v = e^{-x^2/2}$, $du = s \cos(sx) dx$. So,

$$\begin{aligned} F'(s) &= - \int_{-\infty}^{\infty} e^{-x^2/2} x \sin(sx) dx \\ &= uv|_{-\infty}^{\infty} - s \int_{-\infty}^{\infty} e^{-x^2/2} \cos(sx) dx \\ &= -sF(s). \end{aligned}$$

Integrating $F'(s)/F(s) = -s$ over $(0, s)$ yields $\log F(s) = -s^2/2 + \log F(0)$ or $F(s) = F(0)e^{-s^2/2}$.

5.4.5. With $f(a, x) = e^{-x-a/x}/\sqrt{x}$ and $a \geq \epsilon > 0$,

$$|f(a, x)| + \left| \frac{\partial}{\partial a} f(a, x) \right| \leq \begin{cases} e^{-\epsilon/x} \left(\frac{1}{\sqrt{x}} + \frac{1}{x\sqrt{x}} \right), & 0 < x < 1, \\ e^{-x}, & x > 1. \end{cases}$$

But the expression on the right is integrable over $(0, \infty)$. Hence, we may differentiate under the integral sign on (ϵ, ∞) , hence, on $(0, \infty)$. Thus, with $x = a/t$,

$$\begin{aligned} H'(a) &= - \int_0^{\infty} e^{-x-a/x} \frac{dx}{x\sqrt{x}} \\ &= - \int_0^{\infty} e^{-a/t-t} \frac{dt}{\sqrt{at}} = -\frac{1}{\sqrt{a}} H(a). \end{aligned}$$

Integrating $H'(a)/H(a) = -1/\sqrt{a}$, we get $\log H(a) = -2\sqrt{a} + \log H(0)$ or $H(a) = H(0)e^{-2\sqrt{a}}$.

5.4.6. Let $x = y\sqrt{q}$. Then, $dx = \sqrt{q} dy$. Hence,

$$\int_{-\infty}^{\infty} e^{-x^2/2q} dx = \sqrt{q} \int_{-\infty}^{\infty} e^{-y^2/2} dy = \sqrt{2\pi q}.$$

5.4.7. Inserting $g(x) = e^{-x^2\pi}$ and $\delta = \sqrt{t}$ in Exercise 8 of §4.3 and $\pi = 1/2q$ in the previous Exercise yields

$$\lim_{t \rightarrow 0+} \sqrt{t} \psi(t) = \int_0^{\infty} e^{-x^2\pi} dx = \frac{1}{2} \sqrt{2\pi q} = \frac{1}{2}.$$

5.4.8. It is enough to show that ζ is smooth on (a, ∞) for any $a > 1$. Use differentiation N times under the summation sign to get

$$\zeta^{(k)}(s) = \sum_{n=1}^{\infty} \frac{(-1)^k \log^k n}{n^s}, \quad s > a, N \geq k \geq 0.$$

To justify this, let $f_n(s) = n^{-s}$, $n \geq 1$, $s > 1$. Since $\log n/n^\epsilon \rightarrow 0$ as $n \nearrow \infty$, for any $\epsilon > 0$, the sequence $(\log n/n^\epsilon)$ is bounded, which means that there is a constant $C_\epsilon > 0$, such that $|\log n| \leq C_\epsilon n^\epsilon$ for all $n \geq 1$. Hence,

$$\sum_{k=0}^N |f_n^{(k)}(s)| \leq \sum_{k=0}^N \frac{|\log^k n|}{n^s} \leq (N+1) \frac{C_\epsilon^{N+1} n^{N\epsilon}}{n^a} = \frac{C}{n^{a-N\epsilon}}.$$

Then, if we choose ϵ small enough, so that $a - N\epsilon > 1$, the dominating series $\sum g_n = C \sum n^{N\epsilon-a}$ converges.

5.4.9. Again, we show that ψ is smooth on (a, ∞) for all $a > 0$. Use differentiation N times under the summation sign to get

$$\psi^{(k)}(t) = \sum_{n=1}^{\infty} (-1)^k \pi^k n^{2k} e^{-n^2 \pi t}, \quad t > a, N \geq k \geq 0.$$

To justify this, let $f_n(t) = e^{-n^2 \pi t}$, $n \geq 1$, $t > a$. Since $x^{N+1} e^{-x} \rightarrow 0$ as $x \rightarrow \infty$, the function $x^{N+1} e^{-x}$ is bounded. Thus, there is a constant $C_N > 0$, such that $x^N e^{-x} \leq C_N/x$ for $x > 0$. Inserting $x = n^2 \pi t$, $t > a$,

$$\sum_{k=0}^N |f_n^{(k)}(t)| \leq \sum_{k=0}^N n^{2k} \pi^k e^{-n^2 \pi t} \leq \frac{(N+1)C_N}{n^2 \pi \cdot a^{N+1}}.$$

Then, the dominating series $\sum g_n = \sum (N+1)C_N/\pi a^{N+1} n^2$ converges.

5.4.10. Differentiating under the integral sign leads only to bounded functions of t . So J_ν is smooth. Computing, we get

$$J'_\nu(x) = \frac{1}{\pi} \int_0^\pi \sin t \sin(\nu t - x \sin t) dt$$

and

$$J''_\nu(x) = -\frac{1}{\pi} \int_0^\pi \sin^2 t \cos(\nu t - x \sin t) dt.$$

Now, integrate by parts with $u = -x \cos t - \nu$, $dv = (\nu - x \cos t) \cos(\nu t - x \sin t) dt$, $du = x \sin t dt$, and $v = \sin(\nu t - x \sin t)$:

$$\begin{aligned} x^2 J''_\nu(x) + (x^2 - \nu^2) J_\nu(x) &= \frac{1}{\pi} \int_0^\pi (x^2 \cos^2 t - \nu^2) \cos(\nu t - x \sin t) dt \\ &= \frac{1}{\pi} \int_0^\pi u dv = \frac{1}{\pi} uv \Big|_0^\pi - \frac{1}{\pi} \int_0^\pi v du \\ &= -\frac{1}{\pi} \int_0^\pi x \sin t \sin(\nu t - x \sin t) dt \\ &= -x J'_\nu(x). \end{aligned}$$

Here, ν must be an integer to make the uv term vanish at π .

5.4.11. Differentiating under the integral sign,

$$F^{(n)}(s) = \int_{-\infty}^{\infty} x^n e^{sx} e^{-f(x)} dx.$$

Since $|x|^n \leq n!e^{|x|}$, with $h(s, x) = e^{sx} e^{-f(x)}$ and $|s| < b$,

$$\sum_{n=0}^N \left| \frac{\partial^n h}{\partial s^n} \right| \leq \sum_{n=0}^N |x|^n e^{s|x|} e^{-f(x)} \leq (N+1)! e^{(b+1)|x|} e^{-f(x)} = g(x),$$

and g is integrable by Exercise 11 of §4.3. This shows that F is smooth. Differentiating twice,

$$[\log F(s)]'' = \frac{F''(s)F(s) - F'(s)^2}{F(s)^2}.$$

Now, use the Cauchy–Schwarz inequality (Exercise 17 of §4.4) with the functions $e^{(sx-f(x))/2}$ and $xe^{(sx-f(x))/2}$ to get $F''(s)F(s) \geq F'(s)^2$. Hence, $[\log F(s)]'' \geq 0$, or $\log F(s)$ is convex.

5.4.12. With $u = e^{-sx}$ and $dv = \sin x(dx/x)$, $du = -se^{-sx} dx$, and $v = F(t) = \int_0^t \sin r(dr/r)$. So, integration by parts yields the first equation. Now, change variables $y = sx$, $sdx = dy$ in the integral on the right yielding

$$\int_0^b e^{-sx} \frac{\sin x}{x} dx = -e^{-sb} F(b) + \int_0^{b/s} e^{-y} F(y/s) dy.$$

Let $b \rightarrow \infty$. Since F is bounded,

$$\int_0^{\infty} e^{-sx} \frac{\sin x}{x} dx = \int_0^{\infty} e^{-y} F(y/s) dy.$$

Now, let $s \rightarrow 0+$, and use the dominated convergence theorem. Since $F(y/s) \rightarrow F(\infty)$, as $s \rightarrow 0+$, for all $y > 0$,

$$\lim_{s \rightarrow 0+} \int_0^{\infty} e^{-sx} \frac{\sin x}{x} dx = \int_0^{\infty} e^{-y} F(\infty) dy = F(\infty) = \lim_{b \rightarrow \infty} \int_0^b \frac{\sin x}{x} dx.$$

But, from the text, the left side is

$$\lim_{x \rightarrow 0+} \arctan\left(\frac{1}{x}\right) = \arctan(\infty) = \frac{\pi}{2}.$$

Exercises 5.5

5.5.1. Without loss of generality, assume that $a = \max(a, b, c)$. Then, $(b/a)^n \leq 1$, and $(c/a)^n \leq 1$. So,

$$\lim_{n \nearrow \infty} (a^n + b^n + c^n)^{1/n} = a \lim_{n \nearrow \infty} (1 + (b/a)^n + (c/a)^n)^{1/n} = a.$$

For the second part, replace a, b, c in the first part by e^a, e^b , and e^c . Then, take the log.

5.5.2. The relative error is about .08%.

5.5.3. In the asymptotic for $\binom{n}{k}$, replace n by $2n$ and k by n . Since $t = n/2n = 1/2$ and $H(1/2) = 0$, we get $1/\sqrt{\pi n}$.

5.5.4. Straight computation.

5.5.5. $H'(t, p) = \log(t/p) - \log[(1-t)/(1-p)]$ equals zero when $t = p$. Since $H''(t, p) = 1/t + 1/(1-t)$, H is convex. So, $t = p$ is a global minimum.

5.5.6. Straight computation.

5.5.7. Since $(q^n)^{x^2} = e^{nx^2 \log q}$, the limit is

$$\sup\{x^2 \log q : a < x < b\} = a^2 \log q,$$

by the theorem. Here, $\log q < 0$.

5.5.8. Since $\Gamma(s+1) = s\Gamma(s)$,

$$\begin{aligned} f(s+1) &= 3^{3s+3} \frac{\Gamma(s+1)\Gamma(s+1+1/3)\Gamma(s+1+2/3)}{\Gamma(3s+3)} \\ &= \frac{3^3 s(s+1/3)(s+2/3)}{(3s+2)(3s+1)3s} \cdot f(s) = f(s). \end{aligned}$$

Inserting the asymptotic for $\Gamma(s+n)$ yields $2\pi\sqrt{3}$ for the limit. The general case is given by

$$n^{ns} \frac{\Gamma(s)\Gamma(s+1/n)\dots\Gamma(s+(n-1)/n)}{\Gamma(ns)} = \sqrt{n}(2\pi)^{(n-1)/2}.$$

5.5.9. Here,

$$L_n(ns) = \int_{-\infty}^{\infty} e^{n(sx-f(x))} dx,$$

and $\max\{sx - f(x) : x \in \mathbf{R}\}$ exists by superlinearity and is attained at some c . Write $L_n(ns) = I_n^- + I_n^0 + I_n^+ = \int_{-\infty}^a + \int_a^b + \int_b^{\infty}$. Let M be a positive number, to be chosen below. Then, by superlinearity, we may choose $a < c$ sufficiently negative, such that $f(x) > -Mx$ for $x < a$. Similarly, we may choose $b > c$ sufficiently positive, such that $f(x) > Mx$ for $x > b$. The first theorem in the text applies to I_n^0 , whereas

$$I_n^- \leq \int_{-\infty}^a e^{n(M+s)x} dx = \frac{e^{n(M+s)a}}{n(M+s)}, \quad M+s > 0,$$

and

$$I_n^+ \leq \int_b^{\infty} e^{-n(M-s)x} dx = \frac{e^{-n(M-s)b}}{n(M-s)}, \quad M-s > 0.$$

Thus,

$$\lim_{n \nearrow \infty} (1/n) \log(I_n^-) \leq (M+s)a,$$

$$\lim_{n \nearrow \infty} (1/n) \log(I_n^+) \leq -(M - s)b.$$

Choosing M large enough (remember a is negative and b is positive), by the second part of Exercise 1, we conclude that

$$\begin{aligned} \lim_{n \nearrow \infty} \frac{1}{n} \log L_n(ns) &= \lim_{n \nearrow \infty} \frac{1}{n} \log (I_n^- + I_n^0 + I_n^+) \\ &= \max \left[(M + s)a, \max\{sx - f(x) : a < x < b\}, (-M + s)b \right] \\ &= \max\{sx - f(x) : x \in \mathbb{R}\}. \end{aligned}$$

5.5.10. The log of the duplication formula is

$$2s \log 2 + \log \Gamma(s) + \log \Gamma(s + 1/2) - \log \Gamma(2s) = \log(2\sqrt{\pi}).$$

Differentiating,

$$2 \log 2 + \frac{\Gamma'(s)}{\Gamma(s)} + \frac{\Gamma'(s + 1/2)}{\Gamma(s + 1/2)} - 2 \frac{\Gamma'(2s)}{\Gamma(2s)} = 0.$$

Inserting $s = 1/2$, we obtain the result.

5.5.11. Insert $s = 1/4$ in the duplication formula to get

$$\sqrt{2} \frac{\Gamma(1/4)\Gamma(3/4)}{\Gamma(1/2)} = 2\sqrt{\pi}.$$

Now recall that $\Gamma(1/2) = \sqrt{\pi}$. To obtain the formula for $1/M(1, 1/\sqrt{2})$, replace $\Gamma(3/4)$ in the formula in the text by $\pi\sqrt{2}/\Gamma(1/4)$.

Exercises 5.6

5.6.1. $\zeta(4) = \pi^4/90$, $\zeta(6) = \pi^6/945$, and $\zeta(8) = \pi^8/9450$.

5.6.2. Let $b_k = B_k/k!$, and suppose that $|b_k| \leq 2^k$ for $k \leq n - 2$. Then, (6.7) reads

$$\sum_{k=0}^{n-1} \frac{b_k (-1)^{n-1-k}}{(n-k)!} = 0$$

which implies $(n! \geq 2^{n-1})$

$$\begin{aligned} |b_{n-1}| &\leq \sum_{k=0}^{n-2} \frac{|b_k|}{(n-k)!} \leq \sum_{k=0}^{n-2} \frac{2^k}{(n-k)!} \\ &\leq \sum_{k=0}^{n-2} \frac{2^k}{2^{n-k-1}} \leq 2^{n-1}. \end{aligned}$$

Thus, $|b_n| \leq 2^n$ for all $n \geq 1$ by induction. Hence, the radius of convergence by the root test is at least $1/2$. Also, from the formula for $\zeta(2n) > 0$, the Bernoulli numbers are alternating.

5.6.3. The left inequality follows from (6.9) since $b_1 = \sum_{n=1}^{\infty} a_n$. For the right inequality, use $1 + a_n \leq e^{a_n}$. So,

$$\prod_{n=1}^{\infty} (1 + a_n) \leq \prod_{n=1}^{\infty} e^{a_n} = \exp \left(\sum_{n=1}^{\infty} a_n \right).$$

5.6.4. From (1.3),

$$\begin{aligned} \Gamma(x) &= \lim_{n \nearrow \infty} \frac{n^x n!}{x(1+x)(2+x) \dots (n+x)} \\ &= \frac{1}{x} \lim_{n \nearrow \infty} \frac{e^{x(\log n - 1 - 1/2 - \dots - 1/n)} e^x e^{x/2} \dots e^{x/n}}{(1+x)(1+x/2) \dots (1+x/n)} \\ &= \frac{e^{-\gamma x}}{x} \prod_{n=1}^{\infty} \left(\frac{e^{x/n}}{1 + \frac{x}{n}} \right). \end{aligned}$$

5.6.5. For $0 < x < 1$, use (1.5) with x and $1-x$ replacing x . Then, $\Gamma(x)\Gamma(1-x)$ equals

$$\begin{aligned} &\lim_{n \nearrow \infty} \frac{n^x n! n^{1-x} n!}{x(1+x)(2+x) \dots (n+x)(1-x)(2-x) \dots (n+1-x)} \\ &= \frac{1}{x} \lim_{n \nearrow \infty} \frac{n(n!)^2}{(1-x^2)(4-x^2) \dots (n^2-x^2)(n+1-x)} \\ &= \frac{1}{x} \lim_{n \nearrow \infty} \frac{1}{(1-x^2)(1-x^2/4) \dots (1-x^2/n^2)(1+(1-x)/n)} \\ &= \left[x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2} \right) \right]^{-1} = \frac{\pi}{\sin(\pi x)}. \end{aligned}$$

5.6.6. The series $B(x)$ is the alternating version of the Bernoulli series (6.6), and the Taylor series for $\sin(x/2)$ is the alternating version of the Taylor series for $\sinh(x/2)$. But the Bernoulli series times $\sinh(x/2)$ equals $(x/2) \cosh(x/2)$. Hence (Exercise 8 of §1.7),

$$B(x) \sin(x/2) = (x/2) \cos(x/2).$$

Dividing by $\sin(x/2)$, we obtain the series for $(x/2) \cot(x/2)$.

5.6.7. If $\beta > 1$, then, $B(x)$ would converge at 2π . But $(x/2) \cot(x/2) = B(x)$ is infinite at 2π .

5.6.8. Taking the log of (6.12),

$$\log[\sin(\pi x)] - \log(\pi x) = \sum_{n=1}^{\infty} \log \left(1 - \frac{x^2}{n^2} \right).$$

Differentiating under the summation sign,

$$\pi \cot(\pi x) - \frac{1}{x} = \sum_{n=1}^{\infty} \frac{2x}{x^2 - n^2}.$$

To justify this, let $f_n(x) = \log(1 - x^2/n^2)$ and let $|x| < b < 1$. Then, $\log(1 - t) = t + t^2/2 + t^3/3 + \dots \leq t + t^2 + t^3 + \dots = t/(1 - t)$. So,

$$|f_n(x)| + |f'_n(x)| \leq \frac{x^2}{n^2 - x^2} + \frac{2|x|}{n^2 - x^2} \leq \frac{b^2 + 2b}{n^2 - b^2} = g_n,$$

which is summable. Since this is true for all $b < 1$, the equality is valid for $|x| < 1$.

5.6.9. By Exercise 13 of §3.5, $\cot x - 2 \cot(2x) = \tan x$. Then, the series for $\tan x$ follows from the series for $\cot x$ in Exercise 6 applied to $\cot x$ and $2 \cot(2x)$.

5.6.10. By Exercise 4,

$$\log \Gamma(x) = -\gamma x - \log x + \sum_{n=1}^{\infty} \left[\frac{x}{n} - \log \left(1 + \frac{x}{n} \right) \right].$$

Differentiating, we obtain the result. To justify this, let $f_n(x) = x/n - \log(1 + x/n)$, $f'_n(x) = 1/n - 1/(x+n)$. Then, $t - \log(1 + t) = t^2/2 - t^3/3 + \dots \leq t^2/2$ for $t > 0$. Hence, $f_n(x) \leq x^2/n^2$. So,

$$|f_n(x)| + |f'_n(x)| \leq \frac{b^2 + b}{n^2} = g_n,$$

which is summable, when $0 < x < b$. Since b is arbitrary, the result is valid for $x > 0$.

5.6.11. Inserting $x = 1$ in the series for $\Gamma'(x)/\Gamma(x)$ yields a telescoping series. So, we get $\Gamma'(1) = -\gamma$. Inserting $x = 2$ yields

$$-\gamma - \frac{1}{2} + 1 - \frac{1}{3} + \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{5} + \frac{1}{4} - \frac{1}{6} + \dots = 1 - \gamma.$$

Since Γ is strictly convex, this forces the min to lie in $(1, 2)$.

5.6.12. Differentiate the series in Exercise 10, and compare with the series in Exercise 13 of §5.1. Here, on $0 < x < b$, we may take $g_n = (b+1)/n^2$, $n \geq 1$.

5.6.13. Substituting $(1-x)/2 \mapsto x$, we see that the stated equality is equivalent to

$$\lim_{x \rightarrow 0+} \left[\frac{\Gamma'(x)}{\Gamma(x)} + \frac{1}{x} \right] = -\gamma.$$

Move the $1/x$ to the left in Exercise 10, and, then, take the limit $x \rightarrow 0+$. Under this limit, the series collapses to zero by the dominated convergence theorem for series (here, $g_n = b/n^2$ for $0 < x < b$).

Exercises 5.7

5.7.1. θ_0 is strictly increasing since it is the sum of strictly increasing monomials. Since θ_0 is continuous and $\theta_0(0+) = 1$, $\theta_0(1-) = \infty$, $\theta_0((0, 1)) = (1, \infty)$. Similarly, for θ_+ .

5.7.2. Multiply (7.16) by $s\theta_0(s)^2 = \theta_0^2(1/s)$. You get (7.17).

5.7.3. The AGM of $\theta_0^2(q)$ and $\theta_-^2(q)$ equals 1: $M(\theta_0^2(q), \theta_-^2(q)) = 1$ for $0 < q < 1$. Since θ_0 is strictly increasing, θ_0^2 is strictly increasing. This forces θ_-^2 to be strictly decreasing. Moreover, $\theta_-(0+) = 1$. Hence, $\theta_-^2(0+) = 1$, and $q \rightarrow 1-$ implies $M(\infty, \theta_-^2(1-)) = 1$ or $\theta_-^2(1-) = 0$. Thus, θ_-^2 maps $(0, 1)$ onto $(0, 1)$. Since θ_- is continuous and $\theta_-(0+) = 1$, we also have θ_- strictly decreasing and $\theta_-((0, 1)) = (0, 1)$.

5.7.4. Since $\sigma(6) = \sigma(7) = 0$ and $\sigma(2n) = \sigma(n)$, $\sigma(12) = \sigma(14) = 0$. Also, $\sigma(13) = 8$ since

$$13 = (\pm 2)^2 + (\pm 3)^2 = (\pm 3)^2 + (\pm 2)^2.$$

To show that $\sigma(4n - 1) = 0$, we show that $4n - 1 = i^2 + j^2$ cannot happen. Note that $4n - 1$ is odd when exactly one of i or j is odd and the other is even. Say $i = 2k$ and $j = 2\ell + 1$. Then,

$$4n - 1 = 4k^2 + 4\ell^2 + 4\ell + 1 = 4(k^2 + \ell^2 + \ell) + 1,$$

an impossibility. Hence, $\sigma(4n - 1) = 0$, so, $\sigma(11) = \sigma(15) = 0$.

5.7.5. Let $f_n(t, x) = e^{-n^2\pi t} \cos(nx)$, $n \geq 1$. Then, $\partial f_n / \partial t = -n^2\pi f_n$, and $\partial^2 f_n / \partial x^2 = -n^2 f_n$, $n \geq 1$. Thus, to obtain the heat equation, we need only justify differentiation under the summation sign. But, for $t \geq 2a > 0$,

$$|f_n| + \left| \frac{\partial f_n}{\partial t} \right| + \left| \frac{\partial f_n}{\partial x} \right| + \left| \frac{\partial^2 f_n}{\partial x^2} \right| \leq 4n^2\pi e^{-2an^2} = g_n,$$

which is summable since xe^{-ax} is bounded for $x > 0$, hence, $xe^{-2ax} \leq e^{-ax}$.

Exercises 5.8

5.8.1. Assume that $1 < x < 2$. The integrand $f(x, t)$ is positive, hence, increasing, hence, $\leq f(2, t)$. By Exercise 15 of §3.4, $f(2, t)$ is asymptotically equal to $t/2$, as $t \rightarrow 0+$ which is bounded. Also, the integrand is asymptotically equal to te^{-t} , as $t \rightarrow \infty$ which is integrable. Since $f(2, t)$ is continuous, $g(t) = f(2, t)$ is integrable. Hence, we may switch the limit with the integral.

5.8.2. The limit of the left side of (8.4) is γ , by (8.3), and $\Gamma(1) = 1$.

5.8.3. Since $\Gamma(x + n + 1) = (x + n)\Gamma(x + n) = (x + n)(x + n - 1) \dots x\Gamma(x)$, for $x \in (-n - 1, -n) \cup (-n, -n + 1)$,

$$(x + n)\Gamma(x) = \frac{\Gamma(x + n + 1)}{x(x + 1) \dots (x + n - 1)}.$$

Letting $x \rightarrow -n$, we get $(x+n)\Gamma(x) \rightarrow (-1)^n/n!$.

5.8.4. For $t \geq 1$,

$$\psi(t) \leq \sum_{n=1}^{\infty} e^{-n\pi t} = \frac{e^{-\pi t}}{1 - e^{-\pi t}} \leq ce^{-\pi t}$$

with $c = 1/(1 - e^{-\pi})$. Now, the integrand $f(x, t)$ in (8.9) is a smooth function of x with $\partial^n f/\partial x^n$ continuous in (x, t) for all $n \geq 0$. Moreover, for $b > 1$ and $0 < x < b$ and each $n \geq 0$,

$$\begin{aligned} \left| \frac{\partial^n f}{\partial x^n} \right| &\leq \psi(t) 2^{-n} |\log t|^n [t^{(1-x)/2} + t^{x/2}] \\ &\leq ce^{-\pi t} 2^{-n+1} |\log t|^n t^{b/2} = g_n(t), \end{aligned}$$

which is integrable over $(1, \infty)$. Hence, we may repeatedly apply differentiation under the integral sign to conclude that the integral in (8.9) is smooth.

5.8.5. Inserting $x = 2n$ in (8.10), we get

$$\begin{aligned} \zeta(1-2n) &= \frac{\pi^{-n}\Gamma(n)\zeta(2n)}{\pi^{n-1/2}\Gamma(-n+1/2)} \\ &= \pi^{-2n+1/2}(n-1)! \\ &\quad \times \frac{(-n+1/2)(-n+3/2)\dots(-n+n-1/2)}{\Gamma(-n+n+1/2)} \\ &\quad \times \frac{(-1)^{n-1}B_{2n}2^{2n-1}\pi^{2n}}{(2n)!} = -\frac{B_{2n}}{2n}. \end{aligned}$$

5.8.6. Here, $f(x, t) = (1 + [t] - t)/t^{x+1}$, and $I(x) = \int_1^\infty f(x, t) dt$, $x > 1$. Since, for $b > x > a > 1$,

$$|f(x, t)| + \left| \frac{\partial f}{\partial x}(x, t) \right| \leq \frac{b+2}{t^{a+1}} = g(t), \quad t > 1,$$

and g is integrable, we may differentiate under the integral sign, obtaining $I'(x) = (x+1)I(x+1)$ for $a < x < b$. Since $a < b$ are arbitrary, this is valid for $x > 0$. Inserting $x = 2$ in (8.6) yields $\pi^2/6 = \zeta(2) = 1 + 2I(2)$ or $I(2) = \pi^2/12 - 1/2$.

5.8.7. The right side of (8.9) is smooth, except at $x = 0, 1$. Hence, $(x-1)\zeta(x)$ is smooth, except (possibly) at $x = 0$. By (8.6),

$$\log((x-1)\zeta(x)) = \log\left(1 + x(x-1)I(x)\right).$$

So,

$$\frac{\zeta'(x)}{\zeta(x)} + \frac{1}{x-1} = \frac{d}{dx} \log((x-1)\zeta(x)) = \frac{(2x-1)I(x) + x(x-1)I'(x)}{1 + x(x-1)I(x)}.$$

Taking the limit $x \rightarrow 1$, we approach $I(1) = \gamma$.

5.8.8. (8.10) says that $\pi^{-x/2}\Gamma(x/2)\zeta(x) = \pi^{-(1-x)/2}\Gamma((1-x)/2)\zeta(1-x)$. Differentiating the log of (8.10) yields

$$-\frac{1}{2}\log\pi + \frac{1}{2}\frac{\Gamma'(x/2)}{\Gamma(x/2)} + \frac{\zeta'(x)}{\zeta(x)} = \frac{1}{2}\log\pi - \frac{1}{2}\frac{\Gamma'((1-x)/2)}{\Gamma((1-x)/2)} - \frac{\zeta'(1-x)}{\zeta(1-x)}.$$

Now, add $1/(x-1)$ to both sides, and take the limit $x \rightarrow 1$. By the previous exercise, the left side becomes $-\log\pi/2 + \Gamma'(1/2)/2\Gamma(1/2) + \gamma$. By Exercise 13 of §5.6, the right side becomes $\log\pi/2 + \gamma/2 - \zeta'(0)/\zeta(0)$. But $\zeta(0) = -1/2$, and, by Exercise 10 of §5.5 and Exercise 11 of §5.6,

$$\begin{aligned}\frac{1}{2}\frac{\Gamma'(1/2)}{\Gamma(1/2)} &= \frac{1}{2}\frac{\Gamma'(1)}{\Gamma(1)} - \log 2 \\ &= -\frac{\gamma}{2} - \log 2.\end{aligned}$$

So,

$$-\frac{\log\pi}{2} + \left(-\frac{\gamma}{2} - \log 2\right) + \gamma = \frac{\log\pi}{2} + \gamma/2 + 2\zeta'(0).$$

Hence, $\zeta'(0) = -\log(2\pi)/2$.

5.8.9. For $0 < a \leq 1/2$,

$$-\log(1-a) = a + \frac{a^2}{2} + \frac{a^3}{3} + \dots \leq a + a^2 + a^3 + \dots = \frac{a}{1-a} \leq 2a.$$

On the other hand, by the triangle inequality, for $|a| \leq 1/2$,

$$\begin{aligned}|- \log(1-a) - a| &= \left| \frac{a^2}{2} + \frac{a^3}{3} + \dots \right| \\ &\leq \frac{|a|^2}{2} + \frac{|a|^3}{3} + \dots \\ &\leq \frac{1}{2}(|a|^2 + |a|^3 + \dots) = \frac{1}{2} \cdot \frac{a^2}{1-|a|} \leq a^2.\end{aligned}$$

5.8.10. m and n are both odd iff mn is odd. So, $\chi_+(m)$ and $\chi_+(n)$ both equal 1 iff $\chi_+(mn) = 1$. Since χ_+ equals 0 or 1, this shows that $\chi_+(mn) = \chi_+(m)\chi_+(n)$. For χ_- , m or n is even iff mn is even. So,

$$(5.1) \quad \chi_-(mn) = \chi_-(m)\chi_-(n)$$

when either n or m is even. If n and m are both odd and $m = 4i + 3$, $n = 4j + 3$, then, $mn = (4i + 3)(4j + 3) = 4(4ij + 3i + 3j + 2) + 1$ which derives (5.1) when $\chi_-(m) = \chi_-(n) = -1$. The other 3 cases are similar.

5.8.11. With $f_n(x) = 1/(4n-3)^x - 1/(4n-1)^x$,

$$L(s, \chi_-) = \sum_{n=1}^{\infty} f_n(x), \quad x > 0.$$

Then, by the mean value theorem, $f_n(x) \leq x/(4n-3)^{x+1}$. Hence, $|f_n(x)| \leq b/(4n-3)^{a+1} = g_n$ for $0 < a < x < b$. Since $4n-3 \geq n$, $\sum g_n \leq b\zeta(a+1)$. So, the dominated convergence theorem applies.

Exercises 5.9

5.9.1. On $n-1 \leq x \leq n$, $q(x) = \sum_{k=0}^n f(x-k)$, so, $q(n) = \sum_{k=0}^n f(n-k)$. On $n \leq x \leq n+1$, $q(x) = \sum_{k=0}^{n+1} f(x-k) = \sum_{k=0}^n f(x-k) + f(x-n-1)$, so, $q(n) = \sum_{k=0}^n f(n-k) + f(-1)$. Since $f(-1) = 0$, q is well defined at all integers. If $n-1 \leq x \leq n$, $q(x) = \sum_{k=0}^n f(x-n)$, and $q(x-1) = \sum_{k=0}^n f(x-1-k) = \sum_{k=1}^{n+1} f(x-k)$. So, $q(x) - q(x-1) = f(x) - f(x-n-1) = f(x)$ since $f = 0$ on $[-2, -1]$. Thus, q solves (9.3). To show that q is smooth on \mathbb{R} , assume that q is smooth on $(-\infty, n)$. Then, $q(x-1) + f(x)$ is smooth on $(-\infty, n+1)$. Hence, so is $q(x)$ by (9.3). Thus, q is smooth on \mathbb{R} .

5.9.2. If $f(x) = 0$ for $x > 1$, the formula reads

$$q(x) = \begin{cases} -f(x+1), & x \geq -1, \\ -f(x+1) - f(x+2), & -1 \geq x \geq -2, \\ -f(x+1) - f(x+2) - f(x+3), & -2 \geq x \geq -3, \\ \text{and so on.} \end{cases}$$

5.9.3. We show, by induction on $n \geq 0$, that

$$(5.2) \quad c(D)[e^{ax}x^n] = \frac{\partial^n}{\partial a^n} [c(a)e^{ax}]$$

for all $|a| < R$ and convergent series $c(a)$ on $(-R, R)$. Clearly, this is so for $n = 0$. Assume that (5.2) is true for $n-1$, and check (by induction over k) that $D^k(e^{ax}x^n) = xD^k(e^{ax}x^{n-1}) + kD^{k-1}(e^{ax}x^{n-1})$, $k \geq 0$. Taking linear combinations, we get $c(D)(e^{ax}x^n) = xc(D)(e^{ax}x^{n-1}) + c'(D)(e^{ax}x^{n-1})$. By the inductive hypothesis applied to c and c' , we obtain

$$\begin{aligned} c(D)(e^{ax}x^n) &= \frac{\partial^{n-1}}{\partial a^{n-1}} [xc(a)e^{ax} + c'(a)e^{ax}] \\ &= \frac{\partial^{n-1}}{\partial a^{n-1}} \cdot \frac{\partial}{\partial a} [c(a)e^{ax}] \\ &= \frac{\partial^n}{\partial a^n} [c(a)e^{ax}]. \end{aligned}$$

Thus, (5.2) is true for all $n \geq 0$.

5.9.4. Change variable $xt = s$, $xdt = ds$. Then, with $|f(t)| \leq C$, $0 < t < 1$,

$$\begin{aligned} \left| x^{n+1} \int_0^1 e^{-xt} f(t) t^n dt \right| &= \left| \int_0^x e^{-s} f(s/x) s^n ds \right| \\ &\leq C \int_0^\infty e^{-s} s^n ds = C\Gamma(n+1). \end{aligned}$$

This shows that the integral is $O(x^{-n-1})$.

5.9.5. Note that $\int_1^\infty e^{-xt} dt = e^{-x}/x$ and, by differentiation under the integral sign,

$$\int_1^\infty e^{-xt} t^p dt = (-1)^p \frac{d^p}{dx^p} \int_1^\infty e^{-xt} dt = (-1)^p \frac{d^p}{dx^p} \frac{e^{-x}}{x} = R(x)e^{-x}$$

for some rational function R . But $e^{-x} = O(x^{-n})$ for all $n \geq 1$ implies $R(x)e^{-x} = O(x^{-n})$ for all $n \geq 1$. Thus, the integral is ≈ 0 .

5.9.6. If the Stirling series converged at some point a , then, $B_n/n(n-1)a^{n-1}$, $n \geq 1$, would be bounded by the n th term test. Then, the Bernoulli series would be dominated by

$$\sum \left| \frac{B_n}{n!} x^n \right| \leq \sum \frac{C}{|a|(n-2)!} (|a|x)^n$$

which converges for all x . But we know (§5.6) that the Bernoulli series converges, at most, on $(-2\pi, 2\pi)$.

References

1. Richard Bellman, *Analytic Number Theory*, Benjamin, Reading, MA, 1980.
2. Garrett Birkhoff, *A Source Book in Classical Analysis*, Harvard University Press, Cambridge, MA, 1973.
3. J. M. Borwein and P. B. Borwein, *Pi and the AGM*, Wiley-Interscience, New York, NY, 1987.
4. D. Bailey, P. Borwein, and S. Plouffe, *On the Rapid Computation of Various Polylogarithmic Constants*, World-Wide Web, <http://www.cecm.sfu.ca/~pborwein/PISTUFF/Apistuff.html>.
5. R. Courant, *Differential and Integral Calculus*, Interscience, New York, NY, 1937.
6. H.-D. Ebbinghaus et al., *Numbers*, Springer-Verlag, New York, NY, 1990.
7. Leonard Euler, *An Introduction to the Analysis of the Infinite*, Springer-Verlag, New York, NY, 1990.
8. Anthony W. Knap, *Group Representations and Harmonic Analysis from Euler to Langlands, Part I*, Notices of the AMS 43 (1996), 410-415.
9. Serge Lang, *Undergraduate Analysis*, Springer-Verlag, New York, NY, 1983.
10. Mark A. Pinsky, *Introduction to Partial Differential Equations with Applications*, McGraw-Hill, New York, NY, 1984.
11. Robert S. Strichartz, *The Way of Analysis*, Jones & Bartlett, Boston, MA, 1995.
12. E. T. Whittaker and G. N. Watson, *Modern Analysis*, Cambridge University Press, London, UK, 1927.

Index

$B(x, y)$, 193
 $M(a, b)$, 182
 $O(g(x))$, 230
 $\Gamma(x)$, 163
 \approx , 230
 β , 207
 γ , 152
 π , 95
 $\psi(x)$, 168, 181, 212, 220
 $\sigma(n)$, 213
 \sim , 186, 218
 $\tau(x)$, 206
 θ , 211
 θ_+ , θ_- , θ_0 , 212
 ζ , 180
 $a \mid b$, 10
 n th root, 14
 $x!$, 164
 $n!$, 28
 \mathbb{N} , 7
 \mathbb{Q} , 9
 \mathbb{R} , 3
 \mathbb{Z} , 9

Absolute value, 13
AGM, 182
AGM curve, 214
algebraic order, 58
angle, 100
 \arccos , 97
Archimedes' constant, 95
 \arcsin , 97
 \arctan , 99
area, 115
 additivity, 123

 affine invariance, 126
 dilation invariance, 117
 monotonicity, 118
 naive, 116
 of a parallelogram, 122
 of a rectangle, 119
 of a triangle, 122
 reflection invariance, 118
 rotation invariance, 122
 subadditivity, 118
 translation invariance, 116
arithmetic progressions, 223
asymptotic
 equality, 186, 197, 218
 expansion, 230, 231

Bailey-Borwein-Plouffe series, 169
bell-shaped curve, 189

Bernoulli

 function, 206
 numbers, 206
 series, 206

Bessel function, 171, 181, 196

beta function, 193

binomial coefficient, 201

binomial theorem, 81

 Newton's generalization, 91

bound

 lower, 4
 upper, 4

Cantor set, 111

 area of the, 126

Cauchy order, 35

Cauchy-Schwarz inequality

Cauchy-Schwarz inequality (*cont.*)

- for integrals, 152
- in the plane, 101
- chain rule, 64
- codomain, 2
- compact, 39, 41
- complement, 2
- completeness property, 4
- continued fractions, 12, 25, 38, 169, 174
- continuity, 45
 - from the left, 50
 - from the right, 50
 - in two variables, 178
 - modulus of, 48
 - under the integral sign, 178
 - under the summation sign, 180
 - uniform, 51
 - uniform modulus of, 51
- convex conjugate, 81
- cosecant, 99
- cosh, 86
- cosine, 94
- cotangent, 99
- coth, 205
- cover, 115
- critical
 - point, 65
 - value, 65

De Morgan's law, 2

- decimal
 - expansions, 27
 - point, 28
- delta function, 143
- derivative, 61
 - partial, 189
- differentiation, 61
 - under the integral sign, 189
 - under the summation sign, 195
- digits, 27
- dilation, 6, 99, 117
 - centered, 117
- discontinuity
 - jump, 47
 - mild, 48
 - removable, 47
 - wild, 48
- distance, 100
- domain, 2

dominated convergence theorem

- for integrals, 170
- for series, 179
- domination, 170
- duplication formula, 201

Entire, 89

- entropy, 201
- euclidean motion, 99
- Euler product, 222
- Euler's constant, 152
- Euler-Maclaurin
 - derivative, 227
 - formula, 229
- even, 9, 67
 - part, 67
- expansion
 - base of, 28
 - binary, 28
 - decimal, 27
 - hexadecimal, 28
 - ternary, 28

Factorial

- of a natural, 28
- of a real, 164
- Fatou's lemma, 167
- Fourier transform, 181, 196
- fractional part, 10
- functional equation
 - AGM, 186
 - theta, 212
 - zeta, 221
- functions, 2
 - algebraic, 55
 - bounded, 127
 - concave, 75
 - continuous, 45
 - convex, 75
 - decreasing, 44
 - differentiable, 61
 - elementary, 229
 - exponential, 57
 - increasing, 44
 - logarithmic, 57
 - monotone, 44
 - negative part, 128
 - nonnegative, 127
 - positive part, 128

- power, 55, 56
- rational, 47
- signed, 128
- smooth, 74
- strictly concave, 75
- strictly convex, 75
- strictly decreasing, 44
- strictly increasing, 44
- strictly monotone, 44
- theta, 212
- transcendental, 56
- fundamental theorem, 145
- Gamma function, 163**
- Gaussian**
 - function, 189
 - integral, 188
- golden mean, 38
- greatest integer, 10
- Heat equation, 217**
- homogeneity, 182
- Inductive, 7**
 - hypothesis, 8
 - step, 8
- inf, 4
- infimum, 4
- infinite product, 203
 - for $\Gamma(x)$, 210
 - for \sin , 209
 - for \sinh , 203
- integrability
 - of nonnegative functions, 128
 - of signed functions, 129
- integral test, 140
- integrals
 - additivity, 134
 - continuity, 139
 - continuity at the endpoints, 137
 - differentiability, 144
 - dilation invariance, 132
 - integration by parts, 148
 - linearity, 147
 - monotonicity, 131
 - of nonnegative functions, 128
 - of signed functions, 129
 - substitution, 148
 - translation invariance, 132
- interior, 113
- intermediate value property
 - for continuous functions, 50
 - for derivatives, 74
- intersection, 2
- interval, 12
 - compact, 13
 - open, 13
 - punctured, 42
- inverse function theorem
 - for continuous functions, 54
 - for differentiable functions, 69
- irrationals, 14
- L'Hopital's rule, 73**
- Laplace transform, 172, 195
 - of τ , 211
- Laplace's theorem, 198
- limits
 - from the left, 43
 - from the right, 43
 - lower, 19
 - of functions, 42
 - of monotone sequences, 18
 - of sequences, 19
 - upper, 19
- log-convex, 167
- Machin's formula, 108**
- Maclaurin series, 85**
- mappings, 2
 - bijective, 2
 - composition of, 2
 - injective, 2
 - invertible, 3
 - surjective, 2
- maximum, 5
 - global, 65
 - local, 65
- maximum principle, 81
- mean
 - arithmetic, 182
 - arithmetic-geometric, 182
 - geometric, 182
- mean value theorem, 66
 - generalized, 72
- method of exhaustion, 110, 153
- minimum, 5
 - global, 65

- minimum (*cont.*)
 - local, 65
- monomial, 46
- monotone convergence theorem
 - for integrals, 164
 - for series, 166
- multiplicative, 224
- Numbers
 - algebraic, 58
 - integers, 9
 - natural, 7
 - rational, 9
 - real, 3
 - transcendental, 58
- Odd, 9, 67
 - part, 67
- Parity, 8
- partial sum, 25
- partition, 44
 - mesh of a , 52
- paving, 115
- Pi, 95
- piecewise
 - constant, 52, 136
 - continuous, 147
 - differentiable, 153
- polar coordinates, 98
- polynomial, 46
 - coefficients of a , 46
 - degree of a , 46
- polynomial growth, 199
- power series
 - differentiability of, 89
- primes, 10
- primitives, 102
 - integration by parts, 104
 - linearity, 104
 - substitution, 105
- Radius of convergence, 88
- range, 2
- ratio test, 88
- rectangle
 - compact, 40, 114
 - open, 40, 114
- reflection, 6
- relation, 2
- remainder, 83
 - Cauchy form, 84
 - integral form, 150
 - Lagrange form, 84
- Riemann integrable, 142
- Riemann sum, 137, 141
- root test, 88
- rotation, 99
- Sawtooth formula, 219
- secant, 99
- sequence, 16
- sequences
 - bounded, 19
 - Cauchy, 31
 - convergent, 23
 - decreasing, 17
 - error, 31
 - increasing, 17
 - lower, 18
 - monotone, 18
 - of sets, increasing, 154
 - sub-, 23
 - upper, 18
- series, 25
 - n th term test, 28
 - absolutely convergent, 31
 - alternating, 30
 - alternating version, 37
 - Cauchy product, 37
 - comparison test, 27
 - conditionally convergent, 31
 - convergent, 26
 - Dirichlet test, 32
 - double, 35
 - geometric, 26
 - harmonic, 28
 - Leibnitz, 108, 169
 - Leibnitz test, 33
 - power, 86
 - signed, 30
 - tail of a , 26
 - telescoping, 29
- set of differences, 162
- sets, 1
 - bounded, 4
 - closed, 161
 - countable, 35

- disjoint, 2
- empty, 2
- finite, 8
- infinite, 8
- interopen, 158
- measurable, 156
- open, 155
- product of, 2
- sub-, 1
- super-, 2
- sine, 94
- sinh, 86
- square root, 10
- Stirling
 - approximation, 197
 - series, 231
- subconverges, 23
- subgraph, 127
- summand, 25
- summation by parts, 32
- summation under the integral sign
 - absolute case, 171
 - alternating case, 171
 - positive case, 165
- sup, 4
- superlinear, 59, 82
- supremum, 4
- Tangency
 - of functions, 63
 - of lines, 64
- tangent, 98
- Taylor series, 85
- Taylor's theorem, 84, 150
- theta function, 211
- translation, 6, 99, 116
- triangle inequality, 13
- Union, 2
- unit circle, 97
- unit disk, 97
 - area of, 147
- Variation, 45
 - bounded, 45
 - total, 45
- Vieta formula, 169
- Wallis product, 169
- well-separated, 123
- Zeta function, 206, 217

Undergraduate Texts in Mathematics

(continued from page ii)

Lang: Calculus of Several Variables.
Third edition.

Lang: Introduction to Linear Algebra.
Second edition.

Lang: Linear Algebra.
Third edition.

Lang: Undergraduate Algebra. Second
edition.

Lang: Undergraduate Analysis.

Lax/Burstein/Lax: Calculus with
Applications and Computing.
Volume 1.

LeCuyer: College Mathematics with
APL.

Lidl/Pilz: Applied Abstract Algebra.

Macki-Strauss: Introduction to Optimal
Control Theory.

Malitz: Introduction to Mathematical
Logic.

Marsden/Weinstein: Calculus I, II, III.
Second edition.

Martin: The Foundations of Geometry
and the Non-Euclidean Plane.

Martin: Transformation Geometry: An
Introduction to Symmetry.

Millman/Parker: Geometry: A Metric
Approach with Models. Second
edition.

Moschovakis: Notes on Set Theory.

Owen: A First Course in the
Mathematical Foundations of
Thermodynamics.

Palka: An Introduction to Complex
Function Theory.

Pedrick: A First Course in Analysis.

Peressini/Sullivan/Uhl: The Mathematics
of Nonlinear Programming.

Prenowitz/Jantosciak: Join Geometries.

Priestley: Calculus: An Historical
Approach.

Protter/Morrey: A First Course in Real
Analysis. Second edition.

Protter/Morrey: Intermediate Calculus.
Second edition.

Roman: An Introduction to Coding and
Information Theory.

Ross: Elementary Analysis: The Theory
of Calculus.

Samuel: Projective Geometry.
Readings in Mathematics.

Scharlau/Opolka: From Fermat to
Minkowski.

Sethuraman: Rings, Fields, and Vector
Spaces: An Approach to Geometric
Constructability.

Sigler: Algebra.

Silverman/Tate: Rational Points on
Elliptic Curves.

Simmonds: A Brief on Tensor Analysis.
Second edition.

Singer/Thorpe: Lecture Notes on
Elementary Topology and Geometry.

Smith: Linear Algebra. Second edition.

Smith: Primer of Modern Analysis.
Second edition.

Stanton/White: Constructive
Combinatorics.

Stillwell: Elements of Algebra:
Geometry, Numbers, Equations.

Stillwell: Mathematics and Its History.

Strayer: Linear Programming and Its
Applications.

Thorpe: Elementary Topics in
Differential Geometry.

Toth: Glimpses of Algebra and
Geometry.

Troutman: Variational Calculus and
Optimal Control. Second edition.

Valenza: Linear Algebra: An
Introduction to Abstract Mathematics.

Whyburn/Duda: Dynamic Topology.

Wilson: Much Ado About Calculus.

Undergraduate Texts in Mathematics

(continued from page ii)

Lang: Calculus of Several Variables.
Third edition.

Lang: Introduction to Linear Algebra.
Second edition.

Lang: Linear Algebra.
Third edition.

Lang: Undergraduate Algebra. Second
edition.

Lang: Undergraduate Analysis.

Lax/Burstein/Lax: Calculus with
Applications and Computing.
Volume 1.

LeCuyer: College Mathematics with
APL.

Lidl/Pilz: Applied Abstract Algebra.

Macki-Strauss: Introduction to Optimal
Control Theory.

Malitz: Introduction to Mathematical
Logic.

Marsden/Weinstein: Calculus I, II, III.
Second edition.

Martin: The Foundations of Geometry
and the Non-Euclidean Plane.

Martin: Transformation Geometry: An
Introduction to Symmetry.

Millman/Parker: Geometry: A Metric
Approach with Models. Second
edition.

Moschovakis: Notes on Set Theory.

Owen: A First Course in the
Mathematical Foundations of
Thermodynamics.

Palka: An Introduction to Complex
Function Theory.

Pedrick: A First Course in Analysis.

Peressini/Sullivan/Uhl: The Mathematics
of Nonlinear Programming.

Prenowitz/Jantosciak: Join Geometries.

Priestley: Calculus: An Historical
Approach.

Protter/Morrey: A First Course in Real
Analysis. Second edition.

Protter/Morrey: Intermediate Calculus.
Second edition.

Roman: An Introduction to Coding and
Information Theory.

Ross: Elementary Analysis: The Theory
of Calculus.

Samuel: Projective Geometry.
Readings in Mathematics.

Scharlau/Opolka: From Fermat to
Minkowski.

Sethuraman: Rings, Fields, and Vector
Spaces: An Approach to Geometric
Constructability.

Sigler: Algebra.

Silverman/Tate: Rational Points on
Elliptic Curves.

Simmonds: A Brief on Tensor Analysis.
Second edition.

Singer/Thorpe: Lecture Notes on
Elementary Topology and Geometry.

Smith: Linear Algebra. Second edition.

Smith: Primer of Modern Analysis.
Second edition.

Stanton/White: Constructive
Combinatorics.

Stillwell: Elements of Algebra:
Geometry, Numbers, Equations.

Stillwell: Mathematics and Its History.

Strayer: Linear Programming and Its
Applications.

Thorpe: Elementary Topics in
Differential Geometry.

Toth: Glimpses of Algebra and
Geometry.

Troutman: Variational Calculus and
Optimal Control. Second edition.

Valenza: Linear Algebra: An
Introduction to Abstract Mathematics.

Whyburn/Duda: Dynamic Topology.

Wilson: Much Ado About Calculus.

This text is intended for an honors calculus course or for an introduction to analysis. Involving rigorous analysis, computational dexterity, and a breadth of application, it is ideal for undergraduate majors.

The book contains many remarkable features:

- complete avoidance of ϵ - δ arguments by instead using sequences;
- definition of the integral as the area under the graph, while area is defined for *every* subset of the plane;
- heavy emphasis on computational problems;
- applications from many parts of analysis, e.g., convexity, Cantor set, continued fractions, Bessel functions, the zeta function, and many more;
- 347 problems with solutions in the back of the book.