

Total No. of Printed Pages—7

**2 SEM TDC MTH M 1**

**2 0 1 8**

( May )

**MATHEMATICS**

( Major )

Course : 201

**( Matrices, Ordinary Differential Equations,  
Numerical Analysis )**

*Full Marks : 80*

*Pass Marks : 32/24*

*Time : 3 hours*

*The figures in the margin indicate full marks  
for the questions*

**GROUP—A**

**( Matrices )**

**( Marks : 20 )**

1. (a) If  $A$  is an  $n$ -rowed non-singular matrix,  
then what is the rank of  $A^T$ ? 1
- (b) Find the rank of the matrix

$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 0 & 1 & 1 \end{pmatrix}$$

2

( Turn Over )

( 2 )

- (c) Reduce the matrix  $A$  to its normal form where

$$A = \begin{pmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -4 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{pmatrix}$$

Hence find the rank of  $A$ .

Or

Find the rank of the matrix

$$\begin{pmatrix} 1 & 1 & -3 & 2 \\ 2 & -1 & 2 & -3 \\ 3 & -2 & 1 & -4 \\ -4 & 1 & -3 & 1 \end{pmatrix}$$

by reducing it to echelon form.

2. Answer any *two* of the following :  $6 \times 2 = 12$

- (a) Define characteristic roots of a matrix. Find the characteristic equation of the matrix

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 3 & 1 & 0 \\ -2 & 1 & 4 \end{pmatrix}$$

and verify Cayley-Hamilton theorem. Hence compute  $A^{-1}$ .

( 3 )

- (b) Find for what values of  $\lambda$ , the equations

$$\begin{aligned} x + y + z &= 1 \\ x + 2y + 4z &= \lambda \\ x + 4y + 10z &= \lambda^2 \end{aligned}$$

have a solution and also solve them completely in each case.

- (c) What do you mean by homogeneous and non-homogeneous linear equations? Show that the system of equations

$$\begin{aligned} 5x + 3y + 7z &= 4 \\ 3x + 26y + 2z &= 9 \\ 7x + 2y + 10z &= 9 \end{aligned}$$

is consistent and solve it.

GROUP—B

( Ordinary Differential Equations )

( Marks : 30 )

3. (a) Find Wronskian of  $\cos bx$  and  $\sin bx$  ( $b \neq 0$ ).

1

- (b) Solve :

2

$$(x + y + 1) \frac{dy}{dx} = 1$$

( 4 )

- (c) Find the complete solution and singular solution of the differential equation

$$y = px + f(p), \text{ where } p = \frac{dy}{dx} \quad 3$$

- (d) Answer any one of the following : 4

(i) Solve :

$$x dx + y dy + \frac{x dy - y dx}{x^2 + y^2} = 0$$

(ii) Prove that Wronskian of the functions  $e^{m_1 x}$ ,  $e^{m_2 x}$ ,  $e^{m_3 x}$  is equal to

$$(m_1 - m_2)(m_2 - m_3)(m_3 - m_1)e^{(m_1 + m_2 + m_3)x}$$

4. (a) Under what condition  $y = e^{ax}$  will be a solution of the equation

$$\frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Qy = 0 ? \quad 1$$

- (b) Show that the roots of the auxiliary equation are 1, 1, -2 of the differential equation

$$x^3 \frac{d^3 y}{dx^3} + 3x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2y = x^2 \quad 2$$

( 5 )

- (c) Solve any one of the following : 3

$$(i) \frac{d^2 y}{dx^2} + y = \cos 2x$$

$$(ii) (D^2 - 4D + 4)y = x^3 e^{2x}$$

- (d) Solve any one of the following : 4

$$(i) (x^2 D^2 - 3x D + 5)y = \sin(\log x) \text{ where } \frac{d}{dx} \equiv D$$

$$(ii) \sin^2 x \cdot \frac{d^2 y}{dx^2} = 2y, \text{ given } y = \cot x \text{ is a solution}$$

5. Answer any two of the following :  $5 \times 2 = 10$

- (a) Solve by removal of the first-order derivative :

$$\frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} + (x^2 + 1)y = x^3 + 3x$$

- (b) Solve by changing the independent variable :

$$\frac{d^2 y}{dx^2} + \cot x \frac{dy}{dx} + 4y \operatorname{cosec}^2 x = 0$$

- (c) Solve by the method of variation of parameters :

$$\frac{d^2 y}{dx^2} + 9y = \sec 3x$$

## GROUP—C

## ( Numerical Analysis )

( Marks : 30 )

6. (a) What is the degree of convergence of the Newton-Raphson method? 1
- (b) Give the geometrical interpretation of Newton-Raphson method. 4
- (c) Solve  $x^3 - 2x - 5 = 0$  for the positive root by iteration method. 5

Or

Solve the equation  $x \tan x + 1 = 0$  by regula falsi method starting with  $a = 2.5$  and  $b = 3$  correct to 3 decimal places.

- (d) Solve by Gauss elimination method : 5

$$\begin{aligned} 2x + 3y - z &= 5; & 4x + 4y - 3z &= 3; \\ & & 2x - 3y + 2z &= 2 \end{aligned}$$

Or

Apply Gauss-Jordan method to find the solution of the following system :

$$\begin{aligned} 10x + y + z &= 12; & 2x + 10y + z &= 13; \\ & & x + y + 5z &= 7 \end{aligned}$$

( Continued ) 8P—3200/501

7. (a) State 'true' or 'false' : 1  
Simpson's one-third rule is better than the trapezoidal rule.
- (b) Show that  $\delta \equiv E^{1/2} - E^{-1/2}$ , where the symbols have their usual meanings. 2
- (c) Evaluate : 2

$$\Delta^3(1-x)(1-2x)(1-3x) \text{ if } h=1$$

- (d) Answer any two of the following :  $5 \times 2 = 10$

(i) The population of a town is as follows :

Year	x	1941	1951	1961	1971	1981	1991
Population in Lakhs y		20	24	29	36	46	51

Estimate the population increase during the period 1946 to 1976.

- (ii) Evaluate

$$\int_0^1 \frac{dx}{1+x}$$

by dividing the range into 10 equal parts correct to four decimal places.

- (iii) Derive the Newton's forward interpolation formula.

- (iv) Deduce the general quadrature formula for equidistant ordinates.

\*\*\*