

Total No. of Printed Pages—8

2 SEM TDC MTH M 1

2019

(May)

MATHEMATICS

(Major)

Course : 201

**(Matrices, Ordinary Differential Equations,
Numerical Analysis)**

Full Marks : 80
Pass Marks : 32/24

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

GROUP—A

(Matrices)

(Marks : 20)

1. (a) Choose the correct option :

If a matrix A has a non-zero minor of order r , then

(i) $\text{rank}(A) = r$

(ii) $\text{rank}(A) \geq r$

(iii) $\text{rank}(A) < r$

(iv) $\text{rank}(A) \leq r$

(Turn Over)

(2)

- (b) For what value of x the rank of the matrix

$$A = \begin{bmatrix} 2 & 4 & 2 \\ 3 & 1 & 2 \\ 1 & 0 & x \end{bmatrix}$$

will be less than 3?

2

- (c) Reduce the matrix A to its normal form where

$$A = \begin{bmatrix} 2 & -2 & 0 & 6 \\ 4 & 2 & 0 & 2 \\ 1 & -1 & 0 & 3 \\ 1 & -2 & 1 & 2 \end{bmatrix}$$

Hence, find the rank of A .

5

Or

Reduce the following matrix into echelon form and find its rank :

$$\begin{bmatrix} 1 & 3 & 1 & -2 & -3 \\ 1 & 4 & 3 & -1 & -4 \\ 2 & 3 & -4 & -7 & -3 \\ 3 & 8 & 1 & -7 & -8 \end{bmatrix}$$

2. (a) Under what condition a system of m homogenous linear equations $AX=0$ in n unknowns will possess infinite number of solutions?

1

24

(3)

- (b) For what value of k the system of equations

$$\begin{aligned} x + 5y - 3z &= -4 \\ -x - 4y + z &= 3 \\ -2x - 7y &= k \end{aligned}$$

is consistent? Solve it.

5

- (c) Show that the matrix

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

satisfies Cayley-Hamilton theorem.

Hence, compute A^{-1} .

4+2=6

Or

What is the degree of characteristic polynomial of an $n \times n$ square matrix? Determine the characteristic roots and characteristic vectors of the matrix

$$A = \begin{bmatrix} 3 & 5 \\ -2 & -4 \end{bmatrix}$$

1+5=6

GROUP—B

(Ordinary Differential Equations)

(Marks : 30)

3. (a) Write the general solution of the differential equation

$$\frac{d^3 y}{dx^3} = 0$$

if 1, x , x^2 are its linearly independent solutions.

- (b) Solve :

$$(1+x^2)\frac{dy}{dx} + y = e^{\tan^{-1} x}$$

- (c) Find the general solution of the differential equation $p = \tan(px - y)$, where

$$p = \frac{dy}{dx}$$

- (d) Answer any one of the following :

- (i) Evaluate Wronskian of the functions e^x and xe^x . Hence, conclude whether or not they are linearly independent. If they are independent, set up the differential equation having them as its independent solutions.

- (ii) Solve :

$$(x^2 + y^2 + x)dx - (2x^2 + 2y^2 - y)dy = 0$$

4. (a) Under what condition $y = x$ is a part of the complementary function of the differential equation

$$\frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Qy = R ?$$

- (b) Find the particular integral of the differential equation

$$\frac{d^2 y}{dx^2} + \frac{dy}{dx} - 6y = x$$

- (c) Answer any one of the following :

- (i) Solve :

$$\frac{d^2 y}{dx^2} + 4y = x \cos x$$

- (ii) Solve :

$$\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 3y = 2e^{3x}$$

(d) Answer any one of the following : 4

(i) Solve :

$$(x^2 D^2 + xD + 1)y = \sin \log x^2,$$

where $D \equiv \frac{d}{dx}$

(ii) Solve

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - 9y = 0$$

given that $y = x^3$ is a solution.

5. Answer any two of the following : 5×2=10

(a) Solve by removal of the first-order derivative :

$$\frac{d^2 y}{dx^2} - 2 \tan x \frac{dy}{dx} + 5y = 0$$

(b) Solve by changing the independent variable :

$$x \frac{d^2 y}{dx^2} + (4x^2 - 1) \frac{dy}{dx} + 4x^3 y = 2x^3$$

(c) Solve by the method of variation of parameters :

$$\frac{d^2 y}{dx^2} + a^2 y = \operatorname{cosec} ax$$

where a is a constant.

GROUP—C

(Numerical Analysis)

(Marks : 30)

6. (a) State True or False : 1
Iteration method is always convergent.

(b) Evaluate $\sqrt{12}$ using Newton-Raphson method by performing two iterations. 4

Or

Describe Newton-Raphson method for finding real roots of an algebraic equation.

(c) Find the real root of the equation $x^3 - x - 1 = 0$ lying between 1 and 2 using bisection method by performing three iterations. 5

Or

Find a real root of the equation $x^3 - 2x - 5 = 0$ using regula-falsi method by performing three iterations.

(d) Solve by Gauss elimination method : 5

$$2x + 2y + 4z = 14$$

$$3x - y + 2z = 13$$

$$5x + 2y - 2z = 2$$

(Turn Over)

Or

Solve by Gauss-Seidel method by performing two iterations :

$$5x + 2y + z = 12$$

$$x + 4y + 2z = 15$$

$$x + 2y + 5z = 20$$

7. (a) What is the degree of the interpolating polynomial in Simpson's $\frac{3}{8}$ rule? 1
- (b) Show that $(1 + \Delta)(1 - \nabla) = 1$, where the symbols have their usual meanings. 2
- (c) If $f(x) = \frac{1}{x^2}$, find the divided difference $f(a, b)$. 2
- (d) Answer any two of the following questions : $5 \times 2 = 10$

(i) Derive Newton's forward interpolation formula.

(ii) The population of a town is as follows :

Year	x	: 1891	1901	1911	1921	1931
Population in lakh	y	: 46	66	81	93	101

Estimate the population for the year 1925.

- (iii) Deduce Simpson's $\frac{1}{3}$ rule for numerical integration.
- (iv) Find the form of the function given by

x	:	1	2	5
$f(x)$:	1	4	10

★ ★ ★