

Total No. of Printed Pages—11

4 SEM TDC CHM M 1 (N/O)


2016

(May)

CHEMISTRY

(Major)

Course : 401



(Physical Chemistry—I)

*The figures in the margin indicate full marks
for the questions*

(New Course)

Full Marks : 48

Pass Marks : 14

Time : 2 hours

1. Select the correct answer :

1×5=5

(a) Li^+ has a smaller ionic mobility than K^+ because of the

- (i) larger size of Li^+
- (ii) larger radius to charge ratio of Li^+
- (iii) greater degree of hydration of Li^+
- (iv) smaller nuclear charge of Li^+

(2)

(b) Which of the following is not a good conductor of electricity?

- (i) NaCl (aq)
- (ii) NaCl (s)
- (iii) NaCl (molten)
- (iv) Silver metal

(c) The primary cells are

- (i) rechargeable
- (ii) not rechargeable
- (iii) everlasting
- (iv) None of the above

(d) Standard electrode potentials of three metals X, Y and Z are -1.2 V, $+0.5$ V and -3.0 V respectively. The reducing power of these metals will be

- (i) $Z > X > Y$
- (ii) $X > Y > Z$
- (iii) $Y > Z > X$
- (iv) $Y > X > Z$

(e) 2 moles of an ideal gas at 27°C temperature are expanded reversibly from 2 L to 20 L. If the value of R is taken as $2 \text{ cal K}^{-1} \text{ mol}^{-1}$, then the entropy change will be

- (i) 92.1
- (ii) 0
- (iii) 4
- (iv) 9.2

(3)

2. Answer any five of the following questions :

2×5=10

- (a) Equivalent conductance of an electrolyte at finite concentration is less than that of an infinite dilution. Explain.
- (b) Explain why lithium ions move slower than potassium ions in water under an electric field.
- (c) Discuss how the quinhydrone electrode can be used to determine the pH of a solution.
- (d) In conductometric titration, the titre should be always very much concentrated than the solution to be titrated. Explain why.
- (e) Give one example each of electrode concentration cell and electrolyte concentration cell.
- (f) Write the physical significance of Helmholtz free energy and Gibbs free energy.
- (g) Calculate the entropy increase in the evaporation of a mole of water at 100°C .
(Heat of vaporization = 540 cal g^{-1})

(4)

UNIT—I

3. Answer any *two* of the following questions :


4½×2=

(a) Deduce an expression for the entropy changes associated with the changes in temperature and pressure of an ideal gas.

(b) (i) State Carnot theorem.

(ii) A Carnot engine works between the temperatures 27 °C and 127 °C. Calculate the efficiency of the engine.

(iii) Predict whether at 27 °C, the following reaction is spontaneous or not :



Given $\Delta H = +9080 \text{ J mol}^{-1}$ and

$\Delta S = +35.7 \text{ J K}^{-1} \text{ mol}^{-1}$.

(c) (i) State and explain Nernst heat theorem.

(ii) Describe the third law of thermodynamics.

(5)

UNIT—II

4. Answer any *two* of the following questions :

7×2=14

(a) (i) Represent the variation of equivalent conductance of KCl and CH_3COOH with dilution graphically and give explanation for such variation.

5

(ii) Define the term transference number.

2

(b) Explain Wien effect and Debye-Falkenhagen effect.

3½+3½=7

(c) (i) Define molar conductivity and equivalent conductivity.

4

(ii) The equivalent conductances at infinite dilution (λ_0) of HCl, CH_3COONa and NaCl are

426.16, 91.0 and 126.45 $\text{ohm}^{-1} \text{cm}^2 \text{g eqvt}^{-1}$

respectively. Calculate λ_0 of acetic acid.

3

UNIT—III

5. Answer any *two* of the following questions :

5×2=10

(a) What is liquid junction potential? Derive an expression for it.

5

(b) What are fuel cells? Discuss how the e.m.f. is generated in a hydrogen-oxygen fuel cell.

(c) (i) A solution of CuSO_4 is electrolysed for 10 minutes with a current of 1.5 amperes. What is the mass of copper deposited at the cathode?
(Atomic mass of copper = 63.56 u)

(ii) Define standard electrode potential.