### 5 SEM TDC MTH M 2

#### 2016

(November)

### MATHEMATICS

(Major)

Course: 502

# ( Linear Algebra and Number Theory )

Full Marks: 80

Pass Marks: 32 (Backlog)/24 (2014 onwards)

Time: 3 hours

The figures in the margin indicate full marks for the questions

#### GROUP-A

# ( Linear Algebra )

( Marks: 40 )

- (a) Write which of the following statements is 'true' and which is 'false': 1×2=2
  - (i) "The set containing a linearly independent set of vectors is itself linearly independent."
  - (ii) "Intersection of two subspaces of a vector space V is always a subspace of V."

(Turn Over)

3

4

- (b) Examine whether the vector (2, -5, 3)is in the subspace of  $\mathbb{R}^3$  spanned by the vectors (1, -3, 2), (2, -4, -1) and (1, -5, 7).
- Show that the set (c)

 $S = \{(1, 0), (i, 0), (0, 1), (0, i)\}$ forms a basis for the vector space V of ordered pairs of complex numbers over the field of real numbers R, i.e.,  $V = \mathbb{C}^2(\mathbb{R})$ .

- (d) Let V be a finite dimensional vector space of dimension n. Then prove that any set of n linearly independent vectors in V forms a basis for V.
- Let V be any vector space. Prove that the set  $\{v_1, v_2, ..., v_n\}$  is linearly dependent if and only if one of the  $v_i$ 's is a linear combination of the other  $v_j$ 's where
- Define subspace of a vector space. Prove that the set W defined as

 $W = \{(a, b, 0) : a, b \in \mathbb{R}\}\$ 

is a subspace of  $\mathbb{R}^3$ . P7/180

- 2. (a) Let l(p; d) and l(q; d) be two lines passing through p and q respectively Show d. direction l(p; d) = l(q; d) if and only if (q - p) is a having multiple of d.
  - Let T be a linear transformation from a vector space U to a vector space V over the field F. Prove that the range of T is a subspace of V.
  - Show that a linear map T from a vector space to another is one-one if and only (c) if ker  $T = \{0\}$ .
  - (d) Let V be the vector space of all polynomials in x with coefficients in  $\mathbb{R}$ of the form

the form
$$f(x) = a_0 x^0 + a_1 x^1 + a_2 x^2 + a_3 x^3$$
appearator *D* is a lim

The differentiation operator D is a linear transformation on V. Write the matrix of D relative to the ordered basis

ative to the order 
$$B = \{x^0, x^1, x^2, x^3\}$$

Show that the mapping  $T: \mathbb{R}^2 \to \mathbb{R}^3$ defined as T(a, b) = (a+b, a-b, b) is a linear transformation from  $\mathbb{R}^2$  into  $\mathbb{R}^3$ . Find the rank and nullity of T. 2+2+2=6

1

4

3

## GROUP-B

# ( Number Theory )

( Marks: 40 )

- 3. When are two integers said to be relatively prime?
- 4. Answer any two from the following: 3×2=1 Use division algorithm to establish that the square of any integer is either of the form 3k or 3k+1.
  - Prove that if  $a \mid bc$  with gcd(a, b) = 1,
  - Use Euclidean algorithm to obtain (c) integers x and y satisfying the following:

$$\gcd(56, 72) = 56x + 72y$$

- Show that if p is a prime and p|ab then either p|a or p|b.
  - Prove that given any positive integer n, there exist n consecutive composite
  - (c) Find the highest power of 5 dividing 100!.

6. (a) Write a complete set of residues modulo 7.

If  $a \equiv b \pmod{n}$  and the integers a, b, n are all divisible by d > 0, then prove that

$$\frac{a}{d} \equiv \frac{b}{d} \left( \bmod \frac{n}{d} \right)$$

If a is an odd integer, then prove that

$$a^2 \equiv 1 \pmod{8}$$

Solve 18x + 5y = 48.

Solve the following by using Chinese remainder theorem:

$$x \equiv 5 \pmod{4}$$
$$x \equiv 3 \pmod{7}$$

$$x \equiv 2 \pmod{9}$$

7. (a) Evaluate

(i)  $\sigma(210)$ 

(ii) d(63)

where the symbols have their usual

When is an arithmetic function said to be multiplicative? Prove that o is a (b) multiplicative function.

P7/180

P7\_3500/180

1 Continued

5 SEM TDC MTH M 2